

Relationships of Polychlorinated Biphenyls and Dichlorodiphenyldichloroethylene (*p,p'*-DDE) with Testosterone Levels in Adolescent Males

Lawrence M. Schell,^{1,2,3} Mia V. Gallo,^{1,2} Glenn D. Deane,^{1,4} Kyrie R. Nelder,² Anthony P. DeCaprio,⁵ Agnes Jacobs,⁶ and the Akwesasne Task Force on the Environment⁶

¹Center for the Elimination of Minority Health Disparities, and ²Department of Anthropology, University at Albany, Albany, New York, USA; ³Department of Epidemiology and Biostatistics, University at Albany, School of Public Health, Rensselaer, New York, USA; ⁴Department of Sociology, University at Albany, Albany, New York, USA; ⁵Department of Chemistry and Biochemistry, Florida International University, Miami, Florida, USA; ⁶Akwesasne Mohawk Nation, Akwesasne, New York, USA

BACKGROUND: Concern persists over endocrine-disrupting effects of persistent organic pollutants (POPs) on human growth and sexual maturation. Potential effects of toxicant exposures on testosterone levels during puberty are not well characterized.

OBJECTIVES: In this study we evaluated the relationship between toxicants [polychlorinated biphenyls (PCBs), dichlorodiphenyldichloroethylene (*p,p'*-DDE), hexachlorobenzene (HCB), and lead] and testosterone levels among 127 Akwesasne Mohawk males 10 to < 17 years of age with documented toxicant exposures.

METHODS: Data were collected between February 1996 and January 2000. Fasting blood specimens were collected before breakfast by trained Akwesasne Mohawk staff. Multivariable regression models were used to estimate associations between toxicants and serum testosterone, adjusted for other toxicants, Tanner stage, and potential confounders.

RESULTS: The sum of 16 PCB congeners ($\Sigma 16\text{PCBs}$) that were detected in $\geq 50\%$ of the population was significantly and negatively associated with serum testosterone levels, such that a 10% change in exposure was associated with a 5.6% decrease in testosterone (95% CI: $-10.8, -0.5\%$). Of the 16 congeners, the more persistent ones ($\Sigma 8\text{PerPCBs}$) were related to testosterone, whereas the less persistent ones, possibly reflecting more recent exposure, were not. When PCB congeners were subgrouped, the association was significant for the sum of eight more persistent PCBs (5.7% decrease; 95% CI: $-11, -0.4\%$), and stronger than the sum of six less persistent congeners (3.1% decrease; 95% CI: $-7.2, 0.9\%$). *p,p'*-DDE was positively but not significantly associated with serum testosterone (5.2% increase with a 10% increase in exposure; 95% CI: $-0.5, 10.9\%$). Neither lead nor HCB was significantly associated with testosterone levels.

CONCLUSIONS: Exposure to PCBs, particularly the more highly persistent congeners, may negatively influence testosterone levels among adolescent males. The positive relationship between *p,p'*-DDE and testosterone indicates that not all POPs act similarly.

CITATION: Schell LM, Gallo MV, Deane GD, Nelder KR, DeCaprio AP, Jacobs A; Akwesasne Task Force on the Environment. 2014. Relationships of polychlorinated biphenyls and dichlorodiphenyldichloroethylene (*p,p'*-DDE) with testosterone levels in adolescent males. *Environ Health Perspect* 122:304–309; <http://dx.doi.org/10.1289/ehp.1205984>

Introduction

During adolescence the reproductive system matures rapidly, driven by changes in the gonadal steroid hormones testosterone and estradiol (Forest et al. 1976). A disturbance in this system during this critical transitional stage may have long-term repercussions because physiologic parameters reached in adolescence are related to endocrine characteristics in adulthood (Root 1973; Roy et al. 2009; Teilmann et al. 2002).

The possibility that exposure to human-made materials could disrupt this important process is of concern. Polychlorinated biphenyls (PCBs) and dichlorodiphenyltrichloroethane (DDT) are lipophilic organochlorines that bioaccumulate in adipose tissue and fat-rich fluids (Carpenter 1998; Wolff and Anderson 1999). PCBs and DDT [or its metabolite dichlorodiphenyldichloroethylene (*p,p'*-DDE)] have been associated with differences in measures of sexual maturation in

girls in some studies (Denham et al. 2005; Den Hond et al. 2011; Ouyang et al. 2005; Ozen et al. 2012; Su et al. 2012; Vasiliu et al. 2004; Yang et al. 2005), though not all studies (Gladen et al. 2000; Wolff et al. 2008). Such differences may reflect altered steroid hormone levels and associated functional changes (Bourguignon and Parent 2012). PCBs have been shown to inhibit androgen production *in vitro* (in rats) and *in vivo* (Andric et al. 2000; Kovacevic et al. 1995) possibly mediated by the aryl hydrocarbon receptor (AhR), although some PCB congeners that do not bind to the AhR also have some toxic properties (Fischer et al. 1998; Oskam et al. 2003; Yeowell et al. 1987). In addition to the organic compounds, lead is associated with later sexual maturation in girls (Denham et al. 2005; Kim et al. 1995; Selevan et al. 2003).

Less research has dealt with endocrine disruption of male sexual development by metals or organic compounds. In adults, high

levels of lead can interfere with reproduction (Alexander et al. 1996, 1998; Benoff et al. 2003a, 2003b). Some PCB congeners and congener groups were negatively associated with testosterone in a population of Akwesasne Mohawk men with high PCB exposure (Goncharov et al. 2009) and in other adult male populations (Abaci et al. 2009; Dhooge et al. 2006; Pflieger-Bruss et al. 2004). PCBs also have been negatively associated with adult male sperm counts and semen quality (Hauser 2006; Hauser et al. 2003; Mocarelli et al. 2008; Rignell-Hydbom et al. 2004; Toft et al. 2006). These relationships suggest that some persistent organic compounds could affect steroid hormones during adolescent sexual maturation.

Studies of boys' sexual maturation and its regulating hormones during adolescence have produced suggestive but inconsistent evidence of negative relationships with PCBs and *p,p'*-DDE (Den Hond et al. 2002; Mol et al. 2002; Pflieger-Bruss et al. 2004). The

Address correspondence to L.M. Schell, A&S 237, University at Albany, 1400 Washington Ave., Albany, NY 12222 USA. Telephone: (518) 442-4714. E-mail: lmschell@albany.edu

Supplemental Material is available online (<http://dx.doi.org/10.1289/ehp.1205984>).

We acknowledge and thank the Akwesasne Mohawk community, particularly M. Cole, A. Tarbell, D. David, P. Worswick, K. Jock, and C. Arquette for their many contributions, cooperation, and participation in the research we have done together on this topic. In addition, we thank R. Rej for his assistance.

This work was supported by grants from the National Institute of Environmental Health Sciences (ESO4913, ES10904), and the National Institute on Minority Health and Health Disparities, National Institutes of Health (1 P20 MD003373).

The content is solely the responsibility of the authors and does not represent the official views of the National Institute on Minority Health and Health Disparities or the National Institutes of Health.

Members of the Akwesasne community have been involved in a lawsuit against a local industry that has contributed to the pollution of the St. Lawrence River and/or its tributaries. Some of these members are related to the authors who are members of the community (A.J. and the Akwesasne Task Force on the Environment).

The authors declare they have no actual or potential competing financial interests.

Received: 6 September 2012; Accepted: 20 December 2013; Advance Publication: 20 December 2013; Final Publication: 1 March 2014.

elucidation of effects from exposure to these organics is complicated by the presence of lead that can delay boys' sexual maturation (Humblet et al. 2011; Korrick et al. 2011; Williams et al. 2010).

To evaluate the role of specific toxicants on aspects of sexual maturation, an approach that models the various exposures suspected of influencing maturation is necessary. The Akwesasne Mohawk Nation is a community with multiple toxicant exposures [PCBs, *p,p'*-DDE, hexachlorobenzene (HCB), lead] and PCB exposures that are substantially greater than those in the U.S. population as a whole (DeCaprio et al. 2005; Gallo et al. 2011; Schell et al. 2003). The aim of the present research is to estimate the relationship of testosterone levels with concurrent levels of PCBs, *p,p'*-DDE, HCB, and lead among male youth with known toxicant exposure while controlling for maturational stage.

Methods

Setting. The study was conducted in partnership with the Akwesasne Mohawk Nation, a sovereign territory that spans the St. Lawrence River, abutting New York State, USA, and Ontario and Quebec, Canada. Industrialization on the St. Lawrence River and some tributaries in the 1950s produced significant contamination. One federal Superfund site and two New York State Superfund sites are proximate to the Akwesasne Nation territory. The U.S. Food and Drug Administration determined that several local species of fish, fowl, and game contained levels of environmental contaminants above levels safe for human consumption (Fitzgerald et al. 1995; Forti et al. 1995; Lacetti 1993; Sloan and Jock 1990). The Akwesasne population's history of relying on local food sources, which carry the heaviest toxicant burden, suggested that this is a highly exposed population, and this has been confirmed (Schell et al. 2012).

The methods of recruitment, data collection, laboratory analyses, and substitution protocol for toxicant measurements below the limits of detection (LODs) have previously been described in detail (Schell et al. 2003). Members of the Akwesasne Mohawk Nation were trained in data collection techniques by the investigators to collect all anthropometric, interview, and hematological data, but had no prior knowledge of individual's exposure status. The Institutional Review Board of the University at Albany, State University of New York, approved the study protocols, and interviewers obtained informed assent from each participant and informed consent from the parent/guardian.

Participants. Participants were Akwesasne Mohawk youth (10 to < 17 years of age) and their mothers/guardians who lived either on or within 10 miles of the St. Regis Reservation or

Reserve between 1996 and 2000. A description of the sampling protocol has been published [see Schell et al. (2003) for details]. The original study sample consisted of 131 males and 140 females and their mothers/guardians. Four males had insufficient serum for testosterone and biomarker assays and were excluded, five were missing lead values, and two were missing triglycerides and cholesterol levels; the final sample was 120 males in the present analysis.

Blood collection and laboratory analysis. Fasting blood specimens were collected before breakfast (0700–1030 hours) by trained Akwesasne Mohawk staff. Testosterone level and the time of blood collection were not related ($r = 0.08$, $p = 0.38$). Analyses of the PCBs and organochlorines were conducted by parallel dual-column (splitless injection) gas chromatography with electron capture detection on an Agilent 6890 instrument (Agilent Technologies, Santa Clara, CA) at the University at Albany's Exposure Assessment Laboratory; the instrument was capable of capturing 83 separate PCB congeners and 18 additional congeners as pairs or triplets, as well as HCB and *p,p'*-DDE (DeCaprio et al. 2000, 2005). Data were expressed on a whole weight basis (i.e., nanograms per gram serum; parts per billion). Blood lead levels were measured using Zeeman-corrected graphite furnace atomic absorption spectrometry (LOD, 1.0 $\mu\text{g}/\text{dL}$; Mercury Monitor Model 100, Pharmacia Corp., Stockholm, Sweden) by Le Centre de Toxicologie du Quebec in Sainte-Foy Quebec, Quebec, Canada. Serum testosterone, estradiol, cholesterol, and triglyceride concentrations were measured by the Clinical Chemistry and Hematology Laboratory, Wadsworth Center for Laboratories and Research, New York State Department of Health (Albany, NY), as described by Schell et al. (2003).

Testosterone. Total testosterone was measured in unextracted serum specimens. A specific rabbit antibody affixed to polypropylene tubes (Siemens Diagnostics/Diagnostic Products, Los Angeles, CA) was used in the solid-phase radioimmunoassay procedure. A tracer, ^{125}I -labeled testosterone, and a Wallace 1470 Wizard gamma counter (Wallace/PerkinElmer, Waltham, MA) were used to measure radioactivity bound to the tracer. Instrument-based software calculated logit-log transformations, standard curves, and results. The average of duplicate measurements was reported and used. Differences in the duplicate measurements that exceeded 25% (or if the concentration was < 100 ng/dL, then by 25 ng/dL), was the rejection criterion causing re-assay of such specimens. Among samples > 100 ng/dL the variation on duplicate samples was 4.8%, and for samples < 100 ng/dL the variation between duplicates was 7.7%. The functional

sensitivity [limit of quantitation (LOQ)] was 10 ng/dL for testosterone. For statistical analyses all results < 10 ng/dL were set to 5 ng/dL, one-half of the LOQ.

PCBs and organochlorines. In some participants the levels of some PCBs congeners were below the laboratory LOD or minimum detection limit (MDL). Values below the MDL were imputed for each observation by sampling values from the distribution below the MDL, which was estimated based on the observed data using the method described by Gupta (1952), as recommended by the U.S. Environmental Protection Agency (EPA) for distributions where $\geq 50\%$ of the samples have values above the MDL (U.S. EPA 1998). This method provides a more appropriate representation of the sample variance for values below the MDL than do methods that impute a single value for all samples below the MDL. All values of *p,p'*-DDE and HCB were above the MDL.

Several composite exposure variables were evaluated to allow comparison of our results with those from other studies (following Schell et al. 2003): $\Sigma 16\text{PCBs}$ is the sum of all congeners with levels > MDL in $\geq 50\%$ of the study population samples (PCB congeners 52, 70, 74, 84, 87, 95, 99, 101[+90], 105, 110, 118, 149[+123], 138[+163 +164], 153, 180, and 187); $\Sigma 8\text{PerPCBs}$, a persistent PCB group (congeners 74, 99, 105, 118, 138[+163+164], 153, 180, 187); and $\Sigma 6\text{NonPerPCBs}$, a nonpersistent group (congeners 52, 84, 95, 101[+90], 110, and 149[+123]). Brackets indicate "minor" co-eluting congeners based on Aroclor concentration (Hansen 1999). We also estimated exposures with estrogenic PCB congeners ($\Sigma 7\text{EstrogenicPCBs}$: congeners 52, 70, 95, 99, 101, 110, and 153), and with PCB-105, an antiestrogenic congener (Cooke et al. 2001; Wolff et al. 1997).

Three composite variables used in a study of adult Akwesasne males were calculated (Goncharov et al. 2009). Using only congeners detected in $\geq 50\%$ of the sample, we calculated mono-*ortho* ($\Sigma 4\text{MOPCBs}$; congeners 118, 105, 74, and 70), di-*ortho* ($\Sigma 8\text{DOPCBs}$; congeners 52, 87, 99, 110, 153, 180, 138[+163+164], and 90[+101]), and tri- and tetra-*ortho* groups ($\Sigma 3\text{TTOPCBs}$; congeners 95, 187, and 84).

Anthropometric and sociodemographic variables. Each mother–youth pair completed interviews to report their sociodemographic characteristics, child's usual diet, whether the child was breastfed, and the mother's reproductive history (Schell et al. 2003, 2008). Trained staff measured each child's height and weight following standard anthropometric protocols (Gallo et al. 2005; Lohman et al. 1988). Height and weight *z*-scores were computed using EpiInfo 2000 [Centers

for Disease Control and Prevention (CDC) 2001]. Tanner stages (TS) were determined by self-assessment in a private room using drawings employed in a previously validated procedure (Duke et al. 1980; Morris and Udry 1980). Drawings were chosen to provide clearer distinctions between stages and were more acceptable to the community than photographs of the different stages.

Statistical analysis. Multivariable linear regression models were used to estimate associations of natural log (ln)-transformed serum testosterone levels with ln-transformed PCB groupings, controlling for other toxicants (*p,p'*-DDE, HCB, lead) and relevant covariates. We included serum HCB concentrations and blood lead levels (both as ln-transformed continuous variables) and whether the child was breastfed as an infant (1 = yes, 0 = no) in all models based on previous research suggesting associations between these factors and serum testosterone (Denham et al. 2005; Goncharov et al. 2009; Schell et al. 2003). In addition, we included the following factors in all models based on correlations with testosterone ($p < 0.20$): TS (categorical as described below), weight-for-age *z*-score and height-for-age *z*-score (both as continuous variables), child cigarette use (1 = yes, 0 = no), child alcohol use (1 = yes, 0 = no), and ln-transformed serum *p,p'*-DDE, cholesterol, and triglycerides (modeled as continuous variables).

The analytical treatment of TS is complicated by its level of measurement (TS is ordinal rather than interval level) and its expected association with age-related covariates. TSs were grouped based on preliminary analysis of variance (ANOVA) models, which indicated

that mean testosterone levels were comparable and not significantly different between TS I and II or between TS IV and V, but were significantly different between TS I and II (combined), TS III, and TS IV and V (combined). Therefore, we modeled TS using two indicator variables (for TS I and II, and for TS IV and V) with TS III as the referent category. Standardized measures of weight-for-age and height-for-age *z*-scores reduced age-related confounding.

The functional form of the dose response between ln-transformed PCBs and ln-transformed testosterone was assessed by fitting each exposure as a quadratic polynomial (i.e., by modeling ln-PCBs and a squared term) and using the *p*-value of the squared term as a test for statistically significant departures from linearity (data not shown). All *p*-values were > 0.05 ; therefore, exposures were modeled as continuous (ln-transformed) variables. Partial-regression plots (i.e., added-variable plots) and partial-residual plots were also examined for evidence of nonlinearity as well as outliers and influence (Fox 2008). Linearity in the dose response was also supported in these plots (data not shown). Nonadditivity (interactions evaluated as product terms and interpreted based on product term *p*-values: We tested all PCBs groups and HCB, *p,p'*-DDE, lead, and breastfeeding history because relatively few studies have examined the concurrent effects of the most common pollutants to which children may be exposed.

To facilitate the interpretation of the results, coefficients were reported in the regression tables, with the coefficients interpreted in the corresponding text. Results are reported as the percent difference in serum

testosterone associated with a 10% increase in exposure, which, for ln-transformed exposures, is derived by multiplying the coefficient for the association with ln-transformed testosterone by 10. Statistical analyses were conducted using PASW 19 (IBM 2011). An alpha level of $p \leq 0.05$ was used to define statistical significance.

Results

Sample descriptive measures of testosterone, toxicants, covariates, and their patterns of association with TS are shown in Table 1. As expected, mean age consistently increased with increasing TS [$F = 44.62$, df (degrees of freedom) = 122, 4, $p \leq 0.001$]. Mean testosterone levels also increased with TS ($F = 59.94$, $df = 122, 4, p \leq 0.001$), with similar mean values in boys classified as TS I or TS II, and as TS IV or TS V. There were 23 boys in TS I and II with testosterone values below the LOQ of 10 ng/dL, and two boys in TS III were below the LOQ. Height increased monotonically with TS ($F = 32.30$, $df = 122, 4, p \leq 0.001$). Mean weights also differed significantly over TSs ($F = 7.85$, $df = 122, 4, p \leq 0.001$), yet in contrast with height, this association was not monotonic. Forty-eight percent of the males were breastfed as infants, 9% currently smoked, and 6.7% consumed alcohol.

Mean serum PCB, *p,p'*-DDE, HCB, and lead concentrations were similar to values previously reported for the combined sample of Akwesasne male and female adolescents (Schell et al. 2003). The geometric mean concentration of $\Sigma 8$ PerPCBs (0.41 ppb) measured in 1996–2000 in our study population of 10- to < 17-year-old Akwesasne

Table 1. Characteristics of adolescent Akwesasne males: mean \pm SD for all observations combined ($n = 120$) and by Tanner stage.

Covariates and toxicants ^a	All	TS I ($n = 19$)	TS II ($n = 42$)	TS III ($n = 23$)	TS IV ($n = 32$)	TS V ($n = 11$)
Age (years)	13.17 \pm 1.95	11.32 \pm 1.09	11.93 \pm 1.43	13.31 \pm 1.51	14.62 \pm 1.09	16.04 \pm 0.81
Testosterone (ng/dL) ^b	252.18 \pm 279.60	26.41 \pm 35.31	48.71 \pm 90.71	264.70 \pm 252.28	509.88 \pm 203.42	555.70 \pm 299.24
Height (cm)	157.97 \pm 13.39	148.83 \pm 8.27	149.79 \pm 10.61	156.53 \pm 10.45	167.46 \pm 9.10	177.56 \pm 6.35
Height-for-age <i>z</i> -score	0.12 \pm 1.20	0.50 \pm 0.93	-0.01 \pm 1.29	-0.03 \pm 0.91	-0.04 \pm 1.41	0.83 \pm 0.82
Weight (kg)	62.59 \pm 19.04	61.56 \pm 20.95	53.37 \pm 18.75	62.22 \pm 14.54	68.16 \pm 16.68	82.39 \pm 13.51
Weight-for-age <i>z</i> -score	1.48 \pm 1.63	2.73 \pm 2.17	1.26 \pm 1.71	1.37 \pm 1.13	1.13 \pm 1.41	1.59 \pm 1.08
Triglycerides (mg/dL)	84.65 \pm 45.82	103.29 \pm 56.19	100.71 \pm 51.79	74.17 \pm 42.23	68.38 \pm 27.56	68.10 \pm 29.49
Cholesterol (mg/dL)	158.68 \pm 32.80	163.12 \pm 35.69	171.18 \pm 34.88	161.30 \pm 32.75	146.66 \pm 25.61	136.00 \pm 16.63
$\Sigma 16$ PCBs	0.77 \pm 0.41	0.77 \pm 0.37	0.88 \pm 0.56	0.60 \pm 0.21	0.75 \pm 0.35	0.74 \pm 0.17
$\Sigma 8$ PerPCBs	0.47 \pm 0.30	0.41 \pm 0.15	0.55 \pm 0.43	0.38 \pm 0.15	0.46 \pm 0.25	0.46 \pm 0.16
$\Sigma 6$ NonPerPCBs	0.24 \pm 0.16	0.29 \pm 0.19	0.26 \pm 0.21	0.17 \pm 0.07	0.23 \pm 0.12	0.22 \pm 0.06
$\Sigma 7$ EstrogenicPCBs	0.38 \pm 0.22	0.40 \pm 0.23	0.44 \pm 0.29	0.29 \pm 0.11	0.37 \pm 0.17	0.35 \pm 0.08
Antiestrogenic PCB	0.02 \pm 0.02	0.02 \pm 0.01	0.03 \pm 0.02	0.02 \pm 0.02	0.03 \pm 0.01	0.02 \pm 0.01
$\Sigma 4$ MOPCBs	0.15 \pm 0.09	0.16 \pm 0.07	0.17 \pm 0.13	0.13 \pm 0.06	0.15 \pm 0.07	0.14 \pm 0.04
$\Sigma 8$ DOPCBs	0.51 \pm 0.29	0.51 \pm 0.26	0.60 \pm 0.39	0.40 \pm 0.16	0.50 \pm 0.25	0.51 \pm 0.15
$\Sigma 3$ TTOPCBs	0.08 \pm 0.04	0.08 \pm 0.04	0.09 \pm 0.05	0.06 \pm 0.02	0.07 \pm 0.03	0.08 \pm 0.01
<i>p,p'</i> -DDE	0.45 \pm 0.35	0.31 \pm 0.11	0.48 \pm 0.50	0.40 \pm 0.19	0.51 \pm 0.35	0.52 \pm 0.25
HCB	0.04 \pm 0.02	0.03 \pm 0.02	0.04 \pm 0.03	0.04 \pm 0.02	0.05 \pm 0.03	0.03 \pm 0.01
Lead (μ g/dL)	1.59 \pm 0.97	1.48 \pm 0.72	1.67 \pm 0.94	1.25 \pm 0.93	1.79 \pm 1.03	1.70 \pm 1.23

$\Sigma 16$ PCB: congeners with $\geq 50\%$ detection rate, sum of PCBs 52, 70, 74, 84, 87, 95, 99, 101[+90], 105, 110, 118, 138[+163+164], 149[+123], 153, 180, 187; $\Sigma 8$ PerPCBs: sum of PCBs 74, 99, 105, 118, 138[+163+164], 153, 180, 187; $\Sigma 6$ NonPerPCBs: sum of PCBs 52, 84, 95, 101[+90], 110, 149[+123]; $\Sigma 7$ EstrogenicPCBs: sum of PCBs 52, 70, 95, 99, 101[+90], 110, 153 (Cooke et al. 2001); antiestrogenic PCB: PCB-105 (Cooke et al. 2001); $\Sigma 4$ MOPCBs: sum of PCBs 70, 74, 105, 118 (Goncharov et al. 2009); $\Sigma 8$ DOPCBs: sum of PCBs 52, 87, 99, 110, 138[+163+164], 153, 101[+90] (Goncharov et al. 2009); $\Sigma 3$ TTOPCBs: sum of PCBs 84, 95, 187 (Goncharov et al. 2009). Brackets indicate "minor" coeluting congener based on Aroclor concentration (Hansen 1999).

^aValues $<$ MDL were imputed from the estimated distribution $<$ MDL (ppb unless otherwise indicated). ^bThe functional sensitivity for testosterone was 10 ng/dL; for statistical purposes, results below the LOQ was set at 5 ng/dL.

males exceeded the 95th centile (0.40 ppb) of the same congeners (74, 99, 105, 118, 138 [163 + 164], 153, 180, 187) reported by the CDC for 12- to 19-year-old males and females based on NHANES (National Health and Nutrition Examination Study) data collected during 1999–2004 (CDC 2009). Geometric mean *p,p'*-DDE and HCB concentrations (0.39 ppb and 0.04 ppb, respectively) were lower than corresponding values reported by the CDC (1.69 ppb and 0.07 ppb, respectively).

Associations among toxicants and congeners were also examined (see Supplemental Material, Methods, Table S1). As expected, the different PCB congener groups were highly intercorrelated, reflecting the inclusion of many of the same congeners (range of $r = 0.53$ – 0.94). The levels of $\Sigma 16$ PCBs, $\Sigma 8$ PerPCBs, and $\Sigma 7$ EstrogenicPCBs were correlated with the level of *p,p'*-DDE ($r = 0.43$, 0.56 , and 0.34 respectively, $p < 0.01$). HCB was correlated with *p,p'*-DDE ($r = 0.41$, $p \leq 0.01$) and marginally with $\Sigma 8$ PerPCBs ($r = 0.20$, $p \leq 0.05$). Nonpersistent PCB groupings and the antiestrogenic PCBs were not significantly correlated with *p,p'*-DDE. Lead was not correlated with any of the PCB variables or with *p,p'*-DDE or HCB.

Testosterone and toxicant levels. Multivariable regression indicated a significant negative association between $\Sigma 16$ PCBs and serum testosterone, such that a 10% increase in $\Sigma 16$ PCBs was associated with a 5.6% decrease in testosterone (95% CI: -10.8 , -0.5%) (for complete model results, including associations with model covariates, see Supplemental Material, Table S2). There was a positive though nonsignificant association between *p,p'*-DDE and testosterone based on the same model, such that a 10% increase in *p,p'*-DDE was associated with a 5.2% increase in testosterone (95% CI: -0.5 , 10.9% , $p = 0.07$).

Other indices of PCB exposure ($\Sigma 8$ PerPCBs and congener groups used in the analysis of adult Akwesasne males) were tested using the same multivariable model (Table 2). Testosterone was negatively associated with $\Sigma 8$ PerPCBs, $\Sigma 4$ MOPCBs, and $\Sigma 7$ EstrogenicPCBs. For every 10% increase in these PCB groups, mean testosterone levels were 5.7% (95% CI: -11.0 , -0.4%), 6.2% (95% CI: -11.2 , -1.2%), and 4.7% (95% CI: -9.2 , -0.1%) lower, respectively. Associations with $\Sigma 8$ DOPCBs and $\Sigma 3$ TTOPCBs were also negative, but not significant (-4.6% ; 95% CI: -9.3 , 0.1% , $p = 0.06$ and -4.8% 95% CI: -10.3 , 0.7% , $p = 0.09$, respectively). The nonpersistent PCBs and the antiestrogenic PCB-105 also were negatively but not significantly associated with testosterone.

There was no clear evidence of non-additivity between PCBs and HCBs,

p,p'-DDE, or lead on associations with testosterone (all interaction p -values > 0.30). Associations between PCBs and testosterone were stronger in boys who were not breastfed compared with boys who were breastfed, though interactions also were not significant. For example, a 10% increase in $\Sigma 16$ PCBs was associated with a 9.7% decrease in testosterone (95% CI: -19.6 , 0.2%) among boys who were not breastfed, compared with a 4.5% decrease (95% CI: -10.7 , 1.6%) among boys who were breastfed (the interaction of breast feeding by $\Sigma 16$ PCBs was not significant; $p = 0.80$).

Discussion

Among male Mohawk youth, testosterone was negatively associated with several groupings of PCB congeners, and positively associated with *p,p'*-DDE. Because this is an observational, cross-sectional study, causality cannot be inferred, and, given the sample size, these observations should be tested in other samples.

The results are strengthened by certain features of the study. First, although focusing on PCBs, we were able to consider several other common toxicants, including a heavy metal (lead), and two pesticides (HCB and *p,p'*-DDE). Also, this study employed congener-specific laboratory analysis that allowed the calculation of PCB levels in subgroups of congeners. This enabled comparisons of associations between PCB subgroups that might differ in structure and/or persistence with testosterone. The more persistent PCBs were clearly associated negatively with testosterone, whereas nonpersistent PCBs were not. The lack of relationships with the nonpersistent congeners suggests that current exposure is not influential or is not great enough to have an observable effect.

Congener-specific laboratory analyses also allowed us to replicate the same structure-based PCB groups as Goncharov et al. (2009)

used in their study of Akwesasne men; our results in youth were similar in direction and statistical significance, suggesting that the adult profile may result from a developmental trajectory evident in adolescence. Furthermore, we were able to test several PCB congener groups used by other investigators to evaluate replicability across studies. It is important to recognize that the congener composition of the PCB composite variables overlap (see footnote of Tables 1 and 2).

An additional strength of this study is the use of TSs as an independent variable in lieu of age or height, which vary widely with markers of maturation during adolescence, and thus are a poor proxy for maturation in statistical models (Tanner 1962). Because timing of maturation itself could be affected by these toxicants, it is important to remove the effect of timing on testosterone levels. By statistically adjusting for maturation as indexed by TS, it is possible to estimate the effects of toxicants on testosterone independent of the effects on timing of sexual maturation.

A weakness, however, is the self-assessment of TS. Some researchers have found significant correlations between self-assessed TSs and hormonal development (Shirtcliff et al. 2009), whereas others have found that male adolescents both under- and overestimate their TS (Desmangles et al. 2006; Taylor et al. 2001; Williams et al. 1988). However, to the best of our knowledge, there is no evidence that over- or underestimation varies with toxicant exposure. Without such evidence, we believe that the error is most likely unbiased, although bias cannot be ruled out.

Negative associations between PCBs and testosterone were weaker for less persistent congeners than for more persistent congeners, which suggests that earlier exposures may be more relevant to the associations. However, it is not possible to assess the temporal relation

Table 2. Predictors of testosterone levels in adolescent males: results of the multivariable regression analysis with PCB groupings ($n = 120$).

Toxicant (ppb) ^a	β (95% CI)	p -Value	Percent change ^b
$\Sigma 16$ PCBs	-0.56 (-1.08 , -0.05)	0.03	5.6
$\Sigma 8$ PerPCBs	-0.57 (-1.10 , -0.04)	0.03	5.7
$\Sigma 6$ NonPerPCBs	-0.31 (-0.72 , 0.09)	0.13	3.1
$\Sigma 7$ EstrogenicPCBs	-0.47 (-0.92 , -0.01)	0.05	4.7
Antiestrogenic PCB	-0.15 (-0.59 , 0.29)	0.50	1.5
$\Sigma 4$ MOPCBs	-0.62 (-1.12 , -0.12)	0.02	6.2
$\Sigma 8$ DOPCBs	-0.46 (-0.93 , 0.01)	0.06	4.6
$\Sigma 3$ TTOPCBs	-0.48 (-1.03 , 0.07)	0.09	4.8

Each model controlled for Tanner stages (TS) I and II vs. III, TS IV and V vs. III, weight-for-age z-score, height-for-age z-score, breastfed as an infant (yes/no), child's alcohol use (yes/no), child's cigarette use (yes/no), triglycerides (mg/dL), cholesterol (mg/dL), lead (μ dL), *p,p'*-DDE (ppb), and HCB (ppb). Testosterone, cholesterol, triglycerides, lead, *p,p'*-DDE, HCB and PCB variables are ln-transformed. $\Sigma 16$ PCB: congeners with $\geq 50\%$ detection rate, sum of PCBs 52, 70, 74, 84, 87, 95, 99, 101[+90], 105, 110, 118, 138[+163+164], 149[+123], 153, 180, 187; $\Sigma 8$ PerPCBs: sum of PCBs 74, 99, 105, 118, 138[+163+164], 153, 180, 187; $\Sigma 6$ NonPerPCBs: sum of PCBs 52, 84, 95, 101[+90], 110, 149[+123]; $\Sigma 7$ EstrogenicPCBs: sum of PCBs 52, 70, 95, 99, 101[+90], 110, 153 (Cooke et al. 2001); antiestrogenic PCB: PCB-105 (Cooke et al. 2001); $\Sigma 4$ MOPCBs: sum of PCBs 70, 74, 105, 118 (Goncharov et al. 2009); $\Sigma 8$ DOPCBs: sum of PCBs 52, 87, 99, 110, 138[+163+164], 153, 101[+90] (Goncharov et al. 2009); $\Sigma 3$ TTOPCBs: sum of PCBs 84, 95, 187 (Goncharov et al. 2009). Brackets indicate "minor" coeluting congener based on Aroclor concentration (Hansen, 1999).

^aValues $< MDL$ were imputed from the estimated distribution $< MDL$. ^bPercent change associated with a 10% increase in exposure.

between exposures and the outcome, given the cross-sectional study design. Finally, because the Akwesasne Mohawk Nation is not federally censused, it is not possible to know what proportion of the community between 10 and < 17 years of age is sampled.

The clinical relevance of the differences in testosterone associated with *p,p'*-DDE and PCB exposures in the study population is not known, because reference values for serum testosterone concentrations are not available (Kronenberg and Williams 2008). Nevertheless, associations between environmental exposures and testosterone levels in the study population are a concern, given that adolescence is a critical period for the establishment of adult hormone homeostasis (Root 1973). Similar negative associations estimated between PCBs and testosterone in adult Mohawk men from the same community suggest the possibility of early-life effects that may persist into adulthood (Goncharov et al. 2009).

An interesting finding is that the associations of PCBs and *p,p'*-DDE with testosterone were stronger among boys who had not been breastfed. It is difficult to attribute the weaker effect among the breastfed youth to a moderating effect of breastfeeding because breastfeeding delivers additional exposure to PCBs and other lipophilic compounds (Greizerstein et al. 1999). In the Akwesasne sample here, the level of PCBs was significantly higher among those who had been breastfed compared with those not breastfed (Gallo et al. 2011; Schell et al. 2003). Toxicants delivered through lactation may not influence some end points if the period of developmental sensitivity of an end point has passed. In such circumstances, lactation essentially delivers an additional toxicant burden randomly with regard to the dependent variable. This may obfuscate the relationship rather than mitigate it. This interpretation is consistent with results regarding thyroid hormones (Schell et al. 2009), but is limited by the small samples of breastfed and not breastfed boys.

Several studies have examined such relationships among boys within a narrow age range. Studies of Flemish adolescents, 14–15 years of age, from areas differing in exposure characteristics examined hormone and toxicant levels measured concurrently. These studies have produced evidence of both stimulated and decreased testosterone in relation to marker PCBs and pollution, but no associations with *p,p'*-DDE (Croes et al. 2009; Den Hond et al. 2002; Dhooge et al. 2011). Finally, a study of neonatal hexachlorobiphenyl exposure in rats found decreased adult serum testosterone levels (Xiao et al. 2011).

Three studies have evaluated relationships between pubertal development and prenatal exposure to PCBs, PCDFs (polychlorinated

dibenzofurans), or *p,p'*-DDE. In a longitudinal study of 304 singleton males born in the early 1960s and followed through adolescence (Gladden et al. 2004), *p,p'*-DDE levels in cord blood were not associated with testosterone (*p*-values for all tests > 0.10). There was some evidence of reduced pubertal testosterone in Yucheng boys (*n* = 21) who had experienced an acute prenatal exposure to a mixture of PCBs and PCDFs when they were compared with matched controls (Hsu et al. 2005). PCB levels measured in cord blood (PCB-138[+163+164], 153, and 180 combined) were not associated with serum testosterone in Faroese boys at 13–14 years of age (Mol et al. 2002). In our analysis, serum PCB and *p,p'*-DDE concentrations were not highly correlated, and mutually adjusted associations with testosterone were in opposite directions.

Two studies of adult men reported significant negative associations between some PCBs (mono-, tri-, and di-*ortho* substituted PCBs) and testosterone (Goncharov et al. 2009; Richthoff et al. 2003). Great Lakes sport fish consumers' PCB levels were negatively associated with SHBG (sex hormone binding globulin)-bound testosterone, though not with total or free testosterone (Persky et al. 2001; Turyk et al. 2006). Other studies of human populations have produced evidence suggesting relationships of organochlorines to measures of reproductive hormones or function (Ferguson et al. 2012; Richthoff et al. 2003; Rignell-Hydbom et al. 2004), or no evidence of a relationship in adult men (Hagmar et al. 2001). Of the studies of adult men with PCB exposures (Ferguson et al. 2012; Goncharov et al. 2009; Richthoff et al. 2003; Rignell-Hydbom et al. 2004), the study of Akwesasne men with high exposure has provided the strongest evidence for a relationship of testosterone with PCBs (Goncharov et al. 2009). Serum PCB levels measured in our study population of Akwesasne boys in 1996–2000 were higher than serum levels measured in a representative sample of U.S. adolescents (age 12–19 years) in 1999–2004, but similar to levels measured among Akwesasne men during the same time period (Goncharov et al. 2009). Differences in exposures among populations could explain differences in associations among studies (Goncharov et al. 2009).

Conclusion

In this study we found significant associations between testosterone and Σ 16PCBs, with stronger associations estimated for more persistent congeners, and *p,p'*-DDE in a population of adolescent Native American males with relatively high PCB exposures. These results are consistent with experimental studies in animals that found endocrine-disrupting

chemicals, such as PCBs and *p,p'*-DDE, modified serum testosterone levels (Ahmad et al. 2003; Andric et al. 2000; Kovacevic et al. 1995; Xiao et al. 2011), and with studies of humans that found associations consistent with endocrine modification due to those exposures (Croes et al. 2009; Dhooge et al. 2011; Goncharov et al. 2009; Persky et al. 2001; Richthoff et al. 2003; Turyk et al. 2006).

REFERENCES

- Abaci A, Demir K, Bober E, Buyukgebiz A. 2009. Endocrine disruptors—with special emphasis on sexual development. *Pediatr Endocrinol Rev* 6:464–475.
- Ahmad SU, Tariq S, Jalali S, Ahmad MM. 2003. Environmental pollutant Aroclor 1242 (PCB) disrupts reproduction in adult male Rhesus monkeys (*Macaca mulatta*). *Environ Res* 93:272–278.
- Alexander BH, Checkoway H, Costa-Mallen P, Faustman EM, Woods JS, Kelsey KT, et al. 1998. Interaction of blood lead and δ -aminolevulinic acid dehydratase genotype on markers of heme synthesis and sperm production in lead smelter workers. *Environ Health Perspect* 106:213–216.
- Alexander BH, Checkoway H, van Netten C, Muller CH, Ewers TG, Kaufman JD, et al. 1996. Semen quality of men employed at a lead smelter. *Occup Environ Med* 53:411–416.
- Andric SA, Kostic TS, Stojilkovic SS, Kovacevic RZ. 2000. Inhibition of rat testicular androgenesis by a polychlorinated biphenyl mixture Aroclor 1248. *Biol Reprod* 62:1882–1888.
- Benoff S, Centola GM, Millan C, Napolitano B, Marmar JL, Hurley IR. 2003a. Increased seminal plasma lead levels adversely affect the fertility potential of sperm in IVF. *Hum Reprod* 18:374–383.
- Benoff S, Hurley IR, Millan C, Napolitano B, Centola GM. 2003b. Seminal lead concentrations negatively affect outcomes of artificial insemination. *Fertil Steril* 80:517–525.
- Bourguignon JP, Parent AS. 2012. The impact of endocrine disruptors on female timing. In: *Endocrine Disruptors and Puberty* (Diamanti-Kandarakis E, Gore AC, eds). New York:Humana Press, 325–338.
- Carpenter DO. 1998. Polychlorinated biphenyls and human health. *Int J Occup Med Environ Health* 11:291–303.
- CDC (Centers for Disease Control and Prevention). 2001. Epi Info. Part 6.04d. Available: http://www.cdc.gov/epiinfo/html/ei6_downloads.htm [accessed 1 November 2010].
- CDC (Centers for Disease Control and Prevention). 2009. Fourth National Report on Human Exposure to Environmental Chemicals. Available: <http://www.cdc.gov/exposurereport/pdf/fourthreport.pdf> [accessed 1 September 2010].
- Cooke PS, Sato T, Buchanan DL. 2001. Disruption of steroid hormone signaling by PCBs. In: *PCBs Recent Advances in Environmental Toxicology and Health Effects* (Robertson LW, Hansen LG, eds). Louisville, KY:University Press of Kentucky, 257–263.
- Croes K, Baeyens W, Bruckers L, Den Hond E, Koppen G, Nelen V, et al. 2009. Hormone levels and sexual development in Flemish adolescents residing in areas differing in pollution pressure. *Int J Hyg Environ Health* 212:612–625.
- DeCaprio AP, Johnson GW, Tarbell AM, Carpenter DO, Chiarenzelli JR, Morse GS, et al. 2005. Polychlorinated biphenyl (PCB) exposure assessment by multivariate statistical analysis of serum congener profiles in an adult Native American population. *Environ Res* 98:284–302.
- DeCaprio AP, Tarbell AM, Bott A, Wagemaker DL, Williams RL, O'Hehir CM. 2000. Routine analysis of 101 polychlorinated biphenyl congeners in human serum by parallel dual-column gas chromatography with electron capture detection. *J Anal Toxicol* 24:403–420.
- Den Hond E, Dhooge W, Bruckers L, Schoeters G, Nelen V, van de Mierop E, et al. 2011. Internal exposure to pollutants and sexual maturation in Flemish adolescents. *J Expo Sci Environ Epidemiol* 21(3):224–233.
- Den Hond E, Roels HA, Hoppenbrouwers K, Nawrot T, Thijs L, Vandermeulen C, et al. 2002. Sexual maturation in relation to polychlorinated aromatic hydrocarbons: Sharpe and Skakkebaek's hypothesis revisited. *Environ Health Perspect* 110:771–776.

- Denham M, Schell LM, Deane G, Gallo MV, Ravenscroft J, DeCaprio AP, et al. 2005. Relationship of lead, mercury, mirex, dichlorodiphenylchloroethylene, hexachlorobenzene, and polychlorinated biphenyls to timing of menarche among Akwesasne Mohawk girls. *Pediatrics* 115:e127-e134.
- Desmangles JC, Lappe JM, Lipaczewski G, Haynatzki G. 2006. Accuracy of pubertal Tanner staging self-reporting. *J Pediatr Endocrinol Metab* 19:213–221.
- Dhooge W, Den Hond E, Bruckers L, Schoeters G, Nelen V, van de Mieroop E, et al. 2011. Internal exposure to pollutants and sex hormone levels in Flemish male adolescents in a cross-sectional study: associations and dose-response relationships. *J Expo Sci Environ Epidemiol* 21:224–233.
- Dhooge W, van Larebeke N, Koppen G, Nelen V, Schoeters G, Vlietinck R, et al. 2006. Serum dioxin-like activity is associated with reproductive parameters in young men from the general Flemish population. *Environ Health Perspect* 114:1670–1676; doi:10.1289/ehp.9224.
- Duke PM, Litt IF, Gross RT. 1980. Adolescents' self-assessment of sexual maturation. *Pediatrics* 66:918–920.
- Ferguson KK, Hauser R, Altshul L, Meeker JD. 2012. Serum concentrations of *p,p'*-DDE, HCB, PCBs and reproductive hormones among men of reproductive age. *Reprod Toxicol* 34(3):429–435; doi:10.1016/j.reprotox.2012.04.006.
- Fischer LJ, Seegal RF, Ganey PE, Pessah IN, Kodavanti PR. 1998. Symposium overview: toxicity of non-coplanar PCBs. *Toxicol Sci* 41:49–61.
- Fitzgerald EF, Hwang S-A, Brix KA, Bush B, Cook K, Worswick P. 1995. Fish PCB concentrations and consumption patterns among Mohawk women at Akwesasne. *J Expo Anal Environ Epidemiol* 5:1–19.
- Forest MG, De Peretti E, Bertrand J. 1976. Hypothalamic-pituitary-gonadal relationships in man from birth to puberty. *Clin Endocrinol (Oxf)* 5:551–569.
- Forti A, Bogdan KG, Horn E. 1995. Health Risk Assessment for The Akwesasne Mohawk Population from Exposure to Chemical Contaminants in Fish and Wildlife. Albany, NY:New York State Department of Health.
- Fox, J. 2008. *Applied Regression Analysis and Generalized Linear Models*. 2nd ed. Los Angeles, CA:Sage.
- Gallo MV, Schell LM, Akwesasne Task Force On The Environment. 2005. Height, weight and body mass index among Akwesasne Mohawk youth. *Am J Hum Biol* 17:269–279.
- Gallo MV, Schell LM, DeCaprio AP, Jacobs A. 2011. Levels of persistent organic pollutants and their predictors among young adults. *Chemosphere* 83:1374–1382.
- Gladen BC, Klebanoff MA, Hediger ML, Katz SH, Barr DB, Davis MD, et al. 2004. Prenatal DDT exposure in relation to anthropometric and pubertal measures in adolescent males. *Environ Health Perspect* 112:1761–1767; doi:10.1289/ehp.7287.
- Gladen BC, Ragan NB, Rogan WJ. 2000. Pubertal growth and development and prenatal and lactational exposure to polychlorinated biphenyls and dichlorodiphenyl dichloroethene. *J Pediatr* 136:490–496.
- Goncharov A, Rej R, Negoita S, Schymura M, Santiago-Rivera A, Morse G, et al. 2009. Lower serum testosterone associated with elevated polychlorinated biphenyl concentrations in Native American men. *Environ Health Perspect* 117:1454–1460; doi:10.1289/ehp.0800134.
- Greizerstein HB, Stinson C, Mendola P, Buck GM, Kostyniak PJ, Vena JE. 1999. Comparison of PCB congeners and pesticide levels between serum and milk from lactating women. *Environ Res* 80:280–286.
- Gupta AK. 1952. Estimation of the mean and standard deviation of a normal population from a censored sample. *Biometrika* 39(3/4):260–273.
- Hagmar L, Bjork J, Sjodin A, Bergman A, Erfurth EM. 2001. Plasma levels of persistent organohalogen and hormone levels in adult male humans. *Arch Environ Health* 56:138–143.
- Hansen LG. 1999. *The ortho Side of PCBs: Occurrence and Disposition*. Boston, MA:Kluwer Academic Publishers.
- Hauser R. 2006. The environment and male fertility: recent research on emerging chemicals and semen quality. *Semin Reprod Med* 24:156–167.
- Hauser R, Chen Z, Pothier L, Ryan L, Altshul L. 2003. The relationship between human semen parameters and environmental exposure to polychlorinated biphenyls and *p,p'*-DDE. *Environ Health Perspect* 111:1505–1511; doi:10.1289/ehp.6175.
- Hsu PC, Lai TJ, Guo NW, Lambert GH, Leon Guo Y. 2005. Serum hormones in boys prenatally exposed to polychlorinated biphenyls and dibenzofurans. *J Toxicol Environ Health A* 68:1447–1456.
- Humblet O, Williams PL, Korrick SA, Sergeev O, Emond C, Birnbaum LS, et al. 2011. Dioxin and polychlorinated biphenyl concentrations in mother's serum and the timing of pubertal onset in sons. *Epidemiology* 22:827–835.
- IBM. 2011. *PASW Statistical Package for the Social Sciences*. Part 19. New York:IBM Corporation.
- Kim R, Hu H, Rotnitzky A, Bellinger DC, Needleman HL. 1995. A longitudinal study of chronic lead exposure and physical growth in Boston children. *Environ Health Perspect* 103:952–957.
- Korrick SA, Lee MM, Williams PL, Sergeev O, Burns JS, Patterson DG, et al. 2011. Dioxin exposure and age of pubertal onset among Russian boys. *Environ Health Perspect* 119:1339–1344; doi:10.1289/ehp.1003102.
- Kovacevic R, Vojinovic-Miloradov M, Teodorovic I, Andric SA. 1995. Effect of PCBs on androgen production by suspension of adult rat Leydig cells *in vitro*. *J Steroid Biochem Mol Biol* 52:595–597.
- Kronenberg H, Williams RH. 2008. *Williams Textbook of Endocrinology*. 11th ed. Philadelphia, PA:Saunders/Elsevier.
- Lacetti G. 1993. *Public Health Assessment. General Motors/Central Foundry Division*. Albany:New York State Department of Health.
- Lohman TG, Roche AF, Martorell R. 1988. *Anthropometric Standardization Reference Manual*. Champaign, IL:Human Kinetics Books.
- Mocarelli P, Gerthoux PM, Patterson DG Jr, Milani S, Limonta G, Bertona M, et al. 2008. Dioxin exposure, from infancy through puberty, produces endocrine disruption and affects human semen quality. *Environ Health Perspect* 116:70–77; doi:10.1289/ehp.10399.
- Mol NM, Sorensen N, Weihe P, Andersson AM, Jorgensen N, Skakkebaek NE, et al. 2002. Spermatid and serum hormone concentrations at the age of puberty in boys prenatally exposed to polychlorinated biphenyls. *Eur J Endocrinol* 146:357–363.
- Morris NM, Udry JR. 1980. Validation of a self-administered instrument to assess stage of adolescent development. *J Youth Adolesc* 9:271–280.
- Oskam IC, Ropstad E, Dahl E, Lie E, Derocher AE, Wiig O, et al. 2003. Organochlorines affect the major androgenic hormone, testosterone, in male polar bears (*Ursus maritimus*) at Svalbard. *J Toxicol Environ Health A* 66:2119–2139.
- Ouyang F, Perry MJ, Venners SA, Chen C, Wang B, Yang F, et al. 2005. Serum DDT, age at menarche, and abnormal menstrual cycle length. *Occup Environ Med* 62:878–884.
- Ozen S, Darcan S, Bayindir P, Karasulu E, Simsek D, Gurler T. 2012. Effects of pesticides used in agriculture on the development of precocious puberty. *Environ Monit Assess* 184:4223–4232.
- Persky V, Turyk M, Anderson HA, Hanrahan LP, Falk C, Steenport DN, et al. 2001. The effects of PCB exposure and fish consumption on endogenous hormones. *Environ Health Perspect* 109:1275–1283.
- Pflieger-Bruss S, Schuppe HC, Schill WB. 2004. The male reproductive system and its susceptibility to endocrine disrupting chemicals. *Andrologia* 36:337–345.
- Richthoff J, Rylander L, Jönsson BA, Akesson H, Hagmar L, Nilsson-Ehle P, et al. 2003. Serum levels of 2,2',4,4',5,5'-hexachlorobiphenyl (CB-153) in relation to markers of reproductive function in young males from the general Swedish population. *Environ Health Perspect* 111:409–413; doi:10.1289/ehp.5767.
- Rignell-Hydbom A, Rylander L, Giwercman A, Jonsson BA, Nilsson-Ehle P, Hagmar L. 2004. Exposure to CB-153 and *p,p'*-DDE and male reproductive function. *Hum Reprod* 19:2066–2075.
- Root AW. 1973. *Endocrinology of puberty*. I. Normal sexual maturation. *J Pediatr* 83:1–19.
- Roy JR, Chakraborty S, Chakraborty TR. 2009. Estrogen-like endocrine disrupting chemicals affecting puberty in humans—a review. *Med Sci Monit* 15:RA137–RA145.
- Schell LM, Gallo MV, Cook K. 2012. What's not to eat—food adulteration in the context of human biology. *Am J Hum Biol* 24:139–148.
- Schell LM, Gallo MV, Denham M, Ravenscroft J, DeCaprio AP, Carpenter DO. 2008. Relationship of thyroid hormone levels to levels of polychlorinated biphenyls, lead, *p,p'*-DDE, and other toxicants in Akwesasne Mohawk youth. *Environ Health Perspect* 116:806–813; doi:10.1289/ehp.10490.
- Schell LM, Gallo MV, Ravenscroft J, DeCaprio AP. 2009. Persistent organic pollutants and anti-thyroid peroxidase levels in Akwesasne Mohawk young adults. *Environ Res* 109:86–92.
- Schell LM, Hubicki LA, DeCaprio AP, Gallo MV, Ravenscroft J, Tarbell A, et al. 2003. Organochlorines, lead, and mercury in Akwesasne Mohawk youth. *Environ Health Perspect* 111:954–961; doi:10.1289/ehp.5990.
- Selevan SG, Rice DC, Hogan KA, Euling SY, Pfahles-Hutchens A, Bethel J. 2003. Blood lead concentration and delayed puberty in girls. *N Engl J Med* 348:1527–1536.
- Shirtcliff EA, Dahl RE, Pollak SD. 2009. Pubertal development: Correspondence between hormonal and physical development. *Child Dev* 80:327–337.
- Sloan RJ, Jock K. 1990. Chemical Contaminants in Fish from the St. Lawrence River Drainage on Lands of the Mohawk Nation at Akwesasne and Near the General Motors Corporation/Central Foundry Division, Massena, NY Plant. Albany:New York State Department of Environmental Conservation.
- Su PH, Huang PC, Lin CY, Ying TH, Chen JY, Wang SL. 2012. The effect of in utero exposure to dioxins and polychlorinated biphenyls on reproductive development in eight year-old children. *Environ Int* 39:181–187.
- Tanner JM. 1962. *Growth at Adolescence*. Oxford, UK:Blackwell Scientific Publications.
- Taylor SJ, Whincup PH, Hindmarsh PC, Lampe F, Odoki K, Cook DG. 2001. Performance of a new pubertal self-assessment questionnaire: a preliminary study. *Paediatr Perinat Epidemiol* 15:88–94.
- Teilmann G, Juul A, Skakkebaek NE, Toppari J. 2002. Putative effects of endocrine disrupters on pubertal development in the human. *Best Pract Res Clin Endocrinol Metab* 16:105–121.
- Toft G, Rignell-Hydbom A, Tyrkiel E, Shvets M, Giwercman A, Lindh CH, et al. 2006. Semen quality and exposure to persistent organochlorine pollutants. *Epidemiology* 17:450–458.
- Turyk ME, Anderson HA, Freels S, Chatterton R, Jr., Needham LL, Patterson DG Jr, et al. 2006. Associations of organochlorines with endogenous hormones in male Great Lakes fish consumers and nonconsumers. *Environ Res* 102:299–307.
- U.S. EPA (U.S. Environmental Protection Agency). 1998. *Guidance for Data Quality Assessment. Practical Methods for Data Analysis*. QA/G-9. Washington, DC:U.S. EPA.
- Vasiliu O, Muttineni J, Karmaus W. 2004. *In utero* exposure to organochlorines and age at menarche. *Hum Reprod* 19:1506–1512.
- Williams PL, Sergeev O, Lee MM, Korrick SA, Burns JS, Humblet O, et al. 2010. Blood lead levels and delayed onset of puberty in a longitudinal study of Russian boys. *Pediatrics* 125:e1088–e1096.
- Williams RL, Cheyne KL, Houtkooper LK, Lohman TG. 1988. Adolescent self-assessment of sexual maturation. Effects of fatness classification and actual sexual maturation stage. *J Adolesc Health Care* 9:480–482.
- Wolff MS, Anderson HA. 1999. Polybrominated biphenyls: Sources and disposition of exposure among Michigan farm residents, 1976–1980. *Eur J Oncol* 4:645–651.
- Wolff MS, Britton JA, Boguski L, Hochman S, Maloney N, Serra N, et al. 2008. Environmental exposures and puberty in inner-city girls. *Environ Res* 107(3):393–400.
- Wolff MS, Camann D, Gammon M, Stellman SD. 1997. Proposed PCB congener groupings for epidemiological studies. *Environ Health Perspect* 105:13–14.
- Xiao W, Zhang J, Liang J, Zhu H, Zhou Z, Wu Q. 2011. Adverse effects of neonatal exposure to 3,3',4,4',5,5'-hexachlorobiphenyl on hormone levels and testicular function in male Sprague-Dawley rats. *Environ Toxicol* 26:657–668.
- Yang CY, Yu ML, Guo HR, Lai TJ, Hsu CC, Lambert G, Guo YL. 2005. The endocrine and reproductive function of the female Yucheng adolescents prenatally exposed to PCBs/PCDFs. *Chemosphere* 61(3):355–60.
- Yeowell HN, Waxman DJ, Wadhera A, Goldstein JA. 1987. Suppression of the constitutive, male-specific rat hepatic cytochrome P-450 2c and its mRNA by 3,4,5,3',4',5'-hexachlorobiphenyl and 3-methylcholanthrene. *Mol Pharmacol* 32:340–347.