

Structural and Functional Profiling of Environmental Ligands for Estrogen Receptors

Vanessa Delfosse,^{1,2} Marina Grimaldi,^{3,4,5,6} Vincent Cavallès,^{3,4,5,6} Patrick Balaguer,^{3,4,5,6*} and William Bourguet^{1,2*}

¹Inserm (Institut national de la santé et de la recherche médicale) U1054, Montpellier, France; ²CNRS (Centre national de la recherche scientifique) UMR 5048, Universités Montpellier 1 & 2, Centre de Biochimie Structurale, Montpellier, France; ³IRCM (Institut de Recherche en Cancérologie de Montpellier), Montpellier, France; ⁴Inserm, U896, Montpellier, France; ⁵Université Montpellier 1, Montpellier, France; ⁶ICM (Institut régional du Cancer de Montpellier), Montpellier, France

*These authors contributed equally to this work.

BACKGROUND: Individuals are exposed daily to environmental pollutants that may act as endocrine-disrupting chemicals (EDCs), causing a range of developmental, reproductive, metabolic, or neoplastic diseases. With their mostly hydrophobic pocket that serves as a docking site for endogenous and exogenous ligands, nuclear receptors (NRs) can be primary targets of small molecule environmental contaminants. However, most of these compounds are chemically unrelated to natural hormones, so their binding modes and associated hormonal activities are hardly predictable.

OBJECTIVES: We conducted a correlative analysis of structural and functional data to gain insight into the mechanisms by which 12 members of representative families of pollutants bind to and activate the estrogen receptors ER α and ER β .

METHODS: We used a battery of biochemical, structural, biophysical, and cell-based approaches to characterize the interaction between ERs and their environmental ligands.

RESULTS: Our study revealed that the chemically diverse compounds bound to ERs via varied sets of protein–ligand interactions, reflecting their differential activities, binding affinities, and specificities. We observed xenoestrogens binding to both ERs—with affinities ranging from subnanomolar to micromolar values—and acting in a subtype-dependent fashion as full agonists or partial agonists/antagonists by using different combinations of the activation functions 1 and 2 of ER α and ER β .

CONCLUSIONS: The precise characterization of the interactions between major environmental pollutants and two of their primary biological targets provides rational guidelines for the design of safer chemicals, and will increase the accuracy and usefulness of structure-based computational methods, allowing for activity prediction of chemicals in risk assessment.

CITATION: Delfosse V, Grimaldi M, Cavallès V, Balaguer P, Bourguet W. 2014. Structural and functional profiling of environmental ligands for estrogen receptors. *Environ Health Perspect* 122:1306–1313; <http://dx.doi.org/10.1289/ehp.1408453>

Introduction

Endocrine-disrupting chemicals (EDCs) are exogenous substances that interfere with the function of hormonal systems and cause deleterious effects on humans and wildlife (De Coster and van Larebeke 2012; Diamanti-Kandarakis et al. 2009; Schug et al. 2011). Many EDCs are man-made chemicals produced by industry and released into the environment, but some naturally occurring EDCs can also be found in plants or fungi. EDCs can affect the endocrine system of an organism in a wide variety of ways. These include mimicking natural hormones, antagonizing their action, or modifying their synthesis, metabolism, and transport. Moreover, these substances can act via multiple pathways, including membrane receptors, the aryl hydrocarbon receptor, or the enzymes involved in hormone biosynthesis and metabolism (De Coster and van Larebeke 2012). Yet, most of the reported harmful effects of EDCs are ascribed to their interaction with members of the nuclear receptor (NR) family, including the estrogen receptors ER α and ER β for which a large panel of exogenous ligands have been identified. The group of molecules acting as ER environmental ligands is highly heterogeneous and includes natural phytoestrogens or

mycoestrogens, as well as industrial compounds such as pesticides, plasticizers, surfactants, or UV (ultraviolet) filters (Li et al. 2013).

ERs and their endogenous ligand, 17 β -estradiol (E₂), play important roles in the growth and maintenance of a diverse range of tissues. As a consequence, dysfunctional ER signaling (i.e., inappropriate exposure to environmental pollutants) may lead to hormonal cancers, infertility, obesity, or diabetes (Grun and Blumberg 2007; Newbold et al. 2009; Rubin and Soto 2009). ER α is expressed primarily in the uterus, liver, kidney, and heart, whereas ER β is expressed primarily in the ovary, prostate, lung, gastrointestinal tract, bladder, and hematopoietic and central nervous systems. ER α and ER β are also coexpressed in a number of tissues including the mammary, thyroid, or adrenal glands; bone; and some regions of the brain. Although ER α and ER β share similar mechanisms of action, several differences in the transcriptional abilities of each receptor, as well as distinct phenotypes between gene-null animals, have been identified and suggest that these receptors may regulate distinct cellular pathways (Couse and Korach 1999; Curtis et al. 1996). Interestingly, ER β has been shown to antagonize the effects mediated by ER α on cell

proliferation in the breast, uterus, ovary, and prostate (Docquier et al. 2013; Lindberg et al. 2003; Weihua et al. 2000). In this regard, EDCs with selectivity for either ER subtypes may produce different biological outcomes, particularly on cancer cell proliferation.

Like other members of this family, ERs contain three major functional domains, including an N-terminal domain that harbors a transcriptional activation function (AF-1), a DNA-binding domain, and a C-terminal ligand-binding domain (LBD) hosting a ligand-dependent transcriptional activation function (AF-2). The LBD is crucially involved in most of the receptor functions because of its capacity for hormone binding, dimerization, and interaction with coregulators. The LBD also contributes to the modulation of the N-terminal AF-1 through interdomain cross-talk so that both AF-1 and AF-2 domains can recruit a range of coregulatory proteins and act either individually or in a synergistic manner

Address correspondence to W. Bourguet, Centre de Biochimie Structurale, 29 rue de Navacelles, Montpellier 34090 France. Telephone: (33) 467417702. E-mail: bourguet@cbs.cnrs.fr; or P. Balaguer, Institut de Recherche en Cancérologie de Montpellier, Campus Val d'Aurelle, 208 rue des Apothicaires, Montpellier, F34298, France. Telephone: (33) 467612409. E-mail: patrick.balaguer@icm.unicancer.fr

Supplemental Material is available online (<http://dx.doi.org/10.1289/ehp.1408453>).

We acknowledge experimental assistance from the staff of the European Synchrotron Radiation Facility (Grenoble, France) during data collection.

Coordinates and structure factors have been deposited in the Protein Data Bank as follows: 4MG5 (chlordecone), 4MG6 (benzyl-butyl-phthalate), 4MG7 (ferutinine), 4MG8 (α -zearalanol), 4MG9 (butylparaben), 4MGA (4-*tert*-octylphenol), 4MGB (TCBPA; tetrachlorobisphenol-A), 4MGC (benzophenone-2), 4MGD [HPTE; 2,2-bis(*p*-hydroxyphenyl)-1,1,1-trichloroethane].

This study received financial support from the Programme National de Recherche sur les Perturbateurs Endocriniens (PNRPE) (P.B.); the Agence Nationale de la Recherche, programme Contaminants, Environnement, Santé (ANR CESA) (P.B. and W.B.); and the Agence Nationale de Sécurité Sanitaire de l'Alimentation, de l'Environnement et du Travail (ANSES) (P.B. and W.B.).

The authors declare they have no actual or potential competing financial interests.

Received: 21 March 2014; Accepted: 25 September 2014; Advance Publication: 26 September 2014; Final Publication: 1 December 2014.

(Kobayashi et al. 2000; Métivier et al. 2001). It is noteworthy that the diversity of transcriptional coregulators mediating the effect of ERs on gene expression combined with variations in their expression levels and posttranslational modifications affect the specificity of the response depending on the cell type, ligand, or target gene considered (McDonnell and Wardell 2010).

Our recent work has shown that approaches combining structural, biophysical, and cell-based techniques are helpful in understanding how environmental compounds that are structurally and chemically divergent from natural ligands can interact with NRs and impact their signaling pathways (Delfosse et al. 2012; le Maire et al. 2009; Riu et al. 2011). In the present study, we used a similar approach to gain insight into the mechanisms by which 12 contaminants interact with ERs and modulate their AFs. The compounds used in this study (Figure 1) were selected on the basis of their structural diversity and because they belong to the most representative families of ER environmental ligands. These are bisphenol A (BPA) and bisphenol C (BPC), both used as plasticizers; the flame retardant tetrachlorobisphenol A (TCBPA); the preservative butylparaben; the surfactant 4-*tert*-octylphenol (4-OP); the UV filter benzophenone-2 (BP-2); the pesticides 2,2-bis(*p*-hydroxyphenyl)-1,1,1-trichloroethane (HPTE, a methoxychlor metabolite), dichlorodiphenyldichloroethylene (DDE, a DDT metabolite), and chlordecone;

benzylbutylphthalate (BBP); the phytoestrogen ferutinine; and the growth stimulant α -zearalanol (α -ZA), a double reduction product of the mycoestrogen zearalenone.

Materials and Methods

Ligands and peptides. E₂ (CAS number 50-28-2, purity 98%), BP-2 (CAS number 131-55-5, purity 97%), α -ZA (CAS number 247-769-0, purity 97%), 4-OP (CAS number 140-66-9, purity 97%), HPTE (CAS number 2971-36-0, purity 98%), BPA (CAS number 80-05-7, purity 99%), BPC (CAS number 14868-03-2, purity 99%), DDE (CAS number 72-55-9, purity 99%), butylparaben (CAS number 94-26-8, purity 99%), chlordecone (CAS number 143-50-0, purity 99%), and BBP (CAS number 85-68-7, purity 98%) were purchased from Sigma-Aldrich (Saint-Quentin-Fallavier, France). We obtained TCBPA (CAS number 79-95-8, purity 98%) from TCI Europe (Zwijndrecht, Belgium); ferutinine (CAS number 41743-44-6, purity 98%) from Santa Cruz Biotechnology Inc. (Dallas, TX, USA); and 4-hydroxytamoxifen (OHT; CAS number 68392-35-8, purity 99%) from Zeneca (Macclesfield, UK). Compounds were dissolved in DMSO to prepare 10⁻² M stock solutions. The fluorescein-RHKILHRLQEGS peptide corresponding to the NR box 2-binding motif of SRC-1 was purchased from EZbiolab (Westfield, IN, USA).

Reporter cell lines and culture condition. Luciferase and whole-cell ER competitive

binding assays were performed using the reporter cell lines HGELN-ER α , HGELN-ER β , HGELN- Δ AB-ER α , and HGELN- Δ AB-ER β as described by Molina-Molina et al. (2008). For details, see Supplemental Material, "Reporter cell lines and culture conditions."

Protein production and purification. The wt-ER α LBD (wild-type) and the ER α -Y537S LBD mutant (amino acids 302–552) were cloned into a modified pET-15b vector, and the wt-ER β LBD (amino acids 261–502) was cloned into a pET-32a vector. All constructs were expressed in BL21(DE3) cells. Protein domains were purified using a nickel affinity column and size exclusion chromatography, as described in detail in Supplemental Material, "Protein production and purification."

Structure determination. Prior to crystallization assays, the purified ER α -Y537S LBD (final concentration = 0.15 mM) was mixed with 0.3 mM chlordecone, BBP, ferutinine, α -ZA, butylparaben, 4-OP, TCBPA, BP-2, or HPTE, along with 0.3 mM SRC-1 coactivator peptide. Cocrystals were obtained for all complexes in 300–340 mM NaCl, 100 mM HEPES, pH 7.75, and 24–30% polyethylene glycol 3350. Data were collected on the ID23-1, ID23-2, or ID29 beamlines at the European Synchrotron Radiation Facility (Grenoble, France) and processed as described in Supplemental Material, "Data collection and structure determination."

Fluorescence anisotropy measurements. Measurement of the binding affinities of




Figure 1. Chemical structures of the natural agonist E₂, the synthetic antagonist OHT, and the various environmental ER ligands used in the present study.

the fluorescein-labeled SRC-1 NR2 peptide (final concentration of 4 nM) for wild-type ER α and ER β LBDs in both the absence and presence of various ligands was performed using a Safire² microplate reader (TECAN, Männedorf, Switzerland) with the excitation wavelength set at 470 nm and emission measured at 530 nm. The buffer solution for assays was 20 mM Tris, pH 8.0, 150 mM NaCl, 5 mM DTT, and 10% glycerol. All ligands were present at a concentration corresponding to 2 molar equivalents of the highest concentration of protein. The measurements were initiated at the highest concentration of protein (10 μ M for ER α or 20 μ M for ER β). The sample was then diluted successively by a factor of 2, with the buffer containing 8 nM of fluorescent peptide and 20 μ M or 40 μ M of ligand, allowing us to establish the titration curve. Data were analyzed using GraphPad Prism (GraphPad Software Inc., La Jolla, CA, USA).

Results

Compound activities on ER α and ER β .

We monitored the agonistic potential of the compounds using stably transfected HGELN-ER α and -ER β cell lines, allowing for a comparison of the effect of compounds on both human ER subtypes in a similar cellular context. All compounds were first tested on the HGELN parental cell line containing only the reporter gene. We observed some cytotoxicity at ligand concentrations of ≥ 10 μ M but no unspecific modulation of luciferase expression (data not shown). We then characterized the activity of the compounds on HGELN-ER cell lines containing full-length (FL) ER α or ER β . As shown in Figures 2 and 3A, the agonistic potentials depend on the receptor subtype and vary drastically among molecules that range from full agonists to weak agonists or antagonists. Whereas BP-2 acted as a full agonist of both ER subtypes, ferutinine was a selective activator of ER α , and α -ZA efficiently activated both receptors with a slight preference for ER α . The remainder of the compounds can be considered partial agonists with graded effects, with 4-OP being the most active ($\sim 80\%$ of the transactivation seen with E₂) and TCBPA inducing only 17% activity in the HGELN-ER β cell line. Interestingly, some of these compounds displayed different activation capabilities of the two receptor subtypes, as illustrated by TCBPA and chlordecone, which were significantly more efficacious for ER α (50% and 39%, respectively) than for ER β (17% and 19%, respectively). In total, three compounds activated the two ER subtypes equally (BP-2, 4-OP, and BBP), two compounds activated ER β more efficiently than ER α (BPA and butylparaben), and seven compounds activated ER α more efficiently

than ER β (ferutinine, α -ZA, BPC, TCBPA, DDE, chlordecone, and HPTE). The EC₅₀ (median effective concentration) values derived from the transactivation curves suggest that the compounds bound to both ER subtypes with similar affinities (see Supplemental Material, Table S1). To validate this observation, we performed competitive binding assays with (³H)-E₂ (see Supplemental Material, Figure S1 and Table S2), which showed a wide array of affinities ranging from subnanomolar to micromolar values. Together, these experiments show that, in the context of HeLa cells, all the molecules bound to FL-ER α and FL-ER β without subtype selectivity, whereas the functional outcomes of these interactions were, in most cases, subtype specific.

Differential usage of AF-1 and AF-2.

Having characterized the estrogenic potential of the compounds on FL-ER α and FL-ER β , we performed additional cell-based experiments aimed at assessing the relative contribution of the ERs' AF-1 and AF-2 to this activity. We examined the agonistic properties of compounds using HELN cells stably transfected with ERs that had their N-terminal AB (AF-1) region deleted (Δ AB-ER α and Δ AB-ER β). Interestingly, we observed that deletion of the AB domains reduced the xenoestrogen-induced transcriptional activity of ERs, with the effect being remarkably more pronounced for ER β (Figure 2, and compare Figure 3A and 3B). Correlation diagrams representing the maximum activity of FL-ERs versus Δ AB-ERs in the presence of the various ligands clearly illustrate the differential involvement of the two AFs in ER α and ER β (see Supplemental Material, Figure S2). Among the 12 compounds tested, only 2 displayed a fully AF-1-independent activity. These 2 compounds are BP-2, which acted as a full agonist of the entire or truncated ER forms, and more surprisingly, chlordecone, whose partial agonist activity on ER α also relied exclusively on the AF-2. Comparing the decrease in the agonistic potential of the other compounds upon AF-1 removal in ER α and ER β , we observed that most of them retained a significant degree of activity in HGELN- Δ AB-ER α but not in HGELN- Δ AB-ER β cell lines, where the majority of the ER ligands become partially or completely inactive. Together, these data suggest that, in the context of HeLa cells, the environmental compounds bind to ER α and ER β with similar affinities but modulate the transcriptional activity of the two subtypes through different combinations of AF-1 and AF-2, with AF-1 being preeminent in the EDC-induced transcriptional activity of ER β .

Subtype-specific AF-2 modulation by exogenous ER ligands. To evaluate whether the differential involvement of the N- and C-terminal AFs is dependent on the cellular

context or a consequence of intrinsic differences between the two ER subtypes, we used fluorescence anisotropy to monitor the recruitment of a fluorescein-labeled peptide derived from the coactivator SRC-1 by the purified ER α - or ER β -LBDs in either the absence (apo-receptor) or the presence of compounds. This experimental setup allows measuring the influence of a given ligand on a receptor's AF-2 in purely *in vitro* conditions. In keeping with their agonistic or antagonistic activities, E₂ and OHT, respectively, strongly enhanced and decreased the binding affinity of SRC-1 to both ERs, with E₂ being slightly more efficient on ER α (Figure 3C; see also Supplemental Material, Figure S3 and Table S3). The values of the dissociation constants (K_d) obtained in the presence of the various compounds show that based on their impact on the recruitment of the coactivator-derived peptide by ERs, the ligands can be partitioned into three classes. The first class corresponds to strong agonists ($K_d \leq 1$ μ M) and includes α -ZA, ferutinine, and BP-2 for ER α but only BP-2 for ER β . The second class comprises the partial agonists (1 μ M $\leq K_d \leq$ values for apo-ER α or -ER β) and contains most of the remaining molecules for ER α (4-OP, BPA, butylparaben, BBP, TCBPA, chlordecone, and DDE), but only α -ZA, 4-OP, BPA, and butylparaben for ER β . Finally, the third class encompasses the antagonists ($K_d \geq$ values for apo-ER α or -ER β): BPC and HPTE for ER α , and ferutinine, BBP, BPC, TCBPA, chlordecone, HPTE, and DDE for ER β . These fluorescence anisotropy data are in good agreement with the transactivation assays because all of the activity profiles measured in HGELN- Δ AB-ER cell lines (Figure 3B) fell into these three categories of ligands, confirming that the compounds activated ER α AF-2 more efficiently than ER β AF-2, which appears more prone to antagonism. Thus, binding of environmental ligands imposed different structural constraints on the LBDs of the two ER subtypes, which most likely account for the differential contribution of both AFs in ER α and ER β .

Structural analysis. To gain structural insights into the mechanisms by which exogenous ligands bind to and activate ERs, we solved the crystal structures of ER α LBD in complex with the various compounds. Because ER α - and ER β -LBDs share a high degree of homology in their amino acid sequence and are very similar in their ternary architecture, we reasoned that structural information regarding interaction with ER β could be acquired through molecular modeling using the ER α LBD structures. To allow the crystallization of ER α bound to the environmental molecules acting essentially as partial agonists that are unable to induce a stable conformation of the LBD, we used the ER α -Y537S LBD mutant described previously by Nettles




Figure 2. The relative activity of xenoestrogens relies on their different binding modes. (Left) Dose–response curves corresponding to the HGELN-ER α , - Δ AB-ER α , -ER β , and - Δ AB-ER β , luciferase assays of E_2 and xenoestrogens. The maximal activity (100%) was obtained with 10 nM E_2 ; values are mean \pm SD from three separate experiments. (Right) The interaction networks of E_2 and xenoestrogens with LBD residues of ER α . Key for structures: red, oxygen; blue, nitrogen; cyan, carbon; yellow, sulfur; green, chlorine; black dashed lines, hydrogen bonds; red spheres, water molecules. The electron density represents a $F_o - F_c$ simulated annealing omit map contoured at 3 σ .

et al. (2008). In several independent studies, the Y537S surface mutation has been shown to stabilize the active conformation of the receptor, and, in turn, facilitate crystallization of weak agonists without modifying either the overall architecture of the LBD or the binding mode of ligands (Bruning et al. 2010; Delfosse et al. 2012; Nettles et al. 2008). To further stabilize the ER α -Y537S LBD in its active form, a peptide containing the second interaction motif of the coactivator SRC-1 was also added during the crystallization trials. Details of the structure determination and refinement are summarized in Supplemental Material, Table S4.

The structures display the canonical active conformation, with helix H12 capping the ligand-binding pocket (LBP) and the SRC-1 peptide bound to the "AF-2 surface" formed by helices H3, H5, and H12 (Figure 4A). Most compounds could be precisely placed in their respective electron density, revealing different binding modes (Figure 2). Some ligands, such as BP-2, α -ZA, BPA, and

TCBPA, adopted a binding mode reminiscent of that used by E₂, with two phenol groups hydrogen-bonded to three polar residues located at the two ends of the LBP, namely H524 (H11) on one side and E353 (H3) and R394 (H6) on the other side. However, we also noticed significant differences in the geometry of the interactions between H524, E353, R394, and the hydroxyl moieties of E₂ and the ligands with possible functional and/or binding implications. Indeed, none of the compounds recapitulated the exact hydrogen bond network seen in the E₂-containing complex. The remaining contacts involved essentially van der Waals interactions, the number of which varies from one compound to another and accounts in part for the various binding affinities of the ligands. Several compounds did not interact with H524 because they lack a second hydroxyl group (ferutinine, 4-OP, butylparaben, and DDE) or because they adopt a position that draws this hydroxyl moiety toward T347 in H3 (BPC, HPTE, and DDE). Finally, two compounds,

BBP and chlordecone, were not engaged in any direct interaction with either of these polar residues, the latter being indirectly hydrogen bonded to E353 via water molecules. As shown in Figure 2, the position of DDE could not be precisely determined because of the absence of electron density for some regions of the ligand. This poorly defined electron density reflects a higher dynamic for DDE. Finally, the docking of two BBP molecules with distinct positions was necessary to fully account for the observed electron density, indicating that this molecule can adopt two alternate orientations in the LBP.

Structural basis for compound actions in ER α . We next considered how these different binding modes may account for the various activity profiles of compounds toward ER α . Superposition of our 13 structures onto the structure in complex with E₂ revealed that the regions of the LBP occupied by the ligand vary greatly from one compound to another. BP-2, which displayed the highest agonistic activity among the molecules used in this




Figure 3. Differential involvement of AFs in ERs. HGELN-ERs cells (A) and HGELN- Δ AB-ERs cells (B) were incubated with 10 nM E₂ or 10 μ M ER exogenous ligands. The maximal luciferase activity (100%) was obtained with 10 nM E₂; values are the mean \pm SD from three separate experiments. (In B, the horizontal dotted lines highlight the partition of the ligands into three classes based on fluorescence anisotropy data. (C) Fluorescence anisotropy data showing the relative affinity of the SRC-1 NR2 peptide for ER α LBD or ER β LBD in the absence of ligand or in the presence of saturating concentrations of E₂ or xenoestrogens. Ligands are classified as agonists ($K_d \leq 1 \mu$ M), partial agonists (1μ M $\leq K_d \leq$ values for apo-ER α or -ER β), or antagonists ($K_d \geq$ values for apo-ER α or -ER β).

study, occupies almost the same volume as E₂ in the LBP and therefore imposes similar side chain conformations (Figure 4B). In particular, it stabilizes the same conformer of H524 as that seen in the E₂-bound ER α . We and others have previously shown that this residue is involved in a key hydrogen bond network including residues from H3 and H11, which are part of the docking site maintaining H12 in the active position (Bruning et al. 2010; Delfosse et al. 2012; Nettles et al. 2008). α -ZA occupies a bit more space than E₂, especially in the H8 region on one side and the H3, H11, and H12 region on the other side (Figure 4C). Interestingly, ER α accommodates this larger compound with only minimal LBP rearrangements. In contrast, a number of large side chain conformational changes are necessary to accommodate the ferutinine. In particular the bulky methyl-cycloheptene ring pointing toward H8 provokes the enlargement of a small preexisting hydrophobic cavity via the reorientations of M421 and F425 in H8, as well as F404 in the β -strand S1 (Figure 4D). This structure further highlights the plasticity of this particular region of ER α LBP (Nettles et al. 2007), which can accommodate bulky ligands within the confined environment of the active conformation while retaining a substantial level of agonistic activity (Figure 2).

In contrast with the aforementioned compounds, ligands with partial AF-2 agonist activity are generally smaller in size (4-OP, BPA, butylparaben, chlordecone) and/or are more flexible (4-OP, butylparaben, BBP). As a consequence, they are less efficient in stabilizing the LBP side chain conformations required to hold the active form of the receptor. The loss of stabilizing contacts triggered by bisphenols and their functional outcome has previously been discussed in detail (Delfosse et al. 2012). The suboptimal interaction between ER α and the weak compounds reported in this study is exemplified by the structure of the butylparaben-containing complex, where several previously recognized ligand-H11 stabilizing interactions are missing (Figure 4E). We have previously shown that the two chlorine atoms of BPC prevent this compound from adopting the same position as BPA in the ER α LBP (Figure 2) and that the AF-2 antagonistic character of BPC is a direct consequence of this particular binding mode. Indeed, the phenol ring, which adopts a different orientation in BPC, induces a 180° rotation of T347 (H3) and forms a hydrogen bond with the hydroxyl group of this residue. The complete reorientation of the T347 side chain provokes the disruption of a cluster of hydrophobic interactions, which hold together H3, H11, and the loop preceding H12, thereby destabilizing the AF-2 surface. It is noteworthy

that the closely related HPTE adopts a similar binding mode and acts as an AF-2 antagonist, whereas DDE, which contains chlorine atoms instead of hydroxyl groups, does not induce the reorientation of T347 and acts as a partial agonist (Figure 4F). Together, these experimental data provide a structural rationale to explain the functional properties of the environmental ligands of ERs at a near-atomic level.

Structural basis for subtype-specific action of ER ligands. We subsequently looked for a rational explanation for the differential activation of ER α and ER β AF-2 by the compounds (Figure 3B). The LBDs of the two ER subtypes share a high degree of homology

in their primary sequence. Notably, there are only two conservative residue substitutions in the LBPs of the two receptors. These substitutions correspond to the replacement of L384 (ER α) by M336 (ER β) in H6 and M421 (ER α) by I373 (ER β) in H8, which have been shown previously to account, at least in part, for the subtype-specific action of ER ligands (Nettles et al. 2004). Inspection of the crystal structures of both ER subtypes in complex with E₂ shows that the variable amino acids reside on each side of the C and D rings of E₂ and create different space constraints in this portion of ER α and ER β LBPs (see Supplemental Material, Figure S4). Indeed, superposition of our ER α or ER β structures




Figure 4. Xenooestrogens use diverse binding modes. (A) The entire structure of the ER α LBD in complex with E₂ and SRC-1 coactivator peptide (yellow); the structure shows the AF-2 surface formed by helices H3, H5, and H12 (green), and the lower part of the LBD (blue) encloses the ligand-binding pocket (LBP). (B–E) Interaction networks of BP-2 (B; pink), α -ZA (C; orange), ferutinine (D; green), and butylparaben (E; purple) with residues of the LBP compared with that of E₂ (gray). In (E), red dashed lines represent the interactions lost in the butylparaben complex structure. (F) HPTE and DDE adopt the orientation previously observed for BPC allowing HPTE to interact with residue T347. This position results in the disruption of the hydrophobic network involving helices H3 and H11 and the loop preceding H12, thereby destabilizing the AF-2 surface. Color code: red, oxygen; blue, nitrogen; cyan, carbon; yellow, sulfur; green, chlorine; dashed lines, hydrogen bonds and hydrophobic interactions.

contained in the Protein Data Bank (PDB) reveals that in ER α , M421 (H8) can adopt a wide array of conformations to accommodate the ligands, whereas in ER β , M336 (H6) is much less flexible due to strong steric constraints provided by surrounding residues (see Supplemental Material, Figure S4). Thus, with two bulky and rigid residues in H6 (M336) and H8 (I373), ER β might be more sensitive than ER α to variations in the size of the bound ligand.

Accordingly, our structures reveal that most of the compounds insert a bulky feature in this region of the LBP and exhibit a marked subtype-dependent activity as illustrated by ferutinine and BBP (Figures 5 and 3B). Both molecules contain a bulky group that projects toward ER α M421, which, in turn, must undergo a large conformational change. In ER β , the linear amino acid M421 is replaced by the branched residue I373, which is unable to move away from the pocket and make room for the ligands. A likely consequence is that I373 induces a shift of the ligands toward H12, thus lowering the interaction of H12 in the active conformation with the LBD surface and accounting for the weakest agonistic activity of the compounds in ER β (Figure 5). Obviously the strength of the steric constraints applied to ER β H12 varies according to the chemical structure of the bound ligand, as reflected by the graded partial agonistic/antagonistic activity of the compounds (Figure 3B). Finally, it is noteworthy that the weak ligand-induced ER β AF-2 activity can be partially or completely compensated by the N-terminal activation domain, thus confirming the preeminent functional role of ER β AF-1 in HeLa cells (Figure 3A,B).

Discussion

Deregulation of NR-mediated transcription accounts for the deleterious effects of many EDCs. Thus, characterization of the interaction between receptors and environmental compounds both at the structural and functional levels, as well as the development of robust *in vivo*, *in vitro*, and *in silico* screening methods, are important for the assessment of the global hormonal activity of a large number of chemicals (Janosek et al. 2006). In this context, we have used a combination of complementary biochemical, structural, biophysical, and cell-based approaches to provide a mechanistic view for how members of the most recognized pollutant families bind to and activate ER α and ER β , two of their primary biological targets.

We observed that the compounds bind to both ER subtypes with similar affinities but modulate the transcriptional activity of ER α and ER β in different manners. Using HeLa cells stably transfected with Δ AB-ERs and fluorescence anisotropy measurements

with purified ER LBDs, we showed that the environmental compounds activated the C-terminal AF-2 of ER α more efficiently than that of ER β . However, when the N-terminal AF-1 was present (FL-ERs), the global activity of ER β was partially (α -ZA, BPC, DDE, HPTE) or completely (BP-2, 4-OP, BBP) restored compared with the results obtained with ER α , in which two ligands (BPA and butylparaben) activated FL-ER β more efficiently than FL-ER α . In contrast, ferutinine, TCBP, and chlordecone acted as ER β antagonists in both the FL and Δ AB constructs. These data clearly show that the environmental ligands acted in a subtype-specific fashion as full agonists, partial agonists, or antagonists by using different combinations of the N- and C-terminal activation functions of ER α and ER β , the AF-1 being dominant in the latter. They also suggest that the binding of structurally diverse molecules induced specific ER LBD structures and/or dynamics with divergent impacts on AF-1 activity (compare BBP and chlordecone on ER β in Figure 3A,B). However, the precise structural basis of this interdomain communication is still unknown because no three-dimensional (3D) structure of an entire NR has yet been obtained.

The relative agonist/antagonist activity of a given ER ligand is dependent on several parameters, including the nature of the target gene and cell type considered, the latter being critical due to variation in the equipment of transcription coregulators that finely tune the activity of the two AFs of the ERs (McDonnell et al. 2002). Such considerations might explain, for instance, why HPTE is characterized here as an antagonist but reported to display agonistic properties

in other studies (Wilson et al. 2004). In a similar manner, Li et al. (2013) observed that chlordecone activated ER α in HeLa cells, but it was unable to do so in HepG2 cells. It would be therefore interesting to extend the present study by comparing the effects of the various ER ligands on ER activities in different cell types and on the recruitment of several coregulatory proteins. In terms of biological activity, environmental estrogens are comparable to the selective estrogen receptor modulators (SERMs) such as raloxifen and tamoxifen (which also act as partial agonists) and selectively block estrogen action in the breast but not in other tissues, such as bones (Nilsson and Gustafsson 2011). As observed for these SERMs, it appears quite obvious that xenoestrogens presenting partial agonism will manifest different activity depending on the tissues. In line with this, further work will be needed to define the deleterious effects linked to the selective partial agonistic activity of environmental chemicals on key physiological processes regulated by ERs such as bone and metabolic homeostasis, vasomotor symptoms, depression, or neurodegenerative diseases (Nilsson and Gustafsson 2011).

It is well established that ER α is the major driver of the proliferative effects of estrogens in cancers of both breast and ovary, as well as in normal reproductive tissues, with ER β serving largely as a brake for ER α -driven proliferation (Bossard et al. 2012; Madak-Erdogan et al. 2013; Paruthiyil et al. 2004; Strom et al. 2004). Indeed, in ovarian cancer BG1 cells expressing both ERs, we previously observed that ER α -selective agonists activated cell proliferation more efficiently than did ER β -selective ligands or ER pan-agonists (Docquier et al.




Figure 5. Methionine 421 confers plasticity and adaptability to ER α LBP. Structure superposition of E₂-bound ER β LBD (yellow) with (A) ferutinine-bound ER α LBD (green), or (B) BBP-bound ER α LBD (gray). The presence of I373 in ER β instead of M421 in ER α will induce a shift of bulky ligands toward helix H12 thus lowering the stability of the AF-2.

2013). This observation, together with our finding that ER environmental ligands may act in a subtype-specific fashion, suggests that different xenoestrogens could have different impacts on cancer incidence. In this regard, one can predict that pollutants acting as ER α agonists and ER β antagonists in a particular cellular context (e.g., ferutinine in HeLa cells) should stimulate cell proliferation and tumor growth more effectively than compounds activating more ER β than ER α .

Finally, our crystallographic analysis revealed the various mechanisms by which distantly related chemicals bind to and activate ERs. These data will increase the effectiveness of 3D structure-based computational tools aimed at predicting the NR-mediated activity of environmental pollutants (Delfosse et al. 2012; Vuorinen et al. 2013). In addition to providing a better understanding of the differential activities, binding affinities, and specificities of environmental ER ligands, the structures provide rational guidelines for the design of safer chemicals.

REFERENCES

- Bossard C, Busson M, Vindrieux D, Gaudin F, Machelon V, Brigitte M, et al. 2012. Potential role of estrogen receptor beta as a tumor suppressor of epithelial ovarian cancer. *PLoS one* 7(9):e44787; doi:10.1371/journal.pone.0044787.
- Bruning JB, Parent AA, Gil G, Zhao M, Nowak J, Pace MC, et al. 2010. Coupling of receptor conformation and ligand orientation determine graded activity. *Nat Chem Biol* 6(11):837–843.
- Couse JF, Korach KS. 1999. Estrogen receptor null mice: what have we learned and where will they lead us? *Endocr Rev* 20(3):358–417.
- Curtis SW, Washburn T, Sewall C, DiAugustine R, Lindzey J, Couse JF, et al. 1996. Physiological coupling of growth factor and steroid receptor signaling pathways: estrogen receptor knockout mice lack estrogen-like response to epidermal growth factor. *Proc Natl Acad Sci USA* 93(22):12626–12630.
- De Coster S, van Larebeke N. 2012. Endocrine-disrupting chemicals: associated disorders and mechanisms of action. *J Environ Public Health* 2012:713696; doi:10.1155/2012/713696.
- Delfosse V, Grimaldi M, Pons JL, Boulahtouf A, le Maire A, Cavailles V, et al. 2012. Structural and mechanistic insights into bisphenols action provide guidelines for risk assessment and discovery of bisphenol A substitutes. *Proc Natl Acad Sci USA* 109(37):14930–14935.
- Diamanti-Kandaraki E, Bourguignon JP, Giudice LC, Hauser R, Prins GS, Soto AM, et al. 2009. Endocrine-disrupting chemicals: an Endocrine Society scientific statement. *Endocr Rev* 30(4):293–342.
- Docquier A, Garcia A, Savatier J, Boulahtouf A, Bonnet S, Bellet V, et al. 2013. Negative regulation of estrogen signaling by ER β and RIP140 in ovarian cancer cells. *Mol Endocrinol* 27(9):1429–1441.
- Grun F, Blumberg B. 2007. Perturbed nuclear receptor signaling by environmental obesogens as emerging factors in the obesity crisis. *Rev Endocr Metab Disord* 8(2):161–171.
- Janosek J, Hilscherova K, Blaha L, Holoubek I. 2006. Environmental xenobiotics and nuclear receptors—interactions, effects and in vitro assessment. *Toxicol In Vitro* 20(1):18–37.
- Kobayashi Y, Kitamoto T, Masuhiro Y, Watanabe M, Kase T, Metzger D, et al. 2000. p300 mediates functional synergism between AF-1 and AF-2 of estrogen receptor α and β by interacting directly with the N-terminal A/B domains. *J Biol Chem* 275(21):15645–15651.
- le Maire A, Grimaldi M, Roecklin D, Dagnino S, Vivat-Hannah V, Balaguer P, et al. 2009. Activation of RXR-PPAR heterodimers by organotin environmental endocrine disruptors. *EMBO Rep* 10(4):367–373.
- Li Y, Luh CJ, Burns KA, Aroy Y, Jiang Z, Teng CT, et al. 2013. Endocrine-disrupting chemicals (EDCs): *in vitro* mechanism of estrogenic activation and differential effects on ER target genes. *Environ Health Perspect* 121:459–466; doi:10.1289/ehp.1205951.
- Lindberg MK, Movérare S, Skrtic S, Gao H, Dahlman-Wright K, Gustafsson JA, et al. 2003. Estrogen receptor (ER)- β reduces ER α -regulated gene transcription, supporting a “Ying Yang” relationship between ER α and ER β in mice. *Mol Endocrinol* 17(2):203–208.
- Madak-Erdogan Z, Charn TH, Jiang Y, Liu ET, Katzenellenbogen JA, Katzenellenbogen BS. 2013. Integrative genomics of gene and metabolic regulation by estrogen receptors α and β , and their coregulators. *Mol Syst Biol* 9:676; doi:10.1038/msb.2013.28.
- McDonnell DP, Wardell SE. 2010. The molecular mechanisms underlying the pharmacological actions of ER modulators: implications for new drug discovery in breast cancer. *Curr Opin Pharmacol* 10(6):620–628.
- McDonnell DP, Wijayarathne A, Chang CY, Norris JD. 2002. Elucidation of the molecular mechanism of action of selective estrogen receptor modulators. *Am J Cardiol* 90(1A):35F–43F.
- Métivier R, Penot G, Flouriot G, Pakdel F. 2001. Synergism between ER α transactivation function 1 (AF-1) and AF-2 mediated by steroid receptor coactivator protein-1: requirement for the AF-1 α -helical core and for a direct interaction between the N- and C-terminal domains. *Mol Endocrinol* 15(11):1953–1970.
- Molina-Molina JM, Escande A, Pillon A, Gomez E, Pakdel F, Cavailles V, et al. 2008. Profiling of benzophenone derivatives using fish and human estrogen receptor-specific *in vitro* bioassays. *Toxicol Appl Pharmacol* 232(3):384–395.
- Nettles KW, Bruning JB, Gil G, Nowak J, Sharma SK, Hahn JB, et al. 2008. NFKB selectivity of estrogen receptor ligands revealed by comparative crystallographic analyses. *Nat Chem Biol* 4(4):241–247.
- Nettles KW, Bruning JB, Gil G, O'Neill EE, Nowak J, Guo Y, et al. 2007. Structural plasticity in the oestrogen receptor ligand-binding domain. *EMBO Rep* 8(6):563–568.
- Nettles KW, Sun J, Radek JT, Sheng S, Rodriguez AL, Katzenellenbogen JA, et al. 2004. Allosteric control of ligand selectivity between estrogen receptors α and β : implications for other nuclear receptors. *Mol Cell* 13(3):317–327.
- Newbold RR, Jefferson WN, Padilla-Banks E. 2009. Prenatal exposure to bisphenol A at environmentally relevant doses adversely affects the murine female reproductive tract later in life. *Environ Health Perspect* 117:879–885; doi:10.1289/ehp.0800045.
- Nilsson S, Gustafsson JA. 2011. Estrogen receptors: therapies targeted to receptor subtypes. *Clin Pharmacol Ther* 89(1):44–55.
- Paruthiyil S, Parmar H, Kerekatte V, Cunha GR, Firestone GL, Leitman DC. 2004. Estrogen receptor β inhibits human breast cancer cell proliferation and tumor formation by causing a G₂ cell cycle arrest. *Cancer Res* 64(1):423–428.
- Riu A, Grimaldi M, le Maire A, Bey G, Phillips K, Boulahtouf A, et al. 2011. Peroxisome proliferator-activated receptor γ is a target for halogenated analogs of bisphenol A. *Environ Health Perspect* 119:1227–1232; doi:10.1289/ehp.1003328.
- Rubin BS, Soto AM. 2009. Bisphenol A: perinatal exposure and body weight. *Mol Cell Endocrinol* 304(1–2):55–62.
- Schug TT, Janesick A, Blumberg B, Heindel JJ. 2011. Endocrine disrupting chemicals and disease susceptibility. *J Steroid Biochem Mol Biol* 127(3–5):204–215.
- Strom A, Hartman J, Foster JS, Kietz S, Wimalasena J, Gustafsson JA. 2004. Estrogen receptor β inhibits 17 β -estradiol-stimulated proliferation of the breast cancer cell line T47D. *Proc Natl Acad Sci USA* 101(6):1566–1571.
- Vuorinen A, Odermatt A, Schuster D. 2013. *In silico* methods in the discovery of endocrine disrupting chemicals. *J Steroid Biochem Mol Biol* 137:18–26.
- Weihua Z, Saji S, Makinen S, Cheng G, Jensen EV, Warner M, et al. 2000. Estrogen receptor (ER) β , a modulator of ER α in the uterus. *Proc Natl Acad Sci USA* 97(11):5936–5941.
- Wilson VS, Bobseine K, Gray LE Jr. 2004. Development and characterization of a cell line that stably expresses an estrogen-responsive luciferase reporter for the detection of estrogen receptor agonist and antagonists. *Toxicol Sci* 81(1):69–77.