

## Using remotely sensed temperature to estimate climate response functions

This content has been downloaded from IOPscience. Please scroll down to see the full text.

2017 Environ. Res. Lett. 12 014013

(<http://iopscience.iop.org/1748-9326/12/1/014013>)

View [the table of contents for this issue](#), or go to the [journal homepage](#) for more

Download details:

IP Address: 210.77.64.106

This content was downloaded on 30/03/2017 at 10:58

Please note that [terms and conditions apply](#).

You may also be interested in:

[The impact of climate extremes and irrigation on US crop yields](#)

T J Troy, C Kipgen and I Pal

[Implications of regional improvement in global climate models for agricultural impact research](#)

Julian Ramirez-Villegas, Andrew J Challinor, Philip K Thornton et al.

[Placing bounds on extreme temperature response of maize](#)

Christopher J Anderson, Bruce A Babcock, Yixing Peng et al.

[Intensity of heat stress in winter wheat—phenology compensates for the adverse effect of global warming](#)

Ehsan Eyshi Rezaei, Stefan Siebert and Frank Ewert

[Water resources transfers through southern African food trade: water efficiency and climate signals](#)

Carole Dalin and Declan Conway

[More uneven distributions overturn benefits of higher precipitation for crop yields](#)

Ram Fishman

[Particulate matter concentration mapping from MODIS satellite data: a Vietnamese case study](#)

Thanh T N Nguyen, Hung Q Bui, Ha V Pham et al.

[Simulating the impacts of chronic ozone exposure on plant conductance and photosynthesis, and on the regional hydroclimate using WRF/Chem](#)

Jialun Li, Alex Mahalov and Peter Hyde

[Drought effects on US maize and soybean production: spatiotemporal patterns and historical changes](#)

Samuel C Zipper, Jiangxiao Qiu and Christopher J Kucharik

## Environmental Research Letters



## LETTER

## Using remotely sensed temperature to estimate climate response functions

## OPEN ACCESS

## RECEIVED

29 July 2016

## REVISED

9 December 2016

## ACCEPTED FOR PUBLICATION

19 December 2016

## PUBLISHED

17 January 2017

Sam Heft-Neal<sup>1,3</sup>, David B Lobell<sup>1,2</sup> and Marshall Burke<sup>1,2</sup><sup>1</sup> Dept. of Earth System Science, Stanford University, United States<sup>2</sup> Center on Food Security and the Environment, Stanford University, United States<sup>3</sup> Author to whom any correspondence should be addressed.E-mail: [sheftneal@stanford.edu](mailto:sheftneal@stanford.edu)**Keywords:** climate response functions, remote sensing, surface temperature, weather stations, agriculture, air temperature

Original content from this work may be used under the terms of the [Creative Commons Attribution 3.0 licence](https://creativecommons.org/licenses/by/4.0/).

Any further distribution of this work must maintain attribution to the author(s) and the title of the work, journal citation and DOI.



**Abstract**

Temperature data are commonly used to estimate the sensitivity of many societally relevant outcomes, including crop yields, mortality, and economic output, to ongoing climate changes. In many tropical regions, however, temperature measures are often very sparse and unreliable, limiting our ability to understand climate change impacts. Here we evaluate satellite measures of near-surface temperature ( $T_s$ ) as an alternative to traditional air temperatures ( $T_a$ ) from weather stations, and in particular their ability to replace  $T_a$  in econometric estimation of climate response functions. We show that for maize yields in Africa and the United States, and for economic output in the United States, regressions that use  $T_s$  produce very similar results to those using  $T_a$ , despite the fact that daily correlation between the two temperature measures is often low. Moreover, for regions such as Africa with poor station coverage, we find that models with  $T_s$  outperform models with  $T_a$ , as measured by both  $R^2$  values and out-of-sample prediction error. The results indicate that  $T_s$  can be used to study climate impacts in areas with limited station data, and should enable faster progress in assessing risks and adaptation needs in these regions.

**1. Introduction**

Historical data on climatic variables such as temperature and precipitation are key for understanding how human and natural systems respond to climatic change. While many global-scale gridded weather datasets do exist for this purpose [1, 2] and have provided fundamental insights into climatic responses, accuracies are often limited by the underlying station data availability which can vary substantially over time and space. For instance, according to our measure of quality, defined as stations with at least 10 years of data and missing less than 30% of daily observations, high-quality station density in the Global Historical Climatology Network (GHCN) database peaked in Africa in 1976 and peaked globally in 2001 (figure 1). By 2010 the database contained just 215 high-quality weather stations in all of Africa. This combination of low spatial density of stations, and stations that go on and offline at different times, can lead to substantial measurement error in interpolated datasets which in turn can bias estimates of societal impacts [3].

An alternate and less-common approach is to use satellites rather than ground-based measures to study climate variables of interest. For instance, several satellites measure surface emission of thermal energy, which can be converted into estimates of surface skin temperature ( $T_s$ )—a product that the Moderate Resolution Imaging Spectroradiometer (MODIS) has provided at 1 km resolution daily for over a decade. Past studies have evaluated agreement between MODIS and weather stations on daily time scales, often finding weak correlations for daytime temperatures because factors other than  $T_a$ , such as cloudiness and soil moisture, can affect  $T_s$  [4–6]. However, these results could be of limited relevance for estimating how societal outcomes respond to climatic change, since estimates of societal impacts often rely on year-to-year variations in seasonally aggregated measures of temperature exposure, and correlations between station and satellite data tend to increase as the period of aggregation lengthens. For instance, in the United States, the  $R^2$  value associated with regressing daytime  $T_s$  on maximum  $T_a$  is 30% higher for seasonal averages than for 8-day averages (figure 2).



Direct evaluation of  $T_s$  in societal applications thus appears warranted. Although a previous study evaluated  $T_s$  in cross-sectional regressions [7], most econometric studies rely on time-variation to identify climate response functions.

Another motivation for using  $T_s$  is that it may be a more direct measurement of the relevant temperature for certain applications. In agricultural settings,  $T_s$  measures canopy temperature, and the deviation of canopy temperature from  $T_a$  is often used as an indicator of plant water stress for drought monitoring or irrigation scheduling [8, 9].  $T_s$  may therefore better represent environmental conditions for predicting

crop yields than  $T_a$ , as illustrated for wheat experiments in Europe [10].

## 2. Methods

To better understand how satellite-based temperature models could inform our understanding of societal responses to climatic change, we revisited three previous studies that had used standard measures of  $T_a$  to study impacts: maize yields in Africa [11], maize yields in the US [12], and gross domestic product (GDP) in the United States [13]. In each of these

Table 1. Overview of data sources.

Variable	Data	Details
<b>Surface Temperature (<math>T_s</math>)</b>	(Africa and U.S.) MODIS Aqua MYD11C2 v5	8-day composited average day and night land surface temperatures, released as $0.05^\circ \times 0.05^\circ$ grids, available July 2002–present [17].
<b>Air Temperature (<math>T_a</math>) and Precipitation</b>	(Africa) Interpolated ground stations	Daily minimum and maximum temperatures and precipitation for each field trial were estimated by interpolation of daily measurements made in the World Meteorological Organization, World Weather Watch Program, available 1999–2007 [11].
	(U.S.) PRISM Climate Group, Oregon State University	Climatologically-aided interpolation (CAI) from ground weather stations carried out by the PRISM group and released as $2.5' \times 2.5'$ daily grids, available 1981–present [14].
<b>Maize Yields</b>	(Africa) Field trial data	Georeferenced data from more than 25 000 maize experimental field trials across Eastern & Southern Africa, available 1999–2007 [11]. Paper analysis draws on 15 164 trials since 2003.
	(U.S.) United States Department of Agriculture National Agricultural Statistics Service	County-year maize yield data, available 1910–2015. Paper analysis draws on 12 103 observations from 931 maize producing counties between 2003 and 2014.
<b>Per-capita GDP</b>	(U.S.) Bureau of Economic Analysis	County-year per-capita GDP from the Local Area Personal Income Accounts data set, available 1969–2015 [18]. Paper analysis draws on 34 718 observations from 2 747 counties between 2003 and 2014.

studies, we re-estimated relationships using both  $T_a$  and  $T_s$  and compared model performance across temperature measures.

We utilized MODIS Aqua MYD11C2 (8-day) version 5 products as our estimates of land surface temperature ( $T_s$ ) for all three analyses (see table 1 for more information on data sources). MODIS 8-day composited averages have a resolution of  $0.05^\circ$  ( $5.6$  km) and are available from mid 2002 to the present. We used  $T_s$  estimates from the Aqua satellite because it captures images at approximately 1:30 AM and PM local time which more closely approximates the timing of daily temperature extremes than the Terra satellite schedule (10:30 AM and PM) [4]. Missing observations were replaced with inverse distance weighted averages of the nearest four non-missing cells.

For each analysis,  $T_s$  measures were constructed analogously to the  $T_a$  measures that had been used in the previous studies. The Africa analysis drew on more than 15 000 historical maize trials including 12 500 fields under optimal management and 2 500 fields under drought management conditions.  $T_a$  and precipitation data were previously interpolated from publicly available daily weather station data using thin-plate splines [11]. Following this previous work, we estimated a fixed-effects model with quadratic functions of maximum temperature and total precipitation averaged over field-specific 150-day growing seasons:

$$Y_{ist} = T\max_{ist} + T\max_{ist}^2 + Pr_{ist} + Pr_{ist}^2 + \gamma_s + \delta_t + \varepsilon_{it} \quad (1)$$

where  $Y_{ist}$  is the natural logarithm of reported maize yield for the  $i$ th trial at field station  $s$  in year  $t$ ,  $T\max$  is maximum temperature averaged over the 150 days following planting,  $Pr$  is total precipitation around anthesis,  $\gamma$  is a field site fixed effect,  $\delta$  is a year fixed


effect, and  $\varepsilon$  is an error term. For each field site,  $T_s$  observations were constructed by taking the inverse-distance weighted average of the nearest 9 MODIS cells. These values were then averaged over the 150 day growing period at each field site. The precipitation values interpolated from weather stations were included in both  $T_a$  and  $T_s$  regressions.

The analogous analysis in the United States drew on more than 12 000 county-year maize yield observations from USDA's National Agricultural Statistics Service and the PRISM data set that consists of high-resolution gridded daily maximum temperature and precipitation [14]. Regression analysis of temperature impacts on US maize took the form:


$$Y_{it} = T\max_{it} + T\max_{it}^2 + JulyMax_{it} + Pr_{it} + \gamma_i + \delta_t + \varepsilon_{it} \quad (2)$$

where  $Y_{it}$  is log yield in county  $i$  and year  $t$ ,  $T\max_{it}$  is the maximum temperature averaged over the approximate three month maize growing season (JJA),  $JulyMax$  is the maximum temperature averaged across July,  $Pr$  is the total precipitation across the growing season,  $\gamma$  is a county fixed effect,  $\delta$  is a year fixed effect, and  $\varepsilon$  is an error term. We estimated this simple specification in order to facilitate tractable comparison across temperature metrics and because model performance was similar between our model and more flexible growing degree models, such as used in [12, 15]. For both  $T_s$  and  $T_a$ , grid cells were spatially aggregated to the county level using agricultural area weights and temporally averaged over JJA and July. The same PRISM precipitation values were used in both the  $T_a$  and  $T_s$  regressions.

For temperature-GDP relationships in the United States, following [13] we utilized nearly 35 000 county-year observations of GDP from the Bureau of



**Figure 3.** Comparing air and surface temperature response functions for US and African maize. Panels (a) and (b) show results from the estimation of equation 1 drawing on more than 15 000 African field trials carried out between 2002 and 2007. Panel (c) shows results from the estimation of equation 2, an analogous analysis for the United States, based on more than 12 000 county-year observations spanning 2003–2014. Models with  $T_s$  perform better than models with  $T_a$  in Africa as measured by both RMSE and  $R^2$ . In order to plot air and surface temperature on the same axis, surface temperature units were transformed into air temperature units using the quantile mapping procedure described in Methods.



**Figure 4.** Per-capita GDP response to temperature in the United States. Results from estimating equation 3 following the analysis in [13]. Coefficients measure the effect of a day in a given temperature bin, relative to a day in the omitted bin (12°C), on per capita GDP. Data include more than 34 000 county-year observations spanning 2003–2014. In order to plot air and surface temperature on the same axis, surface temperature units were transformed into air temperature units using the quantile mapping procedure described in Methods.

Economic Analysis to estimate the regression:

$$Y_{it} = \rho Y_{i,t-1} + \sum_m (\beta^m T_{it}^m + \alpha^m T_{it-1}^m) + Pr_{it} + Pr_{it}^2 + \gamma_i + \delta_t + \varepsilon_{it} \quad (3)$$

where  $Y_{it}$  is county per-capita GDP in county  $i$  and year  $t$ ,  $Y_{i,t-1}$  is lagged per-capita GDP,  $\beta^m T_{it}^m$  is the number of days in the  $m$ th 3-degree bin<sup>4</sup> in county  $i$  and year  $t$ ,  $Pr$  is total precipitation in the year,  $\gamma$  is a county fixed effect,  $\delta$  is a year fixed effect, and  $\varepsilon$  is an error term. Population weights were used to spatially aggregate MODIS and PRISM grid cells to the county level and PRISM precipitation estimates were used in both regressions.


Estimates of equations (1) through (3) were then used to plot the climate response functions shown in figures 3 and 4. In order to plot the relationships shown in figure 3, splines were fit to mean impacts at each

temperature level estimated by equations (1) and (2). Bootstrapped standard errors were then calculated and used to estimate 95% confidence intervals. Figure 4 shows the coefficients for each temperature bin estimated by equation (3).  $T_s$  has a different support from  $T_a$ . Therefore, in order to facilitate a straightforward comparison, we mapped  $T_s$  to  $T_a$  by matching distribution quantiles and plotting the two response functions with  $T_a$  on the x-axis. The mapping was done by calculating 1 000 equally spaced quantiles separately for  $T_s$  and  $T_a$  then defining a function that matched  $T_s$  quantiles to  $T_a$  quantiles. This procedure transformed the  $T_s$  distribution into the  $T_a$  distribution and allows for a simple comparison in familiar units.

### 3. Results

For maize yields, we find downward sloping responses to temperature for both temperature measures

<sup>4</sup> Since we used 8-day MODIS composites,  $T_s$  values falling into a given bin were counted as 8 days in that bin.



**Figure 5.** Comparing Climate Impacts Estimated with Ground Stations and Satellite Data. Results show the average impact from 1°C warming across the sample populations for different applications. Impacts were estimated with models using air temperatures from ground stations ( $T_a$ ) or surface temperatures from satellite data ( $T_s$ ). Error bars show the bootstrapped 95% confidence intervals.

(figure 3). For the Africa analysis,  $T_s$  had slightly higher explanatory power than  $T_a$ , with  $R^2$  values for  $T_s$  2.7% and 4.2% higher, respectively, for the optimal and drought management trials examined in the original study. For the United States, the  $R^2$  values are virtually the same across models, consistent with a dense and high-quality ground-station network. In order to further assess model performance we also calculated out-of-sample prediction error by repeatedly estimating the models on randomly selected 75% subsets of locations, predicting values for the 25% of locations that had been excluded from estimation, and calculating the RMSE of out-of-sample predicted values relative to actual values. For both optimal and drought management systems in Africa we find that the model with  $T_s$  has lower out-of-sample prediction errors. However, for the United States, the prediction RMSE values are nearly identical across temperature measures. This finding is consistent with our assertion that  $T_s$  is most useful in regions with poor station coverage where  $T_a$  is measured with significant levels of error.

While  $T_s$  predicts crop yields well, it is less clear whether it could be used to estimate response functions for non-agricultural applications. Recent research suggests that a variety of economic activities respond negatively to higher temperatures [13, 16] and our findings suggest that  $T_s$  is, in fact, suitable for estimating economic responses to temperature changes. For our GDP analysis, we find similar non-linear response functions using  $T_a$  and  $T_s$  over most of the temperature support, particularly at the upper end of the temperature distribution where income appears to be most sensitive to temperature (figure 4). The  $R^2$  values for the two models are similar

(0.541 for  $T_a$  model, 0.539 for  $T_s$  model) and the out-of-sample prediction RMSE values are indistinguishable. One apparent difference across temperature measures is that the model with  $T_a$  finds a positive effect of extreme low temperatures on income while the model with  $T_s$  finds no effect over the same range of the temperature distribution. However, the confidence intervals for the two estimates are overlapping at low temperatures.

#### 4. Discussion

Overall, we find that  $T_s$  is a suitable replacement for  $T_a$  in all three applications considered, with  $T_s$  even outperforming  $T_a$  with respect to prediction error in the Africa study, a region of low station density. Another approach to evaluating  $T_s$  performance is to compare the aggregated impacts from 1°C warming estimated with models using  $T_s$  and  $T_a$  (figure 5). In doing so we again find similar estimates for all applications. This overall consistency is perhaps somewhat surprising, given the often low correlations between anomalies in  $T_s$  and  $T_a$  at the daily or 8-day time scale. We view four factors as important in explaining the relative success of  $T_s$ . First, some of the ‘noise’ in  $T_s$  vs.  $T_a$  relationships stems from errors in the  $T_a$  measures, particularly in regions such as Africa where  $T_a$  is often interpolated from anomalies at stations tens of kilometers away. Second, much of the noise likely cancels out when aggregating temperatures to the monthly or seasonal time scales that are used in regressions that relate outcomes to temperature. For applications that require finer temporal resolution of temperature measures, the noise in  $T_s$  may become

more important—although again, whether it is larger than noise in high-temporal-resolution  $T_a$  remains an empirical question. Third, unlike ground measurements, satellite data come from a consistent sensor. Relative spatial variations could therefore be captured more precisely with satellites than with ground measurements from different instruments. Fourth, in vegetated areas much of the noise in the daytime  $T_s$  vs.  $T_a$  relationship arises from anomalous canopy transpiration rates, with stressed canopies often several degrees warmer than  $T_a$  whereas healthy canopies are typically several degrees below  $T_a$  [8, 10]. Thus,  $T_s$  provides a more direct measure of crop condition than  $T_a$ , and this represents an advantage of  $T_s$  for agricultural applications that may compensate for some of its deficiencies.

The substitutability of  $T_s$  for  $T_a$  suggests the potential usefulness of  $T_s$  for future study in areas with limited availability of reliable temperature data. For example, widespread surveys of health and economic activity such as the Demographic and Health Survey (DHS) and Living Standards Measurement Study (LSMS) are available in areas throughout the world with extremely poor weather station availability. Linking these measured outcomes to the MODIS  $T_s$  record, which now spans over 13+ years, will enable improved understanding of how climate trends and extremes affect human livelihoods around the world.

## References

- [1] Harris I, Jones P D, Osborn T J and Lister D H 2014 Updated high-resolution grids of monthly climatic observations—the cru ts3. 10 dataset *Int. J. Clim.* **34** 623–42
- [2] Willmott C J and Kenji M 2015 *Terrestrial Air Temperature: 1900–2014 Gridded Monthly Time Series Version 4.0.1* ([http://climate.geog.udel.edu/~climate/html\\_pages/Global2014/](http://climate.geog.udel.edu/~climate/html_pages/Global2014/))
- [3] Auffhammer M, Hsiang S M, Schlenker W and Sobel A 2013 Using weather data and climate model output in economic analyses of climate change *Rev. Environ. Econ. Policy* **7** 181–98
- [4] Vancutsem C, Ceccato P, Dinku T and Connor S J 2010 Evaluation of modis land surface temperature data to estimate air temperature in different ecosystems over africa *Remote Sens. Environ.* **114** 449–65
- [5] Lakshmi V and Susskind J 2000 Comparison of tovs-derived land surface variables with ground observations *J. Geophys. Res.* **105** 2179–90
- [6] Pinheiro A, Privette J and Bates J 2008 Satellite retrieval of land surface temperature: Challenges and opportunities *20th Conf. on Climate Variability and Change, American Meteorological society* (<https://ams.confex.com/ams/pdfpapers/131227.pdf>)
- [7] Mendelsohn R, Kurukulasuriya P, Basist A, Kogan F and Williams C 2007 Climate analysis with satellite versus weather station data *Clim. Change* **81** 71–83
- [8] Moran M S, Clarke T R, Inoue Y and Vidal A 1994 Estimating crop water deficit using the relation between surface-air temperature and spectral vegetation index *Remote Sens. Environ.* **49** 246–63
- [9] Anderson M C, Hain C, Wardlow B, Pimstein A, Mecikalski J R and Kustas W P 2011 Evaluation of drought indices based on thermal remote sensing of evapotranspiration over the continental united states *J. Clim.* **24** 2025–44
- [10] Siebert S, Ewert F, Rezaei E E, Kage H and Graß R 2014 Impact of heat stress on crop yield on the importance of considering canopy temperature *Environ. Res. Lett.* **9** 044012
- [11] Lobell D, Bänziger M, Magorokosho C and Vivek B 2011 Nonlinear heat effects on african maize as evidenced by historical yield trials *Nat. Clim. Change* **1** 42–5
- [12] Burke M and Emerick K 2016 Adaptation to climate change: Evidence from us agriculture *Am. Econ. J. Econ. Policy* **8** 106–40
- [13] Deryugina T and Hsiang S 2014 Does the environment still matter? daily temperature and income in the united states. *NBER Working Paper 20750* ([www.nber.org/papers/w20750](http://www.nber.org/papers/w20750))
- [14] PRISM Climate Group, Oregon State University. (<http://prism.oregonstate.edu>)
- [15] Schlenker W and Roberts M J 2009 Nonlinear temperature effects indicate severe damages to us crop yields under climate change *Proc. Natl Acad. Sci. USA* **106** 15594–8
- [16] Burke M, Hsiang S and Miguel E 2015 Global non-linear effect of temperature on economic production *Nature* **527** 235–9
- [17] Wan Z 2007 Collection-5 modis land surface temperature products users' guide. ICES, University of California, Santa Barbara ([https://lpdaac.usgs.gov/sites/default/files/public/product\\_documentation/mod11\\_user\\_guide.pdf](https://lpdaac.usgs.gov/sites/default/files/public/product_documentation/mod11_user_guide.pdf))
- [18] Bureau of Economic Analysis, U.S. Department of Commerce ([www.bea.gov/regional/downloadzip.cfm](http://www.bea.gov/regional/downloadzip.cfm))