

Mapping cropland-use intensity across Europe using MODIS NDVI time series

This content has been downloaded from IOPscience. Please scroll down to see the full text.

2016 Environ. Res. Lett. 11 024015

(<http://iopscience.iop.org/1748-9326/11/2/024015>)

View [the table of contents for this issue](#), or go to the [journal homepage](#) for more

Download details:

IP Address: 210.77.64.109

This content was downloaded on 07/04/2017 at 04:35

Please note that [terms and conditions apply](#).

You may also be interested in:

[Hotspots of land use change in Europe](#)

Tobias Kuemmerle, Christian Levers, Karlheinz Erb et al.

[Quantifying yield gaps in wheat production in Russia](#)

Florian Schierhorn, Monireh Faramarzi, Alexander V Prishchepov et al.

[Influence of management and environment on Australian wheat: information for sustainable intensification and closing yield gaps](#)

B A Bryan, D King and G Zhao

[Mapping and analysing cropland use intensity from a NPP perspective](#)

Maria Niedertscheider, Thomas Kastner, Tamara Fetzl et al.

[Mapping the extent of abandoned farmland in Central and Eastern Europe using MODIS time series satellite data](#)

Camilo Alcantara, Tobias Kuemmerle, Matthias Baumann et al.

[Long-term agricultural land-cover change and potential for cropland expansion in the former Virgin Lands area of Kazakhstan](#)

Roland Kraemer, Alexander V Prishchepov, Daniel Müller et al.

[Integrated crop water management might sustainably halve the global food gap](#)

J Jägermeyr, D Gerten, S Schaphoff et al.

[Investigating potential transferability of place-based research in land system science](#)

Tomáš Václavík, Fanny Langerwisch, Marc Cotter et al.

Environmental Research Letters



LETTER

Mapping cropland-use intensity across Europe using MODIS NDVI time series

OPEN ACCESS

RECEIVED

23 April 2015

REVISED

13 October 2015

ACCEPTED FOR PUBLICATION

18 January 2016

PUBLISHED

19 February 2016

Original content from this work may be used under the terms of the [Creative Commons Attribution 3.0 licence](#).

Any further distribution of this work must maintain attribution to the author(s) and the title of the work, journal citation and DOI.

Stephan Estel¹, Tobias Kuemmerle^{1,2}, Christian Levers¹, Matthias Baumann¹ and Patrick Hostert^{1,2}¹ Geography Department, Humboldt-University zu Berlin, Unter den Linden 6, 10099 Berlin, Germany² Integrative Research Institute on Transformations of Human-Environment Systems (IRI THESys), Humboldt-University zu Berlin, Unter den Linden 6, 10099 Berlin, GermanyE-mail: stephan.estel@geo.hu-berlin.de**Keywords:** cropping frequency, crop duration ratio, fallow cycles, Europe, multi-cropping, self-organizing maps, land-use intensitySupplementary material for this article is available [online](#)**Abstract**

Global agricultural production will likely need to increase in the future due to population growth, changing diets, and the rising importance of bioenergy. Intensifying already existing cropland is often considered more sustainable than converting more natural areas. Unfortunately, our understanding of cropping patterns and intensity is weak, especially at broad geographic scales. We characterized and mapped cropping systems in Europe, a region containing diverse cropping systems, using four indicators: (a) cropping frequency (number of cropped years), (b) multi-cropping (number of harvests per year), (c) fallow cycles, and (d) crop duration ratio (actual time under crops) based on the MODIS Normalized Difference Vegetation Index (NDVI) time series from 2000 to 2012. Second, we used these cropping indicators and self-organizing maps to identify typical cropping systems. The resulting six clusters correspond well with other indicators of agricultural intensity (e.g., nitrogen input, yields) and reveal substantial differences in cropping intensity across Europe. Cropping intensity was highest in Germany, Poland, and the eastern European Black Earth regions, characterized by high cropping frequency, multi-cropping and a high crop duration ratio. Contrarily, we found lowest cropping intensity in eastern Europe outside the Black Earth region, characterized by longer fallow cycles. Our approach highlights how satellite image time series can help to characterize spatial patterns in cropping intensity—information that is rarely surveyed on the ground and commonly not included in agricultural statistics: our clustering approach also shows a way forward to reduce complexity when measuring multiple indicators. The four cropping indicators we used could become part of continental-scale agricultural monitoring in order to identify target regions for sustainable intensification, where trade-offs between intensification and the environmental should be explored.

Introduction

Agricultural expansion and intensification have led to marked increases in agricultural production since World War II (Rudel *et al* 2009, Tilman *et al* 2002), albeit at substantial environmental costs (Tscharrntke *et al* 2012). Arguably, demand for agricultural products will have to increase in the future as the world's population grows, diets change, and bioenergy becomes more important (Beringer *et al* 2011, Erb *et al* 2013, Krausmann *et al* 2013). Demand-side strategies such as reducing meat consumption and

curbing food wastage (Bajzelj *et al* 2014, Erb *et al* 2009, Tilman *et al* 2011) are promising, but will likely not be enough. How production increases could be achieved while curbing the environmental impacts of agriculture remains unclear (Butchart *et al* 2010, Foley *et al* 2011, West *et al* 2014).

Expanding agriculture further into the last remaining undeveloped fertile lands in South America and Africa would entail drastic environmental costs (e.g., substantial carbon emissions and biodiversity loss) (Laurance *et al* 2014, Licker *et al* 2010, Ramanakutty *et al* 2002). Crop production can also be

increased by intensifying agriculture on existing cropland, yet underperforming cropland (Godfray *et al* 2010, Johnston *et al* 2011). Such intensification could, for example, entail an optimization of crop rotations (e.g., less fallow) and higher resource efficiency (e.g., nutrient, water use) (Mueller *et al* 2012, Ray and Foley 2013, Siebert *et al* 2010b). Since the environmental impacts of intensification can be substantial (Licker *et al* 2010, Matson *et al* 1997), careful, context-specific assessments of the risks and opportunities of intensification are required (Garnett *et al* 2013).

Information on spatial and temporal patterns of cropland use at multiple geographic scales is required to better understand the potential for intensification. Unfortunately, existing data on cropland-use intensity are mostly coarse in scale, heavily rely on uncertain cropland maps (Fritz *et al* 2011, Fritz *et al* 2013), or are based on national statistics, which themselves may contain uncertainties (Verburg *et al* 2011, Zaks and Kucharik 2011). Many existing datasets represent snapshots in time and cannot reflect the often highly dynamic management intensity of agricultural land (Kuemmerle *et al* 2013, Siebert *et al* 2010b). Substantial progress in mapping indicators of cropland-use intensity has been made recently, including yield gaps (Johnston *et al* 2011, Monfreda *et al* 2008, West *et al* 2014), fertilizer use (Potter *et al* 2010), human appropriation of net primary production (HANPP) (Haberl *et al* 2012), field size (Fritz *et al* 2015) or the extent of irrigated agriculture (Siebert *et al* 2010a, Thenkabail *et al* 2009) or tillage (Johnson 2013). Yet, our understanding of cropping patterns (i.e., temporal dynamics of cropland use), such as crop rotations, multi-cropping (i.e., number of harvests per year), crop duration (i.e., fraction of the year in which the cropland is covered with crops), cropping frequency (i.e., the number of cropped years), or the fallow land extent (Kuemmerle *et al* 2013, Li *et al* 2014, Portmann *et al* 2010, Siebert *et al* 2010b) is limited.

Dense time series of medium-resolution satellite images, such as from the Moderate Resolution Imaging Spectroradiometer (MODIS), can help identifying cropland dynamics across broad geographic extents (Friedl *et al* 2010, Ganguly *et al* 2010, Rogan and Chen 2004, Siebert *et al* 2010b). For example, in the Russian grain belt, cropping frequency was mapped between 2002 and 2009 using phenological metrics (de Beurs and Ioffe 2013). MODIS vegetation indices allow the differentiation of single, double, or triple cropping, for example in Brazil (Galford *et al* 2008, Spera *et al* 2014), India (Biradar and Xiao 2011), the Mekong Delta (Sakamoto *et al* 2009), China (Li *et al* 2014), or the mid-western United States (Wardlow and Egbert 2008). These studies highlight the potential of medium-resolution sensors to map cropping indicators, but used only single indicators over relatively short time periods to describe often

highly dynamic and heterogeneous agricultural systems.

Our main goal here was to characterize European cropping systems by mapping four MODIS-based cropping indicators and by identifying typical cropping clusters. Europe is interesting for assessing patterns of cropland-use intensity for at least three main reasons. First, Europe is characterized by a wide range of agricultural systems, spanning strong environmental and socio-economic gradients, resulting in systems with different degrees of cropland-use intensity (Herzog *et al* 2006, Rounsevell *et al* 2012, Rudel *et al* 2009). Second, Europe has a long land-use history, with most land-use change nowadays happening along gradients of intensity change. Third, Eastern Europe experienced a dramatic decline in cropland-use intensity after the breakdown of socialism (Kuemmerle *et al* 2013, Prishchepov *et al* 2012) and is consistently highlighted as a candidate region for sustainable intensification (Foley *et al* 2011, Müller *et al* 2013). Therefore Europe is a prime example to develop methods that allow capturing and mapping cropland-use intensity and changes therein.

Specifically, we assessed the following research questions:

1. What were the spatial patterns of cropping intensity in Europe from 2001 to 2012, as measured by cropping frequency, multi-cropping, fallow cycles, and crop duration ratio?
2. What are regions of similar cropping systems across Europe?

Data and methods

Our study area included the entire European continent and Turkey to define the cropland extent within this region, we used the GlobCorine land-cover map from 2005 (Defourny *et al* 2010). To overcome scale differences between GlobCorine (300 m) and the MODIS time series (231.6 m), we downscaled the prior to the latter using a nearest neighbour algorithm. Based on this map we focussed on all classes containing cropland, specifically rainfed and irrigated cropland, complex cropland, and mosaic cropland/natural vegetation, and we masked all other classes (figure S1 in the supplementary material).

To map cropping indicators, we used a NDVI time series pre-processed in a previous study following three steps (Estel *et al* 2015). First, using satellite images from both Terra and Aqua satellites from 2000 to 2012 we reduced effects from clouds, water, snow, and ice by excluding poor-quality observations based on the MODIS quality information, land surface temperature, the land-water mask and interpolating missing values. Second, due to the strong climate gradient across Europe and the resulting varying phenology (e.g., earlier green-up and shifted vegetation peak in

Table 1. Intensity classes for each indicator (low, medium, and high, based on terciles) and indicators' class share from the total cropland.

	Intensity class			
	Low	Medium	High	
Cropping frequency	1–4	5–8	9–12	
Multi-cropping	1–4	5–8	9–12	
Fallow cycles	0.33–0.91	0.92–1.49	1.50–2.08	
Crop duration ratio	0.16–0.43	0.44–0.70	0.71–0.98	
Area share [Mha]				
	Low	Medium	High	Total
Cropping frequency	20.7	72.5	306.9	400.1
Multi-cropping	180	37.8	6.4	224.2
Fallow cycles	77.9	26.5	4.6	108.9
Crop duration ratio	47.0	274.3	78.8	400.1
Area share [%]				
	Low	Medium	High	Total
Cropping frequency	5.2	18.1	76.7	100
Multi-cropping	45.0	9.4	1.6	56.0
Fallow cycles	19.5	6.6	1.1	27.2
Crop duration ratio	11.7	68.5	19.7	100.0

the Mediterranean, higher seasonality in the north), we normalized the NDVI time series to make them more comparable. The normalization was twofold and included an accounting for the shifted vegetation peak in Mediterranean environments in Europe, and harmonization the NDVI time series in regards to vegetation maxima and amplitudes across Europe. The normalization procedure is described in detail in Estel *et al* (2015). Third, we gathered an extensive training dataset on active and fallow cropland by interpreting the NDVI time series and high-resolution images from GoogleEarth, and classified each cropland pixel into active (i.e., managed) and fallow farmland (i.e., unmanaged) for each year between 2001 and 2012. The annual active/fallow maps had an average overall accuracy of >90% based on independent validation data (Estel *et al* 2015).


Mapping cropping indicators

We used the pre-processed NDVI time series and the annual fallow/active maps from Estel *et al* (2015) to map four cropland-use intensity indicators: (1) cropping frequency, (2) multi-cropping, (3) fallow cycles, (4) crop duration ratio at a spatial resolution of 231.6 m. To provide an intensity measurement and to better describe and compare the mapped patterns of intensity we divided each indicator into a high, medium and low intensity class (table 1). We

furthermore excluded all pixels labelled as abandoned or permanently fallow based on Estel *et al* (2015). Our final cropland mask included an area of 400 Mha. For the indicators crop duration ratio and multi-cropping, we considered only non-fallow years. We calculated the cropping frequency as the number of years a cropland pixel was cropped over the observation period (see supplementary material). Higher cropping frequencies thus signifies higher cropland-use intensity (table 1). Multi-cropping refers to the number of harvests within a single year (i.e., growing season, Spera *et al* 2014). In Europe, either single or double-cropping occurs. To identify double-cropping, we counted the number vegetation peaks per growing season using TIMESAT (Jönsson and Eklundh 2004, Li *et al* 2014), which detects double peaks based on the amplitude ratio between the primary and the secondary peak (Jönsson and Eklundh 2004). We derived annual single vs. double-cropping maps and summarized the number of double-cropped years (table 1).

Fallow cycles refer to recurring periods of fallow cropland. Longer and frequent fallow periods thus signify less intense land management. We defined fallow as cropland without management (i.e., not sown, cropped, or ploughed) (Estel *et al* 2015). ‘Active fallow’ (e.g., cultivation of legumes for nitrogen-fixation) was not considered. To identify fallow cycles, we screened the active/fallow time series for ‘chain segments’, i.e., a certain number (1, 2 or 3) of consecutive fallow years. We identified chain segments consisting of one (FC1), two (FC2), and three (FC3) fallow years between active years. We counted the occurrence of these chains across the entire time-series per pixel (see figure S3), and summarized all chain segments using a weighting scheme. Weights were calculated as the ratio of the total number of years in the time series (12) and the number of maximally possible chain segments of a particular cycle type (see supplementary material). The resulting index provides information about the level of cyclicity, and thus management intensity. We considered only time series with at least two chain segments (figure 3).

The crop duration ratio is the time a field is cropped in relation to the total length of the growing season (Siebert *et al* 2010b). We derived the total length of the growing season as the number of days with a land surface temperature above 5 °C, i.e., the time between the earliest and the latest MODIS acquisition date when plants are assumed to actively grow (Hickler *et al* 2012, Zhang *et al* 2004). The time period a pixel was cropped was defined as a vegetation signal of at least half the peak of the phenological curve (see supplementary material). The half-maximum is frequently used as a phenological marker for leaf unfolding and the loss of canopy structure of natural vegetation (Bradley *et al* 2007, Fisher *et al* 2006) and we used it as a proxy for crop green-up and harvesting. We then computed the crop duration ratio for each year and calculated the average crop duration ratio from 2001 to 2012. Since



cropping cycles are more dynamic than natural vegetation (e.g., varying timing of ploughing, sowing and harvesting), we carried out a sensitivity analyses to test the robustness of our results in relation to the choice of threshold, and derived crop duration ratios for thresholds of 40%, 45%, 50% (=half-maximum), 55%, and 60%. We then calculated the standard deviation crop duration (see supplementary material).

Mapping typical clusters of cropland-use intensity

To identify similar cropping systems, we used self-organizing maps (SOMs). SOMs are an unsupervised clustering technique based on competitive learning that reduce a high-dimensional dataset to a two dimensional map by grouping observations according to their similarity (Skupin and Agarwal 2008). To identify the optimal number of clusters, we applied a sensitivity analyses with SOM clusters varying from 2×2 to 4×4 clusters (Maulik and Bandyopadhyay 2002) and used the Davies–Bouldin index that compares intra- and inter-cluster variability (Davies and Bouldin 1979) to pick the optimal cluster number. We z-transformed our indicators prior to the clustering, and calculated average values across a 1×1 km² grid (see supplementary material).

Results




In terms of cropping frequency, we found around 166.8 Mha (i.e., 41.7%) of all European croplands cultivated every year during 2001–2012. These areas were mainly located in western and central Europe (northern France, most of Germany, parts of England), northern Italy, eastern and northern Spain, Turkey and the Black Earth regions (i.e. Chernozem, FAO/EC/ISRIC 2003) of southern Russia and south-eastern Ukraine (figure 1). Around 18% of the croplands had medium cropping frequencies (i.e., 5 to 8


cropped years) and occurred mainly in the Mediterranean (e.g., Extremadura, southern Portugal) and north-western Germany (table 1). Around 5% of the croplands had lower cropping frequencies (i.e., 1–4 cropped years) mainly in mountain regions (e.g., Alps, Pyrenees, and Caucasus) and eastern Europe (e.g., Russia, northern Ukraine, Belarus, and the Baltics).

Multi-cropping was widespread in the study area. About 56% (table 1) of all croplands in Europe were double-cropped at least once during 2001–2012. Areas where multi-cropping was high (i.e., 9–12 double-cropped years) accounted for only 2% of the total croplands, and were most widespread in central Europe (i.e., north-eastern Germany, central Poland) and Russia (i.e., Black Earth regions). Medium multi-cropping (i.e., 5–8 double-cropped years) accounted for about 9% of all croplands, mainly in central Europe, central Spain, southern Ukraine, Russia (figure 2).

Our fallow cycle mapping showed that about 27% of all European croplands had one of the three fallow cycles (figure S3). Out of all croplands about 1% had a high fallow cyclicity (upper tercile of index values) and these areas occurred predominately in the southern Iberian Peninsula, north-eastern Turkey, and eastern Europe (table 1). About 26% had a medium (i.e., mid tercile) or low fallow cyclicity (i.e., lower tercile) and occurred all over Europe with concentrations on the Iberian Peninsula, eastern Europe (i.e., northern Ukraine, Russia) and some Mediterranean areas.

Mapping crop duration ratio (figure 4) revealed that about 20% of all European cropland was characterized by high crop duration ratios, mainly in central Europe (e.g., Germany, Poland, eastern France, and southern Hungary). About 69% of all croplands showed medium crop duration ratios (i.e., 0.44–0.70), occurring mainly in European Russia, Ukraine, UK and eastern France (table 1). A few regions (~12%) showed lower crop duration ratios (i.e., <0.43), mainly in Spain, Turkey, Italy, Greece and southern






European Russia. Our sensitivity analyses showed the robustness towards alternative definitions, with standard deviations in crop duration ratio $<20\%$. For some Mediterranean areas, where crop duration ratio is lower than elsewhere in Europe, we found a higher, but still moderate sensitivity (see supplementary material).

We identified six clusters of similar cropping systems (figure 5) using SOMs and a sensitivity analysis across varying cluster numbers (see supplementary material). To describe the magnitude and direction of the different cropping indicators in each cluster (C1–C6), we provided here the deviation (\pm) from mean z-score ($=0$, table S1). Positive and negative numbers thus signify above and below average values respectively, whereas values close to zero mean that a specific indicator is close to the overall mean of the study area (figure 6). Cluster 1 was determined by high cropping frequencies (+0.82%), crop duration ratios slightly above average (+0.22%), and a very low fallow cyclicity (−1.25%). This cluster occurred in northern France, England, Italy, and around the Black Sea in Romania, Ukraine, Russia, and Turkey. Cluster 2 was mainly determined by very high multi cropping (+2.76%), high crop duration ratios (+0.80%) and cropping frequencies (+0.72%), but a low fallow cyclicity (−0.67%). This cluster 2 was mainly located in Germany, central Poland, southern Russia, and western France. Cluster 3 was characterized by very low cropping frequencies (−1.78%), infrequent multi-cropping (−0.80%), a marked fallow cyclicity (+0.73) and occurred mainly in eastern Europe (e.g., European Russia, Baltics, Belarus, and northern Ukraine), the

Mediterranean, and in mountain areas (e.g., Alps, Pyrenees, Caucasus). Cluster 4 had very low crop duration ratios (−1.62%) and a very low fallow cyclicity (−1.23%), and the highest cropping frequencies (+0.85%) of all cluster. This cluster occurred mainly in the Mediterranean, southern Ukraine and southern Russia. Cluster 5 was characterized by a high fallow cyclicity (+0.73%), low multi-cropping (−0.53%) and low crop duration ratios (−0.35%). This cluster occurred mainly in Ukraine and Russia, on the Iberian Peninsula, western France, and Turkey. Cluster 6 had the highest fallow cyclicity (+0.74%) and the highest crop duration ratios (+1.00%) of all cluster. This cluster occurred mainly in central Europe and southern Ukraine.

Discussion

Understanding spatial patterns in cropland-use intensity is important for identifying target regions for intensification, and for assessing its potential environmental trade-offs. Better information on cropping systems, including cropping frequency, fallow cycles, multi-cropping, and crop duration, are important indicators in this context; yet agricultural census do not cover them at fine scale. Using a 12-year MODIS NDVI time series, we mapped these cropping indicators at the continental scale for Europe. The cropping patterns we find correspond well with other indicators of agricultural land management in Europe, highlighting the potential for satellite-based cropping measures to support agricultural monitoring.



Moreover, our indicators capture important aspects of agricultural intensity, highlighting how management intensity varies in space and time. The identified cropping systems can be explained by agro-environmental conditions (e.g., soil quality, water availability), socio-economic conditions (e.g., rural depopulation), management practices (e.g., irrigation), and crop-specific management (e.g., rice growing in northern Italy). Our satellite-based indicators of cropland-use intensity and the mapping of cropping systems may help to identify baselines and candidate region for intensifying croplands sustainably.

The cropping patterns captured by our indicators correspond well with those from other indicators of agricultural management in Europe. For example, existing maps of cropland net primary production, yields, and yield gaps consistently show the highest output intensity where we identified intensive used multi-cropping systems and a long crop duration (Monfreda *et al* 2008, Mueller *et al* 2012, Neumann *et al* 2010). Maps of fertilizer usage also show highest fertilizer use in these regions (Potter *et al* 2010, Temme and Verburg 2011). In contrast, these maps suggest low input and output cropland intensity in eastern Europe, especially in European Russia, Ukraine, and

the Baltic States, congruent with our cropping indicators. A more quantitative comparison among our maps and other land-use intensity indicators is not feasible, given different resolutions and time periods covered, and considering that cropland-use indicators related to inputs and outputs usually represent down-scaling agricultural statistics (usually national scale). Although multi-cropping has increased due the bio-fuel expansion (in Europe mostly based on wheat) in the last decades it is not considered in agricultural statistics (Langeveld *et al* 2014).

That our satellite-based indicators identify, on a general level, the similar spatial patterns of high and low intensity is encouraging, and highlights the potential of satellite-based to observe and monitor agricultural systems more directly and with finer spatial detail (Kuemmerle *et al* 2013, Zaks and Kucharik 2011).

The spatial congruence between our cropping indices and alternative measures of agricultural management, such as fertilizer and yields, also underlines the value of satellite-based cropping indices to more directly measure cropland-use intensity. Yet some of our indicators, for example cropping frequency, (i.e., more cropped years indicates higher intensity) are

more closely related to intensity than others (e.g., crop duration ratio). A few points of caution need to be mentioned when interpreting our individual indicators. First, our multi-cropping indicator captured whether there are one or two vegetation peaks in a given year, but not in all cases a second peak refers to a second crop, nor does it attest to whether the second crop was actually harvested or left in the field as green manure. Second, our fallow cycle indicator captured cropping cycles over multiple years, but rests on a reliable identification of active vs. fallow cropland. Although our accuracy assessment of the active/fallow maps suggests these maps are reliable (>90% overall accuracy, Estel *et al* 2015), uncertainty for some regions (e.g., with small fields) may be higher than for others. Third, we measured crop duration ratio as the length of the vegetation signal on cropland, but lower crop duration ratios may be due to management or climate. To interpret crop duration ratio as a measure of cropland intensity, agro-environmental conditions and irrigation structure should be considered in future work. Finally, validation of our cropping indicators is challenging because retrospective field-level data on cropping do not exist for larger areas, and *in situ* data are currently not feasible to gather at the continental scale. However, we note that the annual fallow/active maps derived in Estel *et al* (2015) and used as input data for the cropping intensity indicators cropping frequency and fallow cycle has been validated extensively (Estel *et al* 2015), and sensitivity analyses for the crop duration indicator attest to the robustness of these indicator.

SOMs were a useful tool to identify regions with similar cropping and thus to help interpret and reduce complexity in our multi-dimension indicator dataset. The six clusters of cropping systems appear to be related to distinctly different agro-environmental and socio-economic conditions across Europe. Cluster 1 characterized moderately intensive rain-fed cropping and included some irrigated areas in southern Europe along rivers and reservoirs (e.g., Ebro-basin in northern Spain, Po Valley in northern Italy, Black Sea area) (Siebert *et al* 2006, Salmon *et al* 2015). Irrigation here led to an uncoupling from climate constraints, allowing for central/western European cropping systems in these areas.

Cluster 2 was clearly linked to the most intensified rain-fed cropping in highly favorable agro-environmental conditions (e.g. in Germany and Denmark, Neumann *et al* 2010). Interestingly, this system was also found in Europe's east, particularly, in southern Ukraine, Romania and southern Russia, where some of the world's most fertile soils are found (Fischer *et al* 2000). High double cropping rates that characterized this cluster (e.g., southern Russia, northern Germany) are linked to the cultivation of winter wheat as main crop. The specific cropping time of winter wheat allows preceding or subsequent crops (e.g., rape,

summer wheat) (Gienapp *et al* 2012, Schierhorn *et al* 2014).

Cluster 3, characterized by frequent fallow years, occurred mainly in water-limited regions (e.g., Extremadura and northern Andalusia in Spain, southern Portugal). Fallowing is characteristic for semi-arid regions to maintain soil moisture and fertility (Boellstorff and Benito 2005). Interestingly, this cropping system also occurred in eastern Europe, although agro-environmental conditions are more favorable there. The higher fallow frequencies there are possibly a legacy of the breakdown of socialism and the subsequent restructuring of agricultural sectors (Fischer *et al* 2000, EU 2005), leading to widespread farmland dis-intensification and abandonment (Rey Benayas 2007, Kuemmerle *et al* 2011, Prishchepov *et al* 2012). Similar trends are ongoing in western Europe's marginal regions (e.g., mountain regions) (Gellrich *et al* 2007, MacDonald *et al* 2000), many of which fell in the same cluster in our analyses.

Cluster 4 was clearly related to rain-fed cropping under water limitation (e.g., the Mediterranean), allowing only for shorter growing seasons (Fischer *et al* 2000). This does not necessarily indicate lower cropland-use intensity since yields in this regions can be high (Mueller *et al* 2012). For example, rice growing areas in the Po Valley (northern Italy) had very short crop durations but are considered among the most intensively used croplands in Europe (Blengini and Busto 2009, Erb *et al* 2013).

Finally, cluster 5 contained much cropland without major constraints (e.g., accessibility, soil fertility), yet that is currently not used to its full potential. Cluster 6, characterized by high fallow rates, occurred mainly in central and Eastern Europe in favorable agro-environmental conditions (e.g., southern Romania, southern Ukraine). Both cluster could entail particularly candidate regions for sustainable intensification.

In sum, our four satellite-based cropping indicators, as well as our cluster analyses to identify similar cropping systems across Europe, appear valuable for broad-scale agricultural monitoring. Satellite-based indicators could therefore complement ground-data that must always remain sample-based with wall-to-wall observations of agricultural management. Our study showed that time series of satellite images can also help to better characterize cropland-use intensity, and thus to assess baselines and potentials for intensifying croplands sustainably. Such information is critical to scrutinize the possible socio-economic and environmental trade-offs, as well as synergies, of intensifying croplands.

Acknowledgements

We gratefully acknowledge financial support by the Einstein Foundation Berlin (Germany), the European

Commission (VOLANTE, No. 265104 and HERCULES, No. 603447), and the DFG (LUCC-BIO, No. 3210 3109 99). We also would like to thank the ESA GlobCORINE Project for making the GlobCORINE map available.

References

- Bajzelj B, Richards K S, Allwood J M, Smith P, Dennis J S, Curmi E and Gilligan C A 2014 Importance of food-demand management for climate mitigation *Nature Clim. Change* **4** 924–9
- Beringer T, Lucht W and Schaphoff S 2011 Bioenergy production potential of global biomass plantations under environmental and agricultural constraints *GCB Bioenergy* **3** 299–312
- Biradar C M and Xiao X 2011 Quantifying the area and spatial distribution of double- and triple-cropping croplands in India with multi-temporal MODIS imagery in 2005 *Int. J. Remote Sens.* **32** 367–86
- Blengini G A and Busto M 2009 The life cycle of rice: LCA of alternative agri-food chain management systems in Vercelli (Italy) *J. Environ. Manag.* **90** 1512–22
- Boellstorff D and Benito G 2005 Impacts of set-aside policy on the risk of soil erosion in central Spain *Agriculture, Ecosystems & Environment* **107** 231–43
- Bradley B A, Jacob R W, Hermance J F and Mustard J F 2007 A curve fitting procedure to derive inter-annual phenologies from time series of noisy satellite NDVI data *Remote Sens. Environ.* **106** 137–45
- Butchart S H M *et al* 2010 Global biodiversity: indicators of recent declines *Science* **328** 1164–8
- Davies D L and Boulidin D W 1979 A cluster separation measure *IEEE Trans. on Pattern Analysis and Machine Intelligence PAMI-1* 224–7
- de Beurs K M and Ioffe G 2013 Use of landsat and MODIS data to remotely estimate Russia's sown area *J. Land Use Sci.* **9** 377–401
- Defourny P, Bontemps S, van Bogaert, E, Weber J L and Soukup T 2010 GlobCorine Validation Report (http://due.esrin.esa.int/files/p114/GLOBCORINE_VR_2.1_20100406.pdf) (accessed 24 March 2015)
- Erb K-H, Haberl H, Jepsen M R, Kuemmerle T, Lindner M, Müller D, Verburg P H and Reenberg A 2013 A conceptual framework for analysing and measuring land-use intensity *Current Opinion in Environmental Sustainability* **5** 464–70
- Erb K-H, Haberl H, Krausmann F, Lauk C, Plutzar C, Steinberger J K, Müller C, Bondeau A, Waha K and Pollack G 2009 *Eating the Planet: Feeding and Fuelling the World Sustainably, Fairly and Humanely—A Scoping Study Social Ecology Working* (Vienna: Institute of Social Ecology) p 116
- Estel S, Kuemmerle T, Alcántara C, Levers C, Prishchepov A and Hostert P 2015 Mapping farmland abandonment and recultivation across Europe using MODIS NDVI time series *Remote Sens. Environ.* **163** 312–25
- EU 2005 *Soil Atlas of Europe* (Luxembourg: European Communities)
- FAO/EC/ISRIC 2003 World Soil Resources (<http://fao.org/soils-portal/soil-survey/soil-maps-and-databases/en/>) (accessed 15 April 2015)
- Fischer G, Velthuis V H and Nachtergaele F O 2000 *Interim Report (IR-00-064): Global Agro-Ecological Zones Assessment: Methodology and Results* International Institute for Applied Systems Analysis (IIASA) (www.iiasa.ac.at/publication/more_IR-00-064.php)
- Fisher J, Mustard J and Vadeboncoeur M 2006 Green leaf phenology at landsat resolution: scaling from the field to the satellite *Remote Sens. Environ.* **100** 265–79
- Foley J A *et al* 2011 Solutions for a cultivated planet *Nature* **478** 337–42
- Friedl M A, Sulla-Menashe D, Tan B, Schneider A, Ramankutty N, Sibley A and Huang X 2010 MODIS collection 5 global land cover: algorithm refinements and characterization of new datasets *Remote Sens. Environ.* **114** 168–82
- Fritz S, See L, McCallum I, Schill C, Obersteiner M, van der Velde M, Boettcher H, Havlík P and Achard F 2011 Highlighting continued uncertainty in global land cover maps for the user community *Environ. Res. Lett.* **6** 044005
- Fritz S *et al* 2015 Mapping global cropland and field size *Glob. Change Biol.* **21** 1980–92
- Fritz S *et al* 2013 The need for improved maps of global cropland *Eos Trans. AGU* **94** 31–2
- Galford G L, Mustard J F, Melillo J, Gendrin A, Cerri C C and Cerri C E P 2008 Wavelet analysis of MODIS time series to detect expansion and intensification of row-crop agriculture in Brazil *Remote Sens. Environ.* **112** 576–87
- Ganguly S, Friedl M A, Tan B, Zhang X and Verma M 2010 Land surface phenology from MODIS: characterization of the collection 5 global land cover dynamics product *Remote Sens. Environ.* **114** 1805–16
- Garnett T *et al* 2013 Sustainable intensification in agriculture: premises and policies *Science* **341** 33–4
- Gellrich M, Baur P, Koch B and Zimmermann N E 2007 Agricultural land abandonment and natural forest re-growth in the Swiss mountains: a spatially explicit economic analysis *Agriculture, Ecosystems & Environment* **118** 93–108
- Gienapp C, Sanftleben P, Jennerich H J and Heilmann H 2012 Beiträge zur Weizenproduktion *Mitteilungen der Landesforschungsanstalt für Landwirtschaft und Fischerei, Landesforschungsanstalt für Landwirtschaft und Fischerei, Dorfplatz 1, 18276 Gülzow* (www.landwirtschaft-mv.de/cms2/LFA_prod/LFA/content/de/_Service/Mitteilungen_der_LFA/Heft_47/index.jsp)
- Godfray H C J, Beddington J R, Crute I R, Haddad L, Lawrence D, Muir J F, Pretty J, Robinson S, Thomas S M and Toulmin C 2010 Food security: the challenge of feeding 9 billion people *Science* **327** 812–8
- Haberl H, Steinberger J K, Plutzar C, Erb K-H, Gaube V, Gingrich S and Krausmann F 2012 Natural and socioeconomic determinants of the embodied human appropriation of net primary production and its relation to other resource use indicators *Ecological Indicators* **23** 222–31
- Herzog F *et al* 2006 Assessing the intensity of temperate European agriculture at the landscape scale *Eur. J. Agro.* **24** 165–81
- Hickler T *et al* 2012 Projecting the future distribution of European potential natural vegetation zones with a generalized, tree species-based dynamic vegetation model *Glob. Ecol. Biogeogr.* **21** 50–63
- Johnson D M 2013 A 2010 map estimate of annually tilled cropland within the conterminous United States *Agricultural Systems* **114** 95–105
- Johnston M, Licker R, Foley J, Holloway T, Mueller N D, Barford C and Kucharik C 2011 Closing the gap: global potential for increasing biofuel production through agricultural intensification *Environ. Res. Lett.* **6** 034028
- Jönsson P and Eklundh L 2004 TIMESAT—a program for analyzing time-series of satellite sensor data *Comput. Geosci.* **30** 833–45
- Krausmann F, Erb K-H, Gingrich S, Haberl H, Bondeau A, Gaube V, Lauk C, Plutzar C and Searchinger T D 2013 Global human appropriation of net primary production doubled in the 20th century *Proc. Natl Acad. Sci. USA* **110** 10324–9
- Kuemmerle T *et al* 2013 Challenges and opportunities in mapping land use intensity globally *Current Opinion in Environmental Sustainability* **5** 484–93
- Kuemmerle T, Olofsson P, Chaskovskyy O, Baumann M, Ostapowicz K, Woodcock C E, Houghton R A, Hostert P, Keeton W S and Radeloff V C 2011 Post-Soviet farmland abandonment, forest recovery, and carbon sequestration in western Ukraine *Glob. Change Biol.* **17** 1335–49
- Langeveld J W A, Dixon J, van Keulen H and Quist-Wessel P M F 2014 Analyzing the effect of biofuel expansion on land use in major producing countries: evidence of increased multiple cropping *Biofuels, Bioproducts and Biorefining* **8** 49–58

- Laurance W F, Sayer J and Cassman K G 2014 Agricultural expansion and its impacts on tropical nature *Trends in Ecology & Evolution* **29** 107–16
- Li L, Friedl M A, Xin Q, Gray J, Pan Y and Frolking S 2014 Mapping crop cycles in China using MODIS-EVI time series *Remote Sensing* **6** 2473–93
- Licker R, Johnston M, Foley J A, Barford C, Kucharik C J, Monfreda C and Ramankutty N 2010 Mind the gap: how do climate and agricultural management explain the ‘yield gap’ of croplands around the world? *Glob. Ecol. Biogeogr.* **19** 769–82
- MacDonald D, Crabtree J R, Wiesinger G, Dax T, Stamou N, Fleury P, Gutierrez Lazpita J and Gibon A 2000 Agricultural abandonment in mountain areas of Europe: environmental consequences and policy response *J. Environ. Manag.* **59** 47–69
- Matson P A, Parton W J, Power A G and Swift M J 1997 Agricultural intensification and ecosystem properties *Science* **277** 504–9
- Maulik U and Bandyopadhyay S 2002 Performance evaluation of some clustering algorithms and validity indices *IEEE Trans. Pattern Anal. Mach. Intell.* **24** 1650–4
- Monfreda C, Ramankutty N and Foley J A 2008 Farming the planet: 2. Geographic distribution of crop areas, yields, physiological types, and net primary production in the year 2000 *Global Biogeochemical Cycles* **22** GB1022
- Mueller N D, Gerber J S, Johnston M, Ray D K, Ramankutty N and Foley J A 2012 Closing yield gaps through nutrient and water management *Nature* **490** 254–7
- Müller D, Leitão P J and Sikor T 2013 Comparing the determinants of cropland abandonment in Albania and Romania using boosted regression trees *Agricultural Systems* **117** 66–77
- Neumann K, Verburg P H, Stehfest E and Müller C 2010 The yield gap of global grain production: a spatial analysis *Agricultural Systems* **103** 316–26
- Portmann F T, Siebert S and Döll P 2010 MIRCA2000—global monthly irrigated and rainfed crop areas around the year 2000: a new high-resolution data set for agricultural and hydrological modeling *Glob. Biogeochem. Cycles* **24** GB1011
- Potter P, Ramankutty N, Bennett E M and Donner S D 2010 Characterizing the spatial patterns of global fertilizer application and manure production *Earth Interactions* **14** 1–22
- Prishchepov A V, Radeloff V C, Baumann M, Kuemmerle T and Müller D 2012 Effects of institutional changes on land use: agricultural land abandonment during the transition from state-command to market-driven economies in post-Soviet Eastern Europe *Environ. Res. Lett.* **7** 024021
- Ramankutty N, Foley J A, Norman J and McSweeney K 2002 The global distribution of cultivable lands: current patterns and sensitivity to possible climate change *Glob. Ecol. Biogeogr.* **11** 377–92
- Ray D K and Foley J A 2013 Increasing global crop harvest frequency: recent trends and future directions *Environ. Res. Lett.* **8** 44041–50
- Rey Benayas J 2007 Abandonment of agricultural land: an overview of drivers and consequences *CAB Reviews: Perspectives in Agriculture, Veterinary Science, Nutrition and Natural Resources* **2** 14
- Rogan J and Chen D 2004 Remote sensing technology for mapping and monitoring land-cover and land-use change *Progress in Planning* **61** 301–25
- Rounsevell M D A *et al* 2012 Challenges for land system science *Land Use Policy* **29** 899–910
- Rudel T K *et al* 2009 Agricultural intensification and changes in cultivated areas, 1970–2005 *Proc. Natl Acad. Sci. USA* **106** 20675–80
- Sakamoto T, Van Phung C, Kotera A, Nguyen K D and Yokozawa M 2009 Analysis of rapid expansion of inland aquaculture and triple rice-cropping areas in a coastal area of the Vietnamese Mekong Delta using MODIS time-series imagery *Landscape and Urban Planning* **92** 34–46
- Salmon J M, Friedl M A, Frolking S, Wisser D and Douglas E M 2015 Global rain-fed, irrigated, and paddy croplands: a new high resolution map derived from remote sensing, crop inventories and climate data *International Journal of Applied Earth Observation and Geoinformation* **38** 321–34
- Schierhorn F, Faramarzi M, Prishchepov A V, Koch F J and Müller D 2014 Quantifying yield gaps in wheat production in Russia *Environ. Res. Lett.* **9** 084017
- Siebert S, Burke J, Faures J M, Frenken K, Hoogeveen J, Döll P and Portmann F T 2010a Groundwater use for irrigation—a global inventory *Hydrol. Earth Syst. Sci.* **14** 1863–80
- Siebert S, Hoogeveen J and Frenken K 2006 Irrigation in Africa, Europe and Latin America—update of the digital global map of irrigation areas to version 4 *Frankfurt Hydrology Paper 05* Institute of Physical Geography, University of Frankfurt, Frankfurt am Main, Germany and Food and Agriculture Organization of the United Nations, Rome, Italy
- Siebert S, Portmann F T and Döll P 2010b Global patterns of cropland use intensity *Remote Sensing* **2** 1625–43
- Skupin A and Agarwal P 2008 Introduction: what is a self-organizing map? *Self-Organising Maps* ed A Skupin and P Agarwal (New York: Wiley) pp 1–20
- Spera S A, Cohn A S, VanWey L K, Mustard J F, Rudorff B F, Risso J and Adami M 2014 Recent cropping frequency, expansion, and abandonment in Mato Grosso, Brazil had selective land characteristics *Environ. Res. Lett.* **9** 064010
- Temme A J A M and Verburg P H 2011 Mapping and modelling of changes in agricultural intensity in Europe *Agriculture, Ecosystems & Environment* **140** 46–56
- Thenkabail P *et al* 2009 Global irrigated area map (GIAM), derived from remote sensing, for the end of the last millennium *Int. J. Remote Sens.* **30** 55
- Tilman D, Balzer C, Hill J and Befort B L 2011 Global food demand and the sustainable intensification of agriculture *Proc. Natl Acad. Sci. USA* **108** 20260–4
- Tilman D, Cassman K G, Matson P A, Naylor R and Polasky S 2002 Agricultural sustainability and intensive production practices *Nature* **418** 671–7
- Tscharntke T, Clough Y, Wanger T C, Jackson L, Motzke I, Perfecto I, Vandermeer J and Whitbread A 2012 Global food security, biodiversity conservation and the future of agricultural intensification *Biological Conservation* **151** 53–9
- Verburg P H, Neumann K and Nol L 2011 Challenges in using land use and land cover data for global change studies *Glob. Change Biol.* **17** 974–89
- Wardlow B D and Egbert S L 2008 Large-area crop mapping using time-series MODIS 250 m NDVI data: an assessment for the US central great plains *Remote Sens. Environ.* **112** 1096–116
- West P C *et al* 2014 Leverage points for improving global food security and the environment *Science* **345** 325–8
- Zaks D P M and Kucharik C J 2011 Data and monitoring needs for a more ecological agriculture *Environ. Res. Lett.* **6** 014017
- Zhang X, Friedl M A, Schaaf C B and Strahler A H 2004 Climate controls on vegetation phenological patterns in northern mid- and high latitudes inferred from MODIS data *Glob. Change Biol.* **10** 1133–45