

Climate change predicted to lengthen transatlantic travel times

This content has been downloaded from IOPscience. Please scroll down to see the full text.

2016 Environ. Res. Lett. 11 031002

(<http://iopscience.iop.org/1748-9326/11/3/031002>)

View [the table of contents for this issue](#), or go to the [journal homepage](#) for more

Download details:

IP Address: 210.77.64.105

This content was downloaded on 05/04/2017 at 03:18

Please note that [terms and conditions apply](#).

You may also be interested in:

[Transatlantic flight times and climate change](#)

Paul D Williams

[Facing unprecedented drying of the Central Andes? Precipitation variability over the period AD 1000–2100](#)

Raphael Neukom, Mario Rohrer, Pierluigi Calanca et al.

[Effects of stratospheric variability on El Niño teleconnections](#)

J H Richter, C Deser and L Sun

[Do responses to different anthropogenic forcings add linearly in climate models?](#)

Kate Marvel, Gavin A Schmidt, Drew Shindell et al.

[Anthropogenic effects on the subtropical jet in the Southern Hemisphere: aerosols versus long-lived greenhouse gases](#)

L D Rotstayn, M A Collier, S J Jeffrey et al.

[Future changes in extratropical storm tracks and baroclinicity under climate change](#)

Jascha Lehmann, Dim Coumou, Katja Frieler et al.

[Influence of climate modes on tropical cyclone formation in the Australian region](#)

A Werner

[Modification of cirrus clouds to reduce global warming](#)

David L Mitchell and William Finnegan

[The economic impact of future increase in tropical cyclones in Japan](#)

Christian Wehersik, M Esteban and T Shibayama

Environmental Research Letters



PERSPECTIVE

Climate change predicted to lengthen transatlantic travel times

OPEN ACCESS

PUBLISHED
25 February 2016

Original content from this work may be used under the terms of the [Creative Commons Attribution 3.0 licence](#).

Any further distribution of this work must maintain attribution to the author(s) and the title of the work, journal citation and DOI.



Keywords: aviation, climate change, atmospheric circulation

Abstract

Among the many consequences of rising atmospheric greenhouse gas concentrations, is a change to the medium through which planes fly. An emerging area of research is the impact that these changes may have on the airline industry, with studies addressing aspects ranging from the impact of rising temperatures on aeroplane weight restrictions (Coffel and Horton 2015 *Wea. Climate Soc.* **7** 94–102) to the impact of changing wind shears on the occurrence of clear air turbulence (Williams and Joshi 2014 *Nat. Clim. Change* **3** 644–8). Now, a recent study by Paul Williams (2016 *Environ. Res. Lett.* **11** 024008), assesses the impact that changing winds may have on the duration of transatlantic flights.

Most people who have taken a round trip flight across the Atlantic are aware that it typically takes longer to travel from east to west than from west to east. This is a consequence of the prevailing westerly winds or jet streams that circle the planet in the mid-latitudes (e.g., figure 1(a)). As the planet warms under rising greenhouse gas concentrations, these winds are expected to change. This has consequences, not only for regional climate and weather in the mid-latitudes, but also, as pointed out by Williams [1], for transatlantic travel times.


The possible impact of changing atmospheric circulation on flight duration was previously assessed in a study last year by Karnauskas *et al* [2] who examined flight durations between Hawaii and the continental US. By comparing interannual variability in the westerly winds with flight duration records, they found, as one might expect, stronger westerly winds favour shorter eastbound journeys and longer westbound journeys. The impacts on the westbound and eastbound legs do not completely cancel, however, leaving a net residual amounting to an overall lengthening of round trip journeys under increased westerly winds. The reason behind this residual remained illusive, although it was key to the overall conclusion that predicted future increases in westerly winds in the East Pacific will lengthen round trip flight durations in that region.

Williams now explains this curious residual through simple arguments, making it clear that, for a flight on a direct great circle path, the residual is a simple consequence of the fact that flight duration is

inversely proportional to the sum of the air speed of the plane and the tail wind speed. strengthened westerly winds in the mid-latitude upper troposphere, as is commonly simulated to occur under climate change (e.g. [3] and figure 1(b)), should then be expected to lengthen round trip transatlantic travel times. As a simple example, take the approximate distance between London and New York as 5500 km, present day westerly wind speeds of 20 m s^{-1} and a future increase in westerly wind speed of 3 m s^{-1} . Assuming the flight time is simply the distance divided by the plane speed plus the tail wind speed, one finds that under the increased westerly wind speed of 3 m s^{-1} , the eastbound journey is shortened by approximately 3 min 53 s, whereas the westbound journey is lengthened by 5 min 16 s. The overall round trip is, therefore, lengthened by 1 min 23 s.

But how does one go about quantifying the magnitude of this effect or the impact it will have? The reality is not as simple as in the above example. The planning of flight routes is complex with the optimal route deviating from the direct great circle path according to day to day variability in the wind field. To consider how flight durations will change in the future, one must not only consider how the winds are expected to change, but also how the optimal flight paths, as determined by flight planners, may change.

To address this issue, Williams applies a flight planning algorithm, used by the airline industry, to examine the optimal routes and flight durations for journeys between London and New York. Wind fields generated from a global climate model under a



doubling of pre-industrial atmospheric CO_2 are fed into this algorithm day by day for 20 years to accumulate statistics of the optimal flight paths and journey times under doubled CO_2 wind conditions. These are then compared with similar calculations performed using wind output from a control simulation at pre-industrial CO_2 levels. Results indicate that round trip journey times will increase throughout the year, with minimum increases during summer of 33 seconds, maximum increases during autumn of 1 min 51 s and an annual average increase of 1 min 6 s. In addition, the probability of eastbound journeys taking under 5 hours 20 minutes is found to increase from 3.5% to 8.1% while the probability of a westbound crossing taking over 7 hours increases from 8.6% to 15.3%.

These changes are small and are unlikely to be noticed by even the most frequent of flyers, but the impact that they could have on fuel consumption and the air industry are not negligible. Considering all transatlantic flights, Williams's estimates that this would amount to an extra 2000 flight hours per year, burning 7.2 million extra US gallons of jet fuel at a cost of \$22 million US dollars to the airline industry and resulting in an additional 70 million kg of CO_2 emitted into the atmosphere. To put these numbers in perspective, US airlines spent a total of \$16.6 billion US dollars on jet fuel for scheduled international flights in 2014¹ and the global airline industry is currently responsible for around 705 billion kg CO_2 emissions annually². So these are relatively small, but non-negligible changes.

The numbers predicted by Williams are not without their uncertainties. A strengthening of the westerly winds in the upper troposphere is expected as a consequence of the fact that, increasing CO_2 concentrations, warm the upper troposphere (and

preferentially so in the tropics) and cool the stratosphere [4]. This leads to a strengthening of the equator-to-pole temperature gradient in the upper troposphere and an associated increase in the westerly winds. However, there is considerable spread among models, as to the exact magnitude and spatial structure of these wind changes, as indicated by figure 1 (c) (see also [3]). The calculations performed by Williams are very computationally intensive and so wind fields from only one such model were used. This uncertain future of the wind field, therefore, leads to uncertainty in the exact values that this calculation produces. Nevertheless, this is one of the first studies that has made a comprehensive attempt to assess the impact that future circulation change may have, through the use of flight planning algorithms. It will no doubt spur future research in this area as well as research into other aspects such as the impact that changes in atmospheric variability or extremes may have on flight delays etc. Such research will improve our overall estimates of the impact that changes to our atmosphere may have on the aviation industry.

References

- [1] Williams P D 2016 Transatlantic flight times and climate change, *Environmental Research Letters Environ. Res. Lett.* **11** 024008
- [2] Karnauskas K B, Donnelly J P, Barkley H C and Martin J E 2015 Coupling between air travel and climate *Nat. Clim. Change*. **5** 1068–73
- [3] Delcambre S C, Lorenz D J, Vimont D J and Martin J E 2013 Diagnosing northern hemisphere jet portrayal in 17 cmip3 global climate models: twenty-first-century projections *J. Clim.* **26** 4930–46
- [4] Manabe S and Wetheral R T 1975 The effect of doubling the CO_2 concentration on the climate of a general circulation model *J. Atmos. Sci.* **32** 3–15

¹ <http://www.transtats.bts.gov/fuel.asp>.

² <http://www.atag.org/facts-and-figures.html>.