

## The UK particulate matter air pollution episode of March–April 2014: more than Saharan dust

This content has been downloaded from IOPscience. Please scroll down to see the full text.

2016 Environ. Res. Lett. 11 044004

(<http://iopscience.iop.org/1748-9326/11/4/044004>)

View [the table of contents for this issue](#), or go to the [journal homepage](#) for more

Download details:

IP Address: 210.77.64.105

This content was downloaded on 31/03/2017 at 11:13

Please note that [terms and conditions apply](#).

You may also be interested in:

[Corrigendum: The UK particulate matter air pollution episode of March–April 2014: more than Saharan dust \(2016 Environ. Res. Lett. 11 044004\)](#)

M Vieno, M R Heal, M M Twigg et al.

[Exceedances of air quality standard level of PM<sub>2.5</sub> in Japan caused by Siberian wildfires](#)

Kohei Ikeda and Hiroshi Tanimoto

[The impact of European legislative and technology measures to reduce air pollutants on air quality, human health and climate](#)

S T Turnock, E W Butt, T B Richardson et al.

[Important role of ammonia on haze formation in Shanghai](#)

Xingnan Ye, Zhen Ma, Jiachen Zhang et al.

[Particulate matter concentration mapping from MODIS satellite data: a Vietnamese case study](#)

Thanh T N Nguyen, Hung Q Bui, Ha V Pham et al.

[Source attribution of particulate matter pollution over North China with the adjoint method](#)

Lin Zhang, Licheng Liu, Yuanhong Zhao et al.

[The impact of synoptic weather on UK surface ozone and implications for premature mortality](#)

R J Pope, E W Butt, M P Chipperfield et al.

[Environmental effects of the recent emission changes in China: implications for particulate matter pollution and soil acidification](#)

Bin Zhao, Shuxiao Wang, Xinyi Dong et al.

## Environmental Research Letters



## CORRIGENDUM

Corrigendum: The UK particulate matter air pollution episode of March–April 2014: more than Saharan dust (2016 *Environ. Res. Lett.* **11** 044004)

## OPEN ACCESS

PUBLISHED  
5 May 2016

Original content from this work may be used under the terms of the [Creative Commons Attribution 3.0 licence](#).

Any further distribution of this work must maintain attribution to the author(s) and the title of the work, journal citation and DOI.



M Vieno<sup>1</sup>, M R Heal<sup>2</sup>, M M Twigg<sup>1</sup>, I A MacKenzie<sup>3</sup>, C F Braban<sup>1</sup>, J J N Lingard<sup>4</sup>, S Ritchie<sup>4</sup>, R C Beck<sup>1</sup>, A Móríng<sup>1,3</sup>, R Ots<sup>1,2</sup>, C F Di Marco<sup>1</sup>, E Nemitz<sup>1</sup>, M A Sutton<sup>1</sup> and S Reis<sup>1,5</sup>

<sup>1</sup> Natural Environment Research Council, Centre for Ecology & Hydrology, Penicuik, UK

<sup>2</sup> School of Chemistry, The University of Edinburgh, UK

<sup>3</sup> School of GeoSciences, The University of Edinburgh, UK

<sup>4</sup> Ricardo Energy and Environment, Harwell, UK

<sup>5</sup> University of Exeter Medical School, European Centre for Environment and Health, Knowledge Spa, Truro, UK

Unfortunately, there was an error in the units of figures 9 and 10 and in the text where the figures were referred to: it should have read ‘ppbv’ instead of ‘ $\mu\text{g m}^{-3}$ ’. The conclusions and interpretation of the results remain unchanged.

## Results and discussion

The paragraph on the altitudinal variation of Saharan dust should have read: ‘To visualise this altitudinal variation in PM composition between the two episodes, a 3D iso-surface of 5 ppbv for  $\text{NO}_3^-$  (green) and Saharan dust (yellow) is shown for 12:00 on 30th March during EP1 (figure 9(b)) and for 3rd April during EP2 (figure 10(b)). Figures 9(a) and 10(a) illustrate the geographic extent of when Saharan dust and  $\text{NO}_3^-$  volume mixing ratio were greater than 1 ppbv for these two occasions.’

The caption of figures 9 and 10 should have read:

**Figure 9.** EMEP4UK modelled Saharan dust (yellow) and  $\text{NO}_3^-$  nitrate (green)  $\text{PM}_{10}$  concentrations at 12:00 on the 30th March 2014: (a) where the volume mixing ratio is greater than 1 ppbv (yellow colour overlays green), and (b) the associated 3D isosurface (5 ppbv). The top of the wireframe box is about 16 km above sea level and the black arrow indicates the north direction. The ppbv mixing ratios for the PM components are based on molecular weights of 62 and 200 for  $\text{NO}_3^-$  and dust, respectively.

**Figure 10.** EMEP4UK modelled Saharan dust (yellow) and particle  $\text{NO}_3^-$  (green)  $\text{PM}_{10}$  concentrations at 12:00 on 3rd April 2014: (a) where the mixing ratio is greater than 1 ppbv (yellow colour overlays green), and (b) the associated 3D isosurface (5 ppbv). The top of the wireframe box is about 16 km above sea level and the black arrow indicates the north direction. The ppbv mixing ratios for the PM components are based on molecular weights of 62 and 200 for  $\text{NO}_3^-$  and dust, respectively.

## Environmental Research Letters



## LETTER

## The UK particulate matter air pollution episode of March–April 2014: more than Saharan dust

## OPEN ACCESS

## RECEIVED

12 October 2015

## REVISED

1 March 2016

## ACCEPTED FOR PUBLICATION

3 March 2016

## PUBLISHED

24 March 2016

M Vieno<sup>1</sup>, M R Heal<sup>2</sup>, M M Twigg<sup>1</sup>, I A MacKenzie<sup>3</sup>, C F Braban<sup>1</sup>, J J N Lingard<sup>4</sup>, S Ritchie<sup>4</sup>, R C Beck<sup>1</sup>, A Móríng<sup>1,3</sup>, R Ots<sup>1,2</sup>, C F Di Marco<sup>1</sup>, E Nemitz<sup>1</sup>, M A Sutton<sup>1</sup> and S Reis<sup>1,5</sup>

<sup>1</sup> Natural Environment Research Council, Centre for Ecology & Hydrology, Penicuik, UK

<sup>2</sup> School of Chemistry, The University of Edinburgh, UK

<sup>3</sup> School of GeoSciences, The University of Edinburgh, UK

<sup>4</sup> Ricardo Energy & Environment, Harwell, UK

<sup>5</sup> University of Exeter Medical School, European Centre for Environment and Health, Knowledge Spa, Truro, UK

E-mail: [vieno.massimo@gmail.com](mailto:vieno.massimo@gmail.com)

**Keywords:** particulate matter, atmospheric chemistry transport model, pollution event

Original content from this work may be used under the terms of the [Creative Commons Attribution 3.0 licence](https://creativecommons.org/licenses/by/3.0/).

Any further distribution of this work must maintain attribution to the author(s) and the title of the work, journal citation and DOI.

**Abstract**

A period of elevated surface concentrations of airborne particulate matter (PM) in the UK in spring 2014 was widely associated in the UK media with a Saharan dust plume. This might have led to over-emphasis on a natural phenomenon and consequently to a missed opportunity to inform the public and provide robust evidence for policy-makers about the observed characteristics and causes of this pollution event. In this work, the EMEP4UK regional atmospheric chemistry transport model (ACTM) was used in conjunction with speciated PM measurements to investigate the sources and long-range transport (including vertical) processes contributing to the chemical components of the elevated surface PM. It is shown that the elevated PM during this period was mainly driven by ammonium nitrate, much of which was derived from emissions outside the UK. In the early part of the episode, Saharan dust remained aloft above the UK; we show that a significant contribution of Saharan dust at surface level was restricted only to the latter part of the elevated PM period and to a relatively small geographic area in the southern part of the UK. The analyses presented in this paper illustrate the capability of advanced ACTMs, corroborated with chemically-speciated measurements, to identify the underlying causes of complex PM air pollution episodes. Specifically, the analyses highlight the substantial contribution of secondary inorganic ammonium nitrate PM, with agricultural ammonia emissions in continental Europe presenting a major driver. The findings suggest that more emphasis on reducing emissions in Europe would have marked benefits in reducing episodic PM<sub>2.5</sub> concentrations in the UK.

**1. Introduction**

An air pollution event of elevated surface concentrations of particulate matter (PM) occurred in the UK between 26 March and 8 April, 2014. Observations from UK air quality monitoring networks showed markedly elevated PM surface concentrations across the majority of the country during this time. The UK Automatic Urban and Rural monitoring Network (AURN) recorded hourly values of PM<sub>10</sub> of up to 100  $\mu\text{g m}^{-3}$ , well above the hourly concentrations of a few tens  $\mu\text{g m}^{-3}$  normally recorded; for example across 45 urban background AURN sites the median, inter-quartile range and 95th percentile of hourly

PM<sub>10</sub> concentrations in 2014 were, respectively, 38, 36–44 and 53  $\mu\text{g m}^{-3}$  (<http://uk-air.defra.gov.uk>).

The spring 2014 PM episode was widely perceived as being associated with an enhanced surface concentration of mineral dust from the Sahara (e.g. as reported by the BBC<sup>6</sup>, the Guardian<sup>7</sup>, and the Daily Mail<sup>8</sup>). An implication is that this was, at least in part, a ‘natural’ event. However, it is known that the UK’s

<sup>6</sup> [www.bbc.co.uk/news/uk-26844425](http://www.bbc.co.uk/news/uk-26844425).

<sup>7</sup> [www.theguardian.com/uk-news/2014/apr/01/london-smog-saharan-dust-storms-downing-street](http://www.theguardian.com/uk-news/2014/apr/01/london-smog-saharan-dust-storms-downing-street).

<sup>8</sup> [www.dailymail.co.uk/wires/ap/article-2594987/Sahara-desert-dust-brings-smog-Britain.html](http://www.dailymail.co.uk/wires/ap/article-2594987/Sahara-desert-dust-brings-smog-Britain.html).

geographic proximity to industrialised countries in Western Europe (e.g. Germany, France and the Netherlands) make it vulnerable to the import of anthropogenic primary and secondary gaseous pollutants and PM. Previous analyses have highlighted this issue for ozone (Vieno *et al* 2010, Francis *et al* 2011) and for secondary inorganic components of PM (Abdalmoghith and Harrison 2005, Jones and Harrison 2006, Yin and Harrison 2008, Charron *et al* 2013, Vieno *et al* 2014). For effective policy action to mitigate the impacts of PM on human health it is essential to understand the composition of the atmosphere and the causes of high levels of PM (e.g. Heal *et al* 2012), of which this episode was an example. Of the studies investigating the origins of elevated PM cited above, all except Vieno *et al* (2014) were based on analysis of measurements of PM chemical components at Harwell or Birmingham in south-central UK. The key feature of the current work was the combination of speciated PM measurements, which are only sparsely undertaken in the UK (as elsewhere), and a regional 3D Eulerian atmospheric chemistry transport model (ACTM) that simulates to 5 km horizontal resolution over the region of study, to assess the validity of attributing this PM event to Saharan dust over the entire UK domain.

## 2. Method

### 2.1. Atmospheric chemistry transport modelling

A Eulerian ACTM, the EMEP4UK version rv4.3 (Vieno *et al* 2010, Simpson *et al* 2012, Vieno *et al* 2014, Vieno *et al* 2016), was used to study the spring 2014 UK air pollution event. The meteorological driver used for EMEP4UK for this purpose was the Weather Research and Forecasting (WRF) model version 3.6. The WRF model included data assimilation (Newtonian nudging) of the numerical weather prediction model meteorological reanalysis from the US National Center for Environmental Prediction/National Center for Atmospheric Research Global Forecast System at 1° resolution, every 6 hours (Saha *et al* 2011). The model domain covers the whole of Europe at a horizontal resolution of 50 km × 50 km with a higher horizontal resolution nested inner domain over the British Isles at 5 km × 5 km, as shown in figure 1.


A full description of the EMEP4UK model can be found in Vieno *et al* (2014) and Simpson *et al* (2012). In addition to gaseous components, the EMEP4UK model simulates the concentrations (at hourly time resolution) of both PM<sub>10</sub> (particles with aerodynamic diameter <10 μm) and PM<sub>2.5</sub> (particles with aerodynamic diameter <2.5 μm), the two PM size fractions forming part of EU air quality legislation (Heal *et al* 2012). The chemical scheme used for the present study is the EMEP-EmChem09 (Simpson *et al* 2012, [www.emep.int](http://www.emep.int)) and the gas/aerosol partitioning was the model for aerosols reacting system (MARS)

(Binkowski and Shankar 1995). In the model version used here, PM<sub>2.5</sub> is derived as the sum of the fine fraction of: ammonium (NH<sub>4</sub><sup>+</sup>), sulphate (SO<sub>4</sub><sup>2-</sup>), nitrate (NO<sub>3</sub><sup>-</sup>), elemental carbon (EC), organic matter (OM), sea salt (SS), mineral dust, and 27% of the coarse nitrate. PM<sub>10</sub> is derived as the sum of PM<sub>2.5</sub> plus the coarse fraction of EC, OM, NO<sub>3</sub><sup>-</sup>, SS, and dust. In the current version of the EMEP4UK model, the split between PM<sub>2.5</sub> and PM<sub>2.5-10</sub> for NO<sub>3</sub><sup>-</sup> is somewhat uncertain as discussed in Aas *et al* (2012). Fine NO<sub>3</sub><sup>-</sup> production is modelled using the MARS thermodynamic model and the coarse NO<sub>3</sub><sup>-</sup> formation in the current model is a function of humidity and HNO<sub>3</sub> using a parametrization derived from Riemer *et al* (2003), as described in Simpson *et al* (2012), but it is not explicitly linked to the surface area of the existing coarse aerosol. Both nitrate generation mechanisms compete for the same HNO<sub>3</sub> therefore constraining the total amount of nitrate produced.

Anthropogenic emissions of NO<sub>x</sub>, NH<sub>3</sub>, SO<sub>2</sub>, primary PM<sub>2.5</sub>, primary coarse PM, CO and non-methane VOC for the UK are derived from the National Atmospheric Emission Inventory (NAEI, <http://naei.defra.gov.uk>) aggregated to 5 km × 5 km resolution. The EMEP 50 km × 50 km resolution emission estimates provided by the Centre for Emission Inventories and Projections (CEIP, [www.ceip.at/](http://www.ceip.at/)) are used for the European domain. Shipping emissions estimates, for the inner domain, are derived from ENTEC (2010).

The boundary conditions at the edge of the European domain (figure 1) are prescribed concentrations adjusted for each simulated year as described in Simpson *et al* (2012). The EMEP4UK model version used here uses the default climatological Saharan dust boundary conditions, based on monthly average dust concentrations for the year 2000 from the global CTM2 at the University of Oslo. This treatment may lead to an inaccurate representation of the source strength of Saharan dust for individual events. The atmospheric flow and deposition processes transporting the dust and controlling its distribution over Europe and the UK during the simulated period are fully represented within the model. For ozone, a 3D field from climatological ozone-sonde data-sets, modified monthly against clean-air surface observations was used as initial and boundary condition for the European domain.

Two EMEP4UK model simulations were carried out to quantify the contributions of UK and regional emissions to the episode of elevated surface PM: (1) A base-case run, called 'BASE' hereafter, and (2) a no-UK anthropogenic emissions run, called 'NoUK' hereafter. The difference between the BASE and the NoUK simulations therefore represents a quantitative measure of the enhancement of the UK PM concentration that is caused by domestic emissions over the European background. The percentage fraction of PM generated by UK emissions ( $f_{UK}$ ) was calculated as:



$$f_{\text{UK}} = 100 \times (\text{BASE} - \text{NoUK}) / \text{BASE}. \quad (1)$$

## 2.2. Measurements


The daily-mean simulated surface concentrations of PM<sub>10</sub> and PM<sub>2.5</sub> across the UK were compared with data from the UK's AURN, which uses the Tapered Element Oscillating Microbalance Filter Dynamic Measurement System (TEOM-FDMS) (Bureau-Veritas 2010). Comparisons were undertaken for all AURN sites with PM<sub>10</sub> and PM<sub>2.5</sub> measurements; for the sake of simplicity this paper presents data for the following four representative sites (uk-air.defra.gov.uk): Aberdeen (North-East Scotland, lat 57.157°, long -2.094°, urban background); Auchencorth Moss (Central Scotland, lat 55.792°, long -3.243°, rural background); Harwell (South-East England, lat 51.571°, long -1.325°, rural background); and London Bloomsbury (Greater London, lat 51.522°, long -0.126°, urban background). The locations of these sites are marked on figure 1. The Harwell and Auchencorth Moss sites are part of the UKEAP (UK Eutrophying and Acidifying Atmospheric Pollutant) network. For this purpose, additional instrumentation includes a Monitor for AeRosols and GAses (MARGA, Metrohm Applikon B. V., NL), which measures hourly concentrations of the major secondary inorganic components of PM (SO<sub>4</sub><sup>2-</sup>, NO<sub>3</sub><sup>-</sup>, and NH<sub>4</sub><sup>+</sup>), as described by Twigg *et al* (2015).

## 3. Results and discussion

The dominant meteorological feature for the duration of the PM pollution event was an area of high pressure located over Europe, with relatively low surface temperatures of 8 °C–14 °C. This limited the evaporation of ammonium nitrate into ammonia and nitric acid, enhancing the lifetime of particle nitrate and the potential influence of its long-range transport.

Figures 2 and 3 compare the modelled and measured daily mean surface concentrations of PM<sub>2.5</sub> and PM<sub>10</sub>, respectively, between 20th March and 13th April 2014 at the four selected AURN sites. All four sites experienced elevated PM during this period when compared with their 2014 annual averages, which for PM<sub>2.5</sub> and PM<sub>10</sub> were: 9 and 14 μg m<sup>-3</sup> (Harwell), 15 and 19 μg m<sup>-3</sup> (London Bloomsbury), 10 and 15 μg m<sup>-3</sup> (Aberdeen) and 7 and 8 μg m<sup>-3</sup> (Auchencorth Moss) (uk-air.defra.gov.uk). The correlation coefficients between model and measurements over the simulation period are greater than 0.77 for all four sites included in this work for both PM<sub>2.5</sub> and PM<sub>10</sub> (figures 2 and 3).

Two distinct episodes are apparent in the PM<sub>2.5</sub> and PM<sub>10</sub> concentrations: one centred on the 30th March (referred to subsequently as episode 1: 'EP1') with an approximate duration of 5 days, and the second centred on the 3rd April ('EP2') with an approximate duration of 4 days. The spatial distributions across the UK and the rest of Western Europe of the daily mean surface PM during these episodes are




**Figure 2.** EMEP4UK modelled daily-mean components of surface PM<sub>2.5</sub> (coloured bars) from 20th March to 12th April 2014 for four AURN sites: Aberdeen (North-East Scotland) and London Bloomsbury (Greater London), both urban background sites; and Auchencorth Moss (Central Scotland) and Harwell (South-East England), both rural background sites. The solid black lines show the observed AURN daily-mean PM<sub>2.5</sub> concentrations. The linear regression between observation and model is shown at the top of each panel, along with the correlation coefficient,  $r$ .

shown in figures 4 and 5 for PM<sub>2.5</sub> and PM<sub>10</sub>, respectively.

Figures 2 and 3 show that during the full period presented, PM<sub>2.5</sub> and PM<sub>10</sub> were dominated by secondary inorganic PM components, especially for PM<sub>2.5</sub>. The estimated proportions of daily-mean PM<sub>2.5</sub> and PM<sub>10</sub> derived from UK emissions are shown in figures 6(a) and 7(a), respectively. The proportions were typically less than 40% from which it can be inferred that EP1 and EP2 both have a large component of PM derived from long-range transport

of secondary inorganic PM and secondary PM precursors. In contrast, on the 1st April, a date between the two episodes, up to 70% of PM<sub>2.5</sub> and PM<sub>10</sub> were estimated to have been generated from UK emissions. The contribution of transboundary transport to PM pollution events in the UK has been reported before; for example, Vieno *et al* (2014) showed that up to 80% of particle NO<sub>3</sub><sup>-</sup> in episodes may be imported into the UK from continental Europe.

To further test the model, figure 8 shows the hourly modelled inorganic composition of PM<sub>10</sub> at




**Figure 3.** EMEP4UK modelled daily-mean components of surface PM<sub>10</sub> (coloured bars) from 20th March to 12th April 2014 for four AURN sites: Aberdeen (North-East Scotland) and London Bloomsbury (Greater London), both urban background sites; and Auchencorth Moss (Central Scotland) and Harwell (South-East England), both rural background sites. The solid black line shows the observed AURN daily-mean PM<sub>10</sub> concentration. The linear regression between observation and model is also shown at the top of each panel, along with the correlation coefficient,  $r$ .

Auchencorth Moss and at Harwell over the full time period compared against the MARGA measurements ([uk-air.defra.gov.uk](http://uk-air.defra.gov.uk)). Both the speciated MARGA measurements and the model follow the TEOM-FDMS PM<sub>10</sub> temporal variation shown in figure 3 relatively well. Gaps in the measurements indicate when the MARGA instrument was offline. As for total daily PM<sub>2.5</sub> and PM<sub>10</sub>, the hourly observations of speciated PM<sub>10</sub> components during EP1 and EP2 are well captured by the model, both in timing and magnitude. The notable enhancement of NO<sub>3</sub><sup>-</sup> concentrations during this event is clearly evident in both the MARGA observations and in the model simulations as shown in figure 8. The fact that the NO<sub>3</sub><sup>-</sup> enhancement was

accompanied by an equivalent enhancement in NH<sub>4</sub><sup>+</sup> supports the conclusion that observed PM<sub>2.5</sub> enhancements are minimally impacted by dust or by NO<sub>3</sub><sup>-</sup> formation on dust surfaces.

With respect to the Saharan dust component, the modelling indicates that the PM<sub>2.5</sub> and PM<sub>10</sub> composition differed between EP1 and EP2. Figures 6(b) and 7(b) show that Saharan dust constituted a greater component of both surface PM<sub>2.5</sub> and PM<sub>10</sub> during EP2 than during EP1. However, the dust was present at substantial surface concentrations only in the southern UK, as illustrated by the data presented here for the representative southern UK monitoring sites at Harwell and London Bloomsbury. Overall, the




percentage of modelled Saharan dust in  $PM_{2.5}$  and  $PM_{10}$  is less than 20% for most of the UK for both EP1 and EP2. The geographic extent of the Saharan dust incursion to the surface (relative dust contribution to  $PM > 20\%$ ) for EP2 is shown in figures 6(b) and 7(b) for  $PM_{2.5}$  and  $PM_{10}$ , respectively. During EP1 the high concentrations of PM in the plume were derived from anthropogenic emissions in continental Europe and advected to the UK as shown in figures 4 and 5. It was only during EP2 that the Saharan dust plume was also present at the surface (for parts of the UK) augmenting the secondary PM from continental Europe. The Saharan dust plume can be observed having evolved and moving north across Europe from the 30th March, arriving at the UK surface level on the 3rd of April (figure 2). Saharan dust in the atmosphere was also predicted by the model during EP1, but at higher altitudes with little impact at the surface. To visualise this altitudinal variation in PM composition between the two episodes, a 3D iso-surface of  $\sim 5 \mu\text{g m}^{-3}$  at the surface for  $\text{NO}_3^-$  (green) and Saharan dust (yellow) is shown for 12:00 on 30th March during EP1 (figure 9(b)) and for 3rd April during EP2 (figure 10(b)). Figures 9(a) and 10(a) illustrate the geographic extent of when Saharan dust and  $\text{NO}_3^-$  concentrations at the surface were greater than  $1 \mu\text{g m}^{-3}$  for these two occasions. Figure 9 shows that during

EP1 there was a low-altitude  $\text{NO}_3^-$  plume advected in from continental Europe and a separate high altitude plume of Saharan dust. The model simulation shows that during EP1 the high altitude Saharan dust did not reach the UK surface apart from a small area in Wales as shown in figure 9(a). In contrast, figure 10 shows that for EP2 the Saharan dust was present at lower altitudes and had reached the surface in the south of the UK on the 3rd of April (the dust plume had also reached the surface in the UK on 2nd April).

From the analysis of these two PM events, which although temporally adjacent have distinct characteristics due to their air mass history, it has been shown that speciation measurements and modelling can allow the clear, quantitative diagnosis of the contributions to PM mass. In this case, the MARGA measurements characterised the inorganic composition and the modelling demonstrated how that composition developed. These measurements do not provide explicit information about the external/internal mixing properties of the PM, e.g. interactions of the secondary PM on the surfaces of mineral dust, which is likely to be occurring as noted in previous studies (Fairlie *et al* 2010). However, since in the EMEP4UK model the production of  $\text{NH}_4\text{NO}_3$  is not linked to the additional particle surface area provided by the dust, the model's ability to reproduce measured concentration





of these compounds suggests that the importance of the Saharan dust promoting secondary aerosol formation is not a key process for this event.


The modelled low levels of Saharan dust in EP1 (<20% of PM mass) coupled with the PM measurements indicate that the UK did not experience the consequences of a Saharan dust episode at that time. Rather, this was mainly an anthropogenic pollution event, exacerbated by the transport of secondary inorganic aerosol precursors emitted outside of the UK, which, in principle, could be mitigated against in future with appropriate international policies to reduce primary air pollutant emissions (especially  $\text{NO}_x$  and  $\text{NH}_3$ ). The importance of long-range transport of ammonium nitrate demonstrated here for this UK  $\text{PM}_{2.5}$  episode corroborates previous work (Vieno *et al* 2014) and confirms similar findings on a European scale presented by Kieseewetter *et al* (2015). The policy implications derived from accurate identification of emission sources contributing to PM concentrations are discussed further in the next section.

#### 4. Policy implications

The findings presented in this paper illustrate the capability of advanced ACTMs to reliably identify the

underlying causes of complex air pollution episodes. The validation of the model results against chemically-specified measurements of PM components further highlights the need for the development and implementation of better integrated model and measurement systems to routinely apportion sources and target the most relevant contributing activities for the design of effective pollution control policies (Reis *et al* 2015). It is also clear that it is feasible to provide robust information on the state of atmospheric composition as an evidence base to policy makers in a more operational manner.


The substantial contribution of long-range transport to ambient concentrations of fine PM in the UK, and the accurate identification of the relevant sources, has far-reaching policy implications. The pollution episode in spring 2014 was widely attributed to Saharan dust in the UK media, thus placing a (false) emphasis on a natural phenomenon, which cannot be addressed by policy action. The focus on an ‘exciting headline’ may be attributable to the complexity of communicating atmospheric processes and the public observations of dust-fall on surfaces. This suggests that while scientific analyses to determine the real causes of particular PM events through measurements and modelling are well established, the translation into communicating outside the atmospheric science arena



needs improvement. Therefore, there is a danger that important messages to high-level policy-makers may be obscured by attention-catching headlines.


The initial mischaracterisation of the PM event may also represent a missed opportunity to inform and educate the general public about the role of anthropogenic emissions, specifically agricultural

emissions of ammonia, as a key contributor to many high PM pollution events in recent years. As a consequence, the lack of public awareness may result in reduced public interest in, and a weaker policy mandate for, any emissions reduction targets for ammonia emissions and their contribution to transboundary fluxes of secondary aerosol components. The



importance of public support has been illustrated by Reis *et al* (2012) specifically for the implementation of air pollution control policies through a robust science-policy interface. The extent to which explicit value

judgements are a further underlying cause for the—to date—only moderate ambition levels in reducing ammonia emissions on a European scale (Voinov *et al* 2014) is difficult to quantify.





**Figure 8.** Hourly means calculated by the EMEP4UK model (dashed lines) and measured by the MARGA instruments (solid lines) at Auchencorth Moss (top) and Harwell (bottom) for the major secondary inorganic components of  $PM_{10}$ :  $NH_4^+$  orange lines,  $NO_3^-$  blue lines and  $SO_4^{2-}$  red lines.

These implications are not only relevant for Europe. The PM pollution episodes in China and other fast growing economies (e.g. Oh *et al* 2015, Yan *et al* 2015, Zhang *et al* 2015, Zhou *et al* 2015, Tan *et al* 2016) are more severe examples of those in the UK, Paris, and other parts of Europe (Gualtieri *et al* 2015). While a major part of urban  $PM_{2.5}$  in these other regions arises from primary  $PM_{2.5}$  emissions from local transport and industrial activities, as has also been demonstrated for the UK (Vieno *et al* 2016), agricultural ammonia emissions have also been identified as key contributors. In these regions also, as for Europe, the need for close collaboration in policy development across provincial or national borders is evident and needs robust modelling and monitoring activities to provide underpinning scientific evidence. Such transboundary collaboration is emerging, e.g. in the case of tackling air quality and ozone pollution in Hong Kong and the Pearl River Delta (Zhong *et al* 2013). The combination of local measures and regional policies to reduce emissions within and far away from urban

areas will likely be more effective in reducing the public health impacts of severe episodes of PM.

## 5. Conclusions

The EMEP4UK regional ACTM has been shown to successfully reproduce elevated surface PM concentrations during the spring of 2014 in the UK. The combined analysis of model results and speciated PM measurements indicates that these elevated PM concentrations were mainly driven by ammonium nitrate, much of which was derived from European  $NO_x$  and  $NH_3$  emissions outside the UK. The contribution of Saharan dust, which was widely reported in the UK media, has been shown to be restricted (at surface level) to the latter part of the elevated PM episode and regionally to the southern UK. Both types of PM and PM precursors arrived in the UK with the same southerly air flow, but originated from quite independent sources. The analyses presented in this paper



illustrate the complementary use of high resolution observations (temporal and chemical speciation) and an advanced ACTM to identify the underlying causes of complex PM air pollution episodes. Specifically, the findings suggest that more emphasis on reducing

ammonia emissions in Europe would have marked benefits in reducing episodic  $\text{PM}_{2.5}$  concentrations and population exposure in the UK. The need for close collaboration in policy development across provincial or national borders applies worldwide.

## Acknowledgments

The authors acknowledge the UK Department for Environment, Food and Rural Affairs (Defra) and the Devolved Administrations, through the projects: development of the EMEP4UK model (AQ0727), the 'UK Eutrophying and Acidifying Atmospheric Pollutants (UKEAP) project' (AQ0647) which funds the MARGA observations, as well as access to the AURN data, which were obtained from [uk-air.defra.gov.uk](http://uk-air.defra.gov.uk) and are subject to Crown 2014 copyright, Defra, licenced under the Open Government Licence (OGL). Partial support for the EMEP4UK modelling from the European Commission FP7 ECLAIRE project is gratefully acknowledged.

## References

- Aas W *et al* 2012 Lessons learnt from the first emep intensive measurement periods *Atmos. Chem. Phys.* **12** 8073–94
- Abdalmoghith S S and Harrison R M 2005 The use of trajectory cluster analysis to examine the long-range transport of secondary inorganic aerosol in the UK *Atmos. Environ.* **39** 6686–95
- Binkowski F S and Shankar U 1995 The regional particulate matter model: I. Model description and preliminary results *J. Geophys. Res.-Atmos.* **100** 26191–209
- Bureau-Veritas 2010 Assessment of UK AURN particulate matter monitoring equipment against the January 2010 guide to demonstration of equivalence, Defra ([http://uk-air.defra.gov.uk/assets/documents/reports/cat14/1101140842\\_Assessment\\_of\\_UK\\_AURN\\_PM\\_Equipment\\_against\\_2010\\_GDE.pdf](http://uk-air.defra.gov.uk/assets/documents/reports/cat14/1101140842_Assessment_of_UK_AURN_PM_Equipment_against_2010_GDE.pdf))
- Charron A, Degrendele C, Laongsri B and Harrison R M 2013 Receptor modelling of secondary and carbonaceous particulate matter at a southern UK site *Atmos. Chem. Phys.* **13** 1879–94
- ENTEC 2010 Defra, UK ship emissions inventory, final report ([http://uk-air.defra.gov.uk/reports/cat15/1012131459\\_21897\\_Final\\_Report\\_291110.pdf](http://uk-air.defra.gov.uk/reports/cat15/1012131459_21897_Final_Report_291110.pdf)) Crown copyright
- Fairlie T D, Jacob D J, Dibb J E, Alexander B, Avery M A, van Donkelaar A and Zhang L 2010 Impact of mineral dust on nitrate, sulfate, and ozone in transpacific asian pollution plumes *Atmos. Chem. Phys.* **10** 3999–4012
- Francis X V, Chemel C, Sokhi R S, Norton E G, Ricketts H M A and Fisher B E A 2011 Mechanisms responsible for the build-up of ozone over south east England during the August 2003 heatwave *Atmos. Environ.* **45** 6880–90
- Gualtieri G, Toscano P, Crisci A, Di Leonardo S, Tartaglia M, Vagnoli C, Zaldei A and Gioli B 2015 Influence of road traffic, residential heating and meteorological conditions on PM10 concentrations during air pollution critical episodes *Environ. Sci. Pollut. R* **22** 19027–38
- Heal M R, Kumar P and Harrison R M 2012 Particles, air quality, policy and health *Chem. Soc. Rev.* **41** 6606–30
- Jones A M and Harrison R M 2006 Assessment of natural components of PM10 at UK urban and rural sites *Atmos. Environ.* **40** 7733–41
- Kiesewetter G, Borken-Kleefeld J, Schopp W, Heyes C, Thunis P, Bessagnet B, Terrenoire E, Fagerli H, Nyiri A and Amann M 2015 Modelling street level PM10 concentrations across Europe: source apportionment and possible futures *Atmos. Chem. Phys.* **15** 1539–53
- Oh H R, Ho C H, Kim J, Chen D L, Lee S, Choi Y S, Chang L S and Song C K 2015 Long-range transport of air pollutants originating in China: a possible major cause of multi-day high-PM10 episodes during cold season in Seoul, Korea *Atmos. Environ.* **109** 23–30
- Reis S *et al* 2012 From acid rain to climate change *Science* **338** 1153–4
- Reis S *et al* 2015 Integrating modelling and smart sensors for environmental and human health *Environ. Modell. Softw.* **74** 238–46
- Riemer N, Vogel H, Vogel B, Schell B, Ackermann I, Kessler C and Hass H 2003 Impact of the heterogeneous hydrolysis of N<sub>2</sub>O<sub>5</sub> on chemistry and nitrate aerosol formation in the lower troposphere under photo-smog conditions *J. Geophys. Res.-Atmos.* **108** 4144
- Saha S *et al* 2011 NCEP Climate Forecast System Version 2 (cfsv2) 6-hourly Products Research Data Archive at the National Center for Atmospheric Research, Computational and Information Systems Laboratory, Boulder, CO
- Simpson D *et al* 2012 The EMEP MSC-W chemical transport model—technical description *Atmos. Chem. Phys.* **12** 7825–65
- Tan J H, Duan J C, Zhen N J, He K B and Hao J M 2016 Chemical characteristics and source of size-fractionated atmospheric particle in haze episode in Beijing *Atmos. Res.* **167** 24–33
- Twigg M M *et al* 2015 Water soluble aerosols and gases at a UK background site: I. Controls of pm2.5 and PM10 aerosol composition *Atmos. Chem. Phys.* **15** 8131–45
- Vieno M *et al* 2010 Modelling surface ozone during the 2003 heatwave in the UK *Atmos. Chem. Phys.* **10** 7963–78
- Vieno M *et al* 2014 The role of long-range transport and domestic emissions in determining atmospheric secondary inorganic particle concentrations across the UK *Atmos. Chem. Phys.* **14** 8435–47
- Vieno M, Heal M R, Williams M L, Carnell E J, Nemitz E, Stedman J R and Reis S 2016 The sensitivities of emissions reductions for the mitigation of UK pm2.5 *Atmos. Chem. Phys.* **16** 265–76
- Voinov A, Seppelt R, Reis S, Nabel J E M S and Shokravi S 2014 Values in socio-environmental modelling: persuasion for action or excuse for inaction *Environ. Modell. Softw.* **53** 207–12
- Yan R C, Yu S C, Zhang Q Y, Li P F, Wang S, Chen B X and Liu W P 2015 A heavy haze episode in Beijing in February of 2014: characteristics, origins and implications *Atmos. Pollut. Res.* **6** 867–76
- Yin J X and Harrison R M 2008 Pragmatic mass closure study for pm1.0, pm2.5 and PM10 at roadside, urban background and rural sites *Atmos. Environ.* **42** 980–8
- Zhang Q Y, Yan R C, Fan J W, Yu S C, Yang W D, Li P F, Wang S, Chen B X, Liu W P and Zhang X Y 2015 A heavy haze episode in Shanghai in December of 2013: Characteristics, origins and implications *Aerosol Air Qual. Res.* **15** 1881–93
- Zhong L J, Louie P K K, Zheng J Y, Yuan Z B, Yue D L, Ho J W K and Lau A K H 2013 Science-policy interplay: air quality management in the Pearl River Delta region and Hong Kong *Atmos. Environ.* **76** 3–10
- Zhou M G *et al* 2015 Smog episodes, fine particulate pollution and mortality in China *Environ. Res.* **136** 396–404