

Steps toward nation-wide monitoring of non-indigenous species in Danish marine waters under the Marine Strategy Framework Directive



NIVA Denmark Water Research
– a subsidiary of the Norwegian Institute for Water Research

REPORT

Main Office	NIVA Region South	NIVA Region East	NIVA Region West	NIVA Denmark
Gaustadalléen 21 NO-0349 Oslo, Norway Phone (47) 22 18 51 00 Telefax (47) 22 18 52 00 Internet: www.niva.no	Jon Lilletuns vei 3 NO-4879 Grimstad, Norway Phone (47) 22 18 51 00 Telefax (47) 37 04 45 13	Sandvikaveien 59 NO-2312 Ottestad, Norway Phone (47) 22 18 51 00 Telefax (47) 62 57 66 53	Thormohlens gate 53 D NO-5006 Bergen Norway Phone (47) 22 18 51 00 Telefax (47) 55 31 22 14	Ørestads Boulevard 73 2300 Copenhagen S Phose (45) 88 96 96 70 www.niva-danmark.dk

Title Steps toward nation-wide monitoring of non-indigenous species in Danish marine waters under the Marine Strategy Framework Directive	Report No. 7022-2016-DK3	Date 7 April 2016
Author(s) Jesper H. Andersen, NIVA Denmark Water Research Emilie Kallenbach, NIVA Denmark Water Research Martin Hesselsøe, Amphi Consult Aps Steen W. Knudsen, Amphi Consult Aps / Natural History Museum Denmark Peter Rask Møller, Natural History Museum Denmark Dorte Bekkevold, DTU Aqua Brian Klitgaard Hansen, DTU Aqua Jens Thaulow, Norwegian Institute for Water Research	Project No. 15389	Pages 122 Price -
Topic group MSFD NIS	Geographical area Denmark North Sea Baltic Sea	Distribution Public

Client(s) Danish Nature Agency	Client ref. UCB
---------------------------------------	--------------------

Abstract This report is the outcome of MONIS 2 – or in full, “Monitoring of Non-Indigenous Species in Danish Marine Water, phase 2” – and includes three deliverable: (1) a national Target Species List including 50 species, (2) a draft Technical Guidance Report, and (3) <i>in silico</i> designed and tested primers and probes for 48 of the 50 species on the Target Species List. The list is based on discussions at a workshop and subsequent scoring and ranking of relevant species. The draft Technical Guidance report is anchored in existing Standard Operating Procedures (i.e. protocols for sampling, storage and analysis) and adapted to the requirements of the Danish NOVANA programme. In addition, the report includes suggestions for next steps to take to implement and improve monitoring and assessment activities in regard to non-indigenous species in Danish marine waters.

4 keywords, Danish	4 keywords, English
1. Havstrategidirektivet 2. Ikke-hjemmehørende arter 3. Overvågning 4. eDNA-baserede metoder	1. Marine Strategy Framework Directive (MSFD) 2. Non-indigenous species (NIS) 3. Monitoring 4. eDNA based methods

Chief Scientist
Jesper Andersen

Research Director
Nikolai Friberg

ISBN 978-82-577-6757-0

MONIS phase 2

**Steps toward nation-wide monitoring of
non-indigenous species in Danish marine waters
under the Marine Strategy Framework Directive**

Client: Danish Nature Agency

Preface

We report the outcome of the project “Monitoring of Non-Indigenous Species in Danish Marine Waters, phase 2” (MONIS 2), which has been initiated and funded by the Danish Nature Agency. The report follows up on MONIS 1, which is published in 2014 by the Danish Nature Agency:

- Andersen, J.H., S.A. Pedersen, J. Thaulow, F. Stuer-Lauridsen & S. Cochrane (2014): Monitoring of non-indigenous species in Danish marine waters. Background and proposals for a monitoring strategy and a monitoring network. Danish Nature Agency. 55 pp.

Three key deliverables included in the report:

- A national Target Species List including 50 species,
- A draft Technical Guidance Report describing in detail how to carry out sampling, storage and analyses, and
- Species-specific *in silico* designed and tested primers- and probe systems for 48 of the 50 species on the Target Species List. All systems have been matched and compared with nucleotide sequences available on the NCBI GenBank database.

In addition to the above, we propose next steps which are meant as a road map for the implementation of national monitoring and assessment activities in regard to Descriptor 2 specific requirements (i.e. non-indigenous species; NIS) under the EU Marine Strategy Framework Directive (MSFD).

The authors would like to thank Henrik Carl, Natural History Museum Denmark for permission to use the photo on the front cover as well as Norman Green for a critical review of an earlier version of this report.

Copenhagen, 7 April 2016

Jesper H. Andersen

Contents

Summary	4
1. Introduction	5
2. What did we do?	6
2.1 Towards a national MSFD D2 Target Species List	6
2.2 Technical Guidance Report for eDNA-based monitoring activities	7
2.3 Development of an eDNA-based single-species detections system	8
3. Results	9
3.1 Target Species List	9
3.2 Technical Guidance Report	9
3.3 A first generation of MSFD D2-related primers	9
4. Discussion and recommendations	11
5. References	13
Appendix A: Minutes of the TSL workshop	16
Appendix B: Draft Technical Guidance Report	19
Appendix C: <i>In silico</i> developed primers	36
Appendix D: Ranking of species	113

Summary

The project “Monitoring of Non-Indigenous Species in Danish Marine Waters, phase 2” (MONIS 2) has been initiated and funded by the Danish Nature Agency in order to follow up on the proposed monitoring strategy and tentatively outlined design of a national monitoring programme targeting non-indigenous species in Danish marine waters. The monitoring activities are anchored in requirement in the EU Marine Strategy Framework Directive (MSFD), specifically in regard to Descriptor 2 concerning non-indigenous species (D2).

In order to make proposals suggested by MONIS 1 operational work has focused on three issues:

1. development of a national MSFD D2-specific Target Species List including 50 species,
2. development and writing of a draft technical Guidance Report based on the existing National Monitoring and Assessment Programme for the Aquatic and Terrestrial Environments (NOVANA) principles and structure for such reports, and
3. *in silico* (i.e. computer-based analysis) to compare nucleotide sequences for the upcoming development of primers for as many of the species on the Targets Species List as possible.

The Target Species List, the *in silico* developed primers and the Technical Guidance Report are included in this report. In combination, they represent important building blocks towards the initiation of nation-wide sampling covering both coastal waters and open marine areas in late spring / early summer 2016.

The process and strategy outlined by the MONIS 1 report can turn into a more practical phase where a Danish MSFD D2-specific monitoring programme can now be implemented.

1. Introduction

Danish marine waters are impacted by multiple human stressors, e.g. nutrient inputs (Ærtebjerg *et al.* 2003, Andersen & Conley 2009), fishing activities (Korpinen *et al.* 2012, Andersen & Stock (eds.) 2013), inputs of hazardous substances (Dahlöf & Andersen 2009), as well as physical modification (Andersen *et al.* 2012). The ecosystem health of Danish marine waters is accordingly assessed as being impaired (Naturstyrelsen 2012), especially in regard to eutrophication (Riemann *et al.* 2015, Andersen *et al.* 2015), hazardous substances (Andersen *et al.* 2016), biodiversity (Andersen *et al.* 2014a) as well as ‘ecosystem health’ (Naturstyrelsen 2012).

Other human activities might also be important, e.g. introductions of non-indigenous species, but there is currently only scarce information about introduction and occurrence of non-native species in Danish marine waters (e.g. Stæhr & Thomsen 2012, Azour *et al.* 2015).

However, the EU Marine Strategy Framework Directive (MSFD) call for an ecosystem-based approach to the management of all European marine waters (Anon. 2008) and consequently Denmark, as well as all EU Member States with a coastline, are required to carry out so-called ‘Initial Assessment’ of pressures and environmental status and to implement specific monitoring activities.

The assessment in regard to ‘good environmental status’ (GES), being an overarching target to achieve in a MSFD-context, is nested, and the groups of indicators addressed are termed Descriptors. In total there are 11 descriptors and the GES definition for descriptor 2 (D2) is “Non-indigenous species introduced by human activities do not adversely alter the ecosystem” (Anon. 2010).

In Denmark, the Nature Agency has initiated a process aiming at a formulation of both a strategy for MSFD D2 monitoring as well as an interim proposal for a monitoring programme. This was done within the MONIS project (Andersen *et al.* 2014b). In order to follow up on this strategy and proposal for a monitoring programme, the Nature Agency published a tender requesting three key products:

1. the development of a national MSFD D2-specific Target Species List,
2. production of a draft Technical Guidance Report describing in details how to carry out sampling, storage and lab analyses, and
3. *in silvo* development and testing of as many species-specific primer-probe assays as possible with regard to the national MSFD D2 Target Species List.

The tender was won by a partnership comprising NIVA Denmark Water Research, AmphiConsult, DTU Aqua and the University of Copenhagen/National Natural History Museum of Denmark as well as an associate partner being Eurofins Miljø AS.

In addition to the above three key products, the report also includes a range of suggestions for the next steps to arrive at an operational MSFD D2-specific monitoring programme.

2. What did we do?

The processes, principles and methods used for (1) development of a national MSFD D2-specific Target Species List, (2) drafting of a Technical Guidance Report, and (3) *in silico* development and testing of as many species-specific primer-probe assays as possible, are described in the following sections.

2.1 Towards a national MSFD D2 Target Species List

The ranking of marine species for a national MSFD D2 target list is based on different authorities' registrations and listing of non-indigenous and/or invasive species in the Danish marine areas (Naturstyrelsen 2008a, 2008b, Stæhr & Thomsen 2012) or within the boundaries of European seas (HELCOM 2015, OSPAR 2015). Additionally, two other lists established by Jensen (2013) and Møller (2015) were included in the making of the target species list, in order to fill in the blanks. As it appears from the overall list, the different lists are in accordance with one another.

Table 1: Summary of the species lists on which the national Target Species List is based.

Full reference
1. Berggren, U. (<i>pers. comm.</i>): Draft short list of marine non-indigenous species.
2. HELCOM (2015): List of Target Species currently in use in the HELCOM area. 1 pp.
3. Jensen, K. (2013): Target Species List 1 (established species) and Target Species List 2 (alert list). 2 pp.
4. Møller, P.R. (<i>pers. comm.</i>): Supplerende liste over marine fisk. (In Danish)
5. Naturstyrelsen (2008a): Sortlisten. Akvatiske arter. 1 pp.
6. Naturstyrelsen (2008b): Observationslisten. Akvatiske arter. 1 pp.
7. OSPAR (2015): OSPAR Target Species List (OSPAR Agreement 2015-10). 2 pp.
8. Stæhr, P.A. & Thomsen, M.S. (2012): Opgørelse over rumlig udbredelse, tidslig udvikling og tæthed af ikke-hjemmehørende arter i danske farvande. Fagligt notat fra DCE - Nationalt Center for Miljø og Energi. 14 pp.
9. Jensen, K. (<i>pers. comm.</i>)

Ranking of the target species is based on the following five criteria:

1. established non-indigenous species (NIS),
2. potential NIS,
3. invasiveness,
4. ease of determination by conventional methods, and
5. ease of determination by eDNA techniques.

These criteria are described in the following sections:

- **Criterion 1 – Established NIS:** The scores are based on the list of established species in Table 1. If a species is registered on one of the lists, it gets the score 8, whereas it is assigned the score 10, if it is appears on two lists (Jensen, K. (*pers. Comm.*) and Stæhr & Thomsen (2012)). If it is not registered on any of those two, it gets the score 0, and is considered as not yet established in Denmark (cf. Table 1).
- **Criterion 2 – Potential NIS:** If not yet established, a species' potential invasiveness is assessed. This assessment is rooted in the lists from OSPAR (2015) and HELCOM (2015). OSPAR and HELCOM cover the North-East Atlantic and Baltic Sea respectively and thereby indicate the largest dispersal areas for this study. For each of the lists the species occurs on, the species gets one point. In addition, a score from 1 to 3 is given if the species is mentioned in the Jensen (2013) List 2 of not yet established ("alert") species depending on the score given them. The Jensen score is derived from an appraisal of the potential for being invasive. Furthermore, if present on EU's regulation list (Berggren *pers. comm.*), it gets the score 3.

- **Criterion 3 – Invasiveness:** The third criterion describes the degree of invasiveness a species is assessed to have. The scores 2, 4, 6 can be obtained, with 6 being the most invasive. The scores are based on a literature study, on the species effect on: 1) colonization of conservational valuable habitats, 2) impact on native species, 3) effect on ecosystem functioning and 4) economical as well as public health effect (Madsen *et al.* 2014). Furthermore, if literature was not available, expert judgements was applied.
- **Criterion 4 – Ease of determination:** The score given under this criterion is based on knowledge of the degree of difficulty in the characterization of the species. If the species is easy to confuse with similar looking species, it received the score of 1, whereas, if it is very characteristic or peculiar and hence, easy to identify, it received the score of 3.
- **Criterion 5 – Ease of determination by environmental DNA (eDNA):** Different species exhibit different levels of complexity in regard to determination with eDNA-based methods (see Díaz-Ferguson *et al.* 2014, Goldberg *et al.* 2015, Taberlet *et al.* 2012, Thomsen & Willerslev 2015, Thomsen *et al.* 2012, Wilcox *et al.* 2013). Species with many sympatric species generally more difficult to determine with eDNA methods than species with no or few close relatives. The ease of determination by eDNA also depends on the availability of DNA reference sequences in public data facilities (e.g. National Center for Biotechnical Information (NCBI), The Barcoding Of Life Data System (BOLD)). The expert judgement of the ease of determination is the basis for the scoring of values between 1 and 3.

The above criteria and the ranking process was planned early in th MONIS 2 process (see Appendix A). Each species is evaluated and scored in regard to these 5 criteria. Finally, the scores are summed together. The 51 species getting the highest scores are those included in the MSFD D2-specific Target Species List. The 37th ranked species, a parasite *Anguillicola crassus*, was excluded from the final list as the chances of detecting a parasite by using eDNA-based methods were anticipated to be low. The total number of species on the final list being passed on to further analysis thus came out at 50.

2.2 Technical Guidance Report for eDNA-based monitoring activities

A robust and scientifically credible Technical Guidance Report is needed to prevent both false positive (e.g. due to cross-contamination or non-specific primers) or false negative (e.g. due to technical error during sampling, filtration, extraction or non-amplifying primers) registration of species presence.

False positives and negatives are of special concern when estimating the potential detrimental effect of non-indigenous species. Large scale eDNA surveys conducted in the USA (Jerde *et al.* 2013) and Japan (Fukumoto *et al.* 2015) serve as good examples of the stringency of Standard Operating Procedures (SOPs) needed when multiple people are involved in sample collection and possibly also laboratory analyses. Thus, for documentation, the SOPs for every step in the Technical Adcisory TA have to be written with the utmost detail for the entire process from sampling to laboratory analysis.

The Techincal Guidance provided for this MONIS phase 2 report considers all 48-50 primer-probe assays systems reviewed as tentative. None of the systems have yet been tested on actual DNA extracts from target-species and sympatric non-target species (i.e. *in vitro* test). Without knowledge of how these systems perform on actual DNA, it is unknown whether any of the suggested systems in fact are able to provide true positive detection of eDNA from the target-species. It should be kept in mind that the *in silico* test only is based on nucleotide sequence comparison. Therefore, preliminary *in vitro* tests for each system that is developed is mandatory to estimate the specificity of each system, before any direct tests can be completed in environmental samples.

2.3 Development of an eDNA-based single-species detections system

An eDNA detection system was based on *in silico* development and testing using a polymerase chain reaction (PCR) detection system that is only based on available DNA-sequence information from target and non-target species.

We have focused on developing a ‘single-species detection system’ for each of the 50 species on the Target Species List. The ‘single-species detection system’ is based on the design of a species-specific primer-probe assay developed for quantitative PCR (qPCR) platforms. This approach will during applications circumvent the need for routinely sequencing of PCR products. In addition, species-specific qPCR assays may potentially enable future quantitative applications of eDNA, i.e. it may be possible to estimate the abundance of eDNA copies from a target species per volume of filtered water. This may enable correlations between the prevalence of eDNA with population assessments or other conventional monitoring data. Also, species-specific qPCR systems are considered more sensitive when targeting eDNA fragments with low abundance and this will reduce the risk of false negatives.

Contrary to the invertebrate and vertebrate species on the Target Species List, phytoplankton and macro algae have until now received relative moderate attention in regards to eDNA or direct species identification from filtrated water samples.

For each of the species on the Target Species List relevant mitochondrial DNA (e.g. CytB, CO1) or nuclear sequences (e.g. 28S) were identified via the online NCBI database (www.ncbi.nlm.nih.gov) and aligned using e.g. the BioEdit software, with the same sequence for closely related species or other species showing sequence comparison. Primers and probes were positioned in sequence sections where the difference between the target and non-target species was the largest. Once positioned, each assay was screened for unknown compatibility in other non-target species by conducting a nucleotide BLAST search via the online NCBI database. If non-related, non-target species were identified during this screening, the primer-probe assay was re-designed to improve the species-specification of the assay. Along with the sequence information, the degree of base pair differentiation for a representative number of non-target species is given. This was done to indicate the likelihood that the individual primers and the probe would not amplify a non-target species.

3. Results

The results in regard to (1) development of a national MSFD D2-specific Target Species List, (2) drafting of a Technical Guidance Report, and (3) *in silico* development of as many species-specific primers as possible, are described in the following sections.

3.1 Target Species List

By using the method described above, a species list of 50 species was deduced (Table 2). In total 21 invertebrates, 12 fish, 12 species of submerged aquatic vegetation, and 5 species of phytoplankton appears on the list. With the exception of 4 species, all of the 50 species, which constitute the Non-indigenous target species list, are established in Denmark.

The MONIS 2 draft Target Species List can be found in Table 2 with supplementary information in Appendix D.

3.2 Technical Guidance Report

A draft Technical Guidance Report has been written in Danish with the tentative title: “Teknisk Anvisning for indsamling af marine vandprøver og analyse for eDNA (*environmental DNA*)” and focuses on the following elements:

1. Sampling and filtration of water
2. Preservation of filters
3. Storage of filters
4. Extraction of DNA
5. Species-specific eDNA detection using qPCR

For details, please confer with the draft text in Appendix B.

3.3 A first generation of MSFD D2-related primers

MONIS, phase 2 reports a total of 48 primer and probe assays out of the 50 species on the Species Target List. Details can be seen in Appendix D.

Table 2: Proposed Target Species List for monitoring of non-indigenous species in Danish marine waters.

No.	Group	Species
1.	Fish	<i>Neogobius melanostomus</i> (Pallas, 1814)
2.	Invertebrate	<i>Crassostrea gigas</i> (Thunberg, 1793)
3.	Invertebrate	<i>Crepidula fornicata</i> (Linnaeus, 1758)
4.	Invertebrate	<i>Teredo navalis</i> Linnaeus, 1758
5.	Phytoplankton	<i>Karenia mikimotoi</i> (Miyake & Kominami ex Oda, 1935) (G. Hansen & Moestrup, 2000)
6.	Fish	<i>Oncorhynchus mykiss</i> (Walbaum, 1792)
7.	Zooplankton	<i>Mnemiopsis leidyi</i> Agassiz, 1865
8.	Invertebrate	<i>Eriocheir sinensis</i> Milne-Edwards, 1853
9.	Invertebrate	<i>Rhithropanopeus harrisi</i> (Gould, 1841)
10.	Phytoplankton	<i>Prorocentrum minimum</i> (Pavillard) J. Schiller, 1933
11.	Subm. Veg.	<i>Gracilaria vermiculophylla</i> (Ohmi) Papenfuss, 1967
12.	Subm. Veg.	<i>Sargassum muticum</i> (Yendo) (Fensholt, 1955)
13.	Invertebrate	<i>Cordylophora caspia</i> (Pallas, 1771)
14.	Invertebrate	<i>Styela clava</i> Herdman, 1881
15.	Invertebrate	<i>Dreissena polymorpha</i> (Pallas, 1771)
16.	Fish	<i>Acipenser baerii</i> Brandt, 1869
17.	Fish	<i>Acipenser gueldenstaedtii</i> Brandt & Ratzeburg, 1833
18.	Fish	<i>Acipenser ruthenus</i> Linnaeus, 1758
19.	Fish	<i>Acipenser stellatus</i> Pallas, 1771
20.	Fish	<i>Huso huso</i> (Linnaeus, 1758)
21.	Fish	<i>Oncorhynchus gorbuscha</i> (Walbaum, 1792)
22.	Fish	<i>Salvelinus fontinalis</i> (Mitchill, 1814)
23.	Subm. Veg.	<i>Colpomenia peregrina</i> (Sauvageau, 1927)
24.	Invertebrate	<i>Ensis americanus</i> (Gould, 1870)
25.	Subm. Veg.	<i>Dasya baillouviana</i> (Gmelin) Montagne, 1841
26.	Subm. Veg.	<i>Heterosiphonia japonica</i> Yendo, 1920
27.	Subm. Veg.	<i>Spartina anglica</i> Hubbard
28.	Phytoplankton	<i>Heterosigma akashiwo</i> (Y.Hada ex Y.Hara & M.Chihara, 1987)
29.	Phytoplankton	<i>Pseudochattonella sarcimen</i> (W. Eikrem, B. Edvardsen & J. Thronsen) W. Eichrem, 2009
30.	Invertebrate	<i>Molgula manhattensis</i> (de Kay, 1843)
31.	Zooplankton	<i>Cercopagis pengoi</i> (Ostroumov, 1891)
32.	Invertebrate	<i>Homarus americanus</i> Milne-Edwards, 1837
33.	Invertebrate	<i>Paralithodes camtschaticus</i> (Tilesius, 1815)
34.	Invertebrate	<i>Didemnum vexillum</i> Romanov, 1989
35.	Subm. Veg.	<i>Fucus evanescens</i> C. Agardh, 1820
36.	Invertebrate	<i>Petricolaria pholadiformis</i> (Lamarck, 1818)
37.	Fish	<i>Percottus glenii</i> Dybowski, 1877
38.	Invertebrate	<i>Elminius modestus</i> Darwin, 1854
39.	Invertebrate	<i>Ficopomatus enigmatus</i> (Fauvel, 1923)
40.	Invertebrate	<i>Marenzelleria viridis</i> (Verrill, 1873)
41.	Invertebrate	<i>Ocenebra inornata</i> (Récluz, 1851)
42.	Invertebrate	<i>Potamopyrgus antipodarum</i> (Gray, 1843)
43.	Phytoplankton	<i>Pseudochattonella verriculosa</i> (Y.Hara & M.Chihara) S.Tanabe-Hosoi, D.Honda, S.Fukaya, Y.Inagaki & Y.Sako, 2007
44.	Subm. Veg.	<i>Codium fragile</i> subsp. <i>fragile</i> (Suringar) Hariot, 1889
45.	Fish	<i>Carassius auratus</i> (Linnaeus, 1758)
46.	Fish	<i>Cyprinus carpio</i> (Linnaeus, 1758)
47.	Invertebrate	<i>Mya arenaria</i> Linnaeus, 1758
48.	Zooplankton	<i>Penilia avirostris</i> Dana, 1849
49.	Invertebrate	<i>Diadumene lineata</i> (Verrill, 1870)
50.	Subm. Veg.	<i>Bonnemaisonia hamifera</i> Hariot, 1891

4. Discussion and recommendations

The methods used, the results and the suggested next steps in regard to (1) development of a national MSFD D2-specific Target Species List, (2) drafting of a Technical Guidance Report, and (3) *in silico* development and testing of as many species-specific primers as possible, are discussed in the following sections.

The proposed Target Species List (Table 2) is the first of its kind in a national context and based on the currently best available information. However, it should be regarded as a “living” document subject to regular updates as new information and regulation become available.

The draft Technical Guidance Report (Appendix B) is also the first of its kind in the context of the NOVANA programme; the descriptions of eDNA-based methods have so far not been included in NOVANA as a standard methodology. The Technical Guidance Report deals with sampling, preservation and laboratory analyses and does in our opinion open the door for sampling with a specific focus on MSFD D2.

Recommendation 1: The Technical Guidance Report should be discussed and tentatively approved for sampling and storage of filters as this would enable the initiation of sampling activities as soon as possible, but not later than summer of 2016.

As for the proposed Target Species List, the Technical Guidance Report should also be considered a “living” document and updated whenever relevant, e.g. based on acquired experience for the use of it or when methods are further developed and tested for operational monitoring and analyses.

Another reason for commencing the sampling activities is that it has been suggested to store all filters in a national Filter Archive as this – at a later stage – would provide a possibility to reanalyse filters focusing on a broader range of species than those treated here. However, for prolonged storage it is recommended to extract DNA from filters, and archive the extracted DNA. If sampling has already been initiated, it is extremely important that similar types of filters are used, and sampling and storage protocols are uniform to ensure samples are comparable across sampling locality and sampling time.

Recommendation 2: The Danish Nature Agency should liaise with a partner capable of running the suggested Filter Archive as this would secure proper handling of filters sampled in 2016 and onwards.

Some institutions that potentially could host the suggested Filter Archive identified by MONIS-2 are: Eurofins Miljø A/S or Aarhus University. The National History Museum in Denmark (NHMD) has officially been appointed as a national cryo-facility, but this is something that will be implemented and operational at a later stage.

All eDNA-based activities should be coordinated and directly linked to the monitoring based on conventional sampling methods, primarily in regard to phytoplankton, zooplankton, submerged aquatic vegetation, fish, seabirds and marine mammals. This coordination should take place as soon as possible and prior to the implementation of eDNA-based sampling.

Recommendation 3: It is strongly recommended to place the stations for sampling of eDNA in connection with stations or areas where conventional MSFD D2-targeted sampling is being ongoing or planned (see MONIS-1 report; Andersen *et al.* 2015). This will enable cross-comparisons and validation.

Following and implementing the above recommendations 1-3 will in our understanding be a critical first step towards an eDNA-based MSFD D2-specific monitoring programme in Denmark. Next steps should

include further development of primers and probe systems and must include *in vitro* testing of target- and non-target species in 2016 and 2017. This will ultimately lead to the development of reporting guidelines.

Recommendation 4: Initiation of a process leading to more and better primer-probe assays is a prerequisite for successful implementation of a nation-wide MSFD D2 monitoring.

In order to strengthen the basis for the proposed activities further and continuous development of primer-probe assays is required, especially for those species on the Target Species List that are not yet considered ‘operational’ (i.e., where species-specific primer-probe assays could not be designed). In addition, the *in silico*-based primer-probe assays in appendix C should be tested and evaluated in order to support prioritization in regard to the further development of primers. Before any of the 50 provided primer and probe systems can be utilized on environmental samples, initial *in vitro* test is needed. *In vitro* test includes tests on extracted DNA from target –and sympatric non-target species which must be performed to confirm the expected specificity of each system. Without *in vitro* test it is not possible to rule out false positive and false negative detection arising from any of the developed systems. *In vitro* testing requires a comprehensive cryo-archive of tissue samples collected from both target- and non-target species. The NHMD currently holds one of Denmark’s largest collections of tissue samples from various organisms, and it is therefore recommended that the mandatory *in vitro* tests are initiated by the NHMD or in very close collaboration with the NHMD.

Finally, we suggest establishing an informal network in regard to the eDNA-based MSFD D2 monitoring activities in Danish marine waters. Key objective for this activity is to support the Danish Nature Agency in implementing the recommendation and result from the MONIS-1 and MONIS-2 projects and to provide science-based support in relation to marine eDNA-based monitoring activities.

Recommendation 5: It is recommended that the Danish Nature Agency establishes a national network or working group which can provide guidance and scientific support with regards to implementation of eDNA-based method in the NOVANA/MSFD-specific marine monitoring activities in Denmark.

In a longer perspective, the suggested Danish activities would need to be coordinated with more or less identical work in progress in neighbouring countries – primarily Sweden and Germany – in order to avoid duplicate effort and the waste of scarce resources.

Recommendation 6: It is strongly recommended that Danish Nature Agency informs neighbouring countries and relevant Regional Marine Conventions about the Danish progress in establishing an eDNA-based monitoring network and in a longer perspective to coordinate the development of specific primers (*inter alia*, e.g. including cost-sharing).

Based on the work carried out, the following conclusions can be drawn:

1. Sampling activities can in principle – pending tentative approval of the MONIS-2 Technical Guidance Report and the initiation of a Filter Archive – be planned and implemented from mid-summer of 2017.
2. A further testing and development of primer-probe assays is required and should be carried out in 2016 and 2017. This testing should include *in vitro* testing of tissue extracted DNA from target and non-target species.
3. A network or working group should be established to provide guidance and scientific support for the Danish Nature Agency and the implementation of eDNA-based methods in the national marine monitoring programmes and activities.

If any of the above conclusions are not being followed up, the forthcoming 2018 MSFD Initial Assessment may lack scientific credibility and be in conflict with pan-European guiding principles for the assessment and reporting of the MSFD.

5. References

Andersen, J.H. & D.J. Conley (2009): Eutrophication in coastal marine ecosystems: towards better understanding and management strategies. *Hydrobiologia* 629(1):1-4.
<http://www.springerlink.com/content/w707717n84j65571/fulltext.pdf>

Andersen, J.H., C.D. Pommer, J.W. Hansen & P. Dolmer (2012): Foreløbig karakterisering af fysisk skader forårsaget af råstofindvinding og bundtrawling i de danske farvande. Fagligt notat fra DCE - Nationalt Center for Miljø og Energi. 27 pp. (In Danish)

Andersen, J.H. & A. Stock (eds.), S. Heinänen, M. Mannerla & M. Vinther (2013): Human uses, pressures and impacts in the eastern North Sea. Aarhus University, DCE – Danish Centre for Environment and Energy. Technical Report from DCE – Danish Centre for Environment and Energy No. 18. 134 pp.

Andersen, J.H., K. Dahl, C. Göke, M. Hartvig, C. Murray, A. Rindorf, H. Skov, M. Vinther & S. Korpinen (2014a): Integrated assessment of marine biodiversity status using a prototype indicator-based assessment tool. *Frontiers in Marine Science*.

<http://journal.frontiersin.org/Journal/10.3389/fmars.2014.00055/abstract>

Andersen, J.H., S.A. Pedersen, J. Thaulow, F. Stuer-Lauridsen & S. Cochrane (2014b): Monitoring of non-indigenous species in Danish marine waters. Background and proposals for a monitoring strategy and a monitoring network. Danish Nature Agency. 55 pp.

Andersen, J.H., J. Carstensen, D.J. Conley, K. Dromph, V. Fleming, B. Gustafsson, A. Josefson, A. Norkko, A. Villnäs & C. Murray (2015): A Baltic Sea-wide analysis of temporal and spatial trends in eutrophication status. *Biological Reviews*. <http://onlinelibrary.wiley.com/doi/10.1111/brv.12221/epdf>

Andersen, J.H., C. Murray, M.M. Larsen, N. Green, T. Högåsen, K. Gustavson, E. Boalt, E. Garnaga, M. Haarich, J. Manio, J. Strand & S. Korpinen (2015): Development and testing of a prototype tool for integrated assessment of chemical status in aquatic environments. *Environmental Monitoring and Assessment*. <http://link.springer.com/article/10.1007/s10661-016-5121-x>

Anon. (2008): Directive 2008/56/EC of the European Parliament and of the Council of 17 June 2008 establishing a framework for community action in the field of marine environmental policy (Marine Strategy Framework Directive). Official Journal of the European Communities L 164/19.

Anon. (2010): Directive 2010/477/EU Commission Decision of 1 September 2010 on criteria and methodological standards on good environmental status of marine waters. Official Journal of the European Communities L 232/14.

Azour, F., van Deurs, M., Behrens, J., Carl, H., Hüssy, K., Greisen, K., Ebert, R., Möller, P.R. 2015. Invasion rate and population characteristics of the round goby *Neogobius melanostomus*: effects of density and invasion history. *Aquat. Biol.*, 24 (2015), pp. 41–52 <http://dx.doi.org/10.3354/ab00634>

Berggren, U. (pers. comm.): Short list of the marine species to the EU regulation concerning non-indigenous species.

Dahllöf, I. & J.H. Andersen (2009): Hazardous and Radioactive Substances in Danish Marine Waters. Status and Temporal Trends. Danish Spatial and Environmental Planning Agency & National Environmental Research Institute. 110 pp.

http://www2.dmu.dk/pub/OSPAR_Hazardous_Substances_print.pdf

Díaz-Ferguson, E.E. & G.R. Moyer GR. (2014): History, applications, methodological issues and perspectives for the use of environmental DNA (eDNA) in marine and freshwater environments. *Rev. Biol. Trop.* 62(4):1273-84.

Froese, R. & D. Pauly. (Eds.) (2015): FishBase. World Wide Web electronic publication. www.fishbase.org, (10/2015). Accessed Nov-2015.

Fukumoto, S., *et al.* (2015). "A basin-scale application of environmental DNA assessment for rare endemic species and closely related exotic species in rivers: a case study of giant salamanders in Japan." *Journal of Applied Ecology* 52(2): 358-365. <http://onlinelibrary.wiley.com/doi/10.1111/1365-2664.12392/full>

Goldberg, C.S., K.M. Strickler & D.S. Pilliod (2015): Moving environmental DNA methods from concept to practice for monitoring aquatic macroorganisms. *Biological Conservation* 183:1-3.

HELCOM (2015): List of Target Species currently in use in the HELCOM area. 1 pp.

Jerde, C. L., *et al.* (2013). "Detection of Asian carp DNA as part of a Great Lakes basin-wide surveillance program." *Canadian Journal of Fisheries and Aquatic Sciences* 70(4): 522-526.
<http://www.nrcresearchpress.com/doi/abs/10.1139/cjfas-2012-0478#Vv5nvvl97RY>

Jensen, K. (2013): Target Species List 1 (established species) and Target Specis List 2 (alert list). 2 pp.

Jensen, K. (pers. comm.): data matrix providing the basis of Jensen, K. 2013.

Korpinen, S., L. Meski, J.H. Andersen & M. Laamanen (2012): Human pressures and their potential impact on the Baltic Sea ecosystem. *Ecological Indicators* 15:105-114.
<http://dx.doi.org/10.1016/j.ecolind.2011.09.023>

Madsen, C.L., C.M. Dahl, K.B. Thirslund, F. Groussset, V.K. Johannsen & H.P. Ravn (2014): Pathways for non-native species in Denmark. Department of Geosciences and Natural Resource Management, University of Copenhagen, Frederiksberg. 131 pp.

Møller, P.R. (2015): Supplerende liste over marine fisk. (In Danish)

Naturstyrelsen (2008a): Sortlisten. Akvatiske arter. 1 pp. (In Danish)

Naturstyrelsen (2008b): Observationslisten. Akvatiske arter. 1 pp. (In Danish)

Naturstyrelsen (2012): Danmarks Havstrategi – Basisanalyse. Miljøministeriet. 100 pp. (in Danish)

OSPAR (2015): OSPAR Target Species List (OSPAR Agreement 2015-10). 2 pp.

Riemann, B., J. Carstensen, K. Dahl, H. Fossing, J.W. Hansen, H.H. Jakobsen, A.B. Josefson, D. Krause-Jensen, S. Markager, P. Stæhr, K. Timmerman, J. Windolf & J.H. Andersen (2015): Recovery of Danish coastal ecosystems after reductions in nutrient loading: trends and time lags. *Estuaries & Coasts*.
<http://link.springer.com/article/10.1007/s12237-015-9980-0>

Stæhr, P.A. & Thomsen, M.S. (2012): Opgørelse over rumlig udbredelse, tidslig udvikling og tæthed af ikke-hjemmehørende arter i danske farvande: Fagligt notat fra DCE - Nationalt Center for Miljø og Energi. 14 pp. (In Danish)

Taberlet, P., E. Coissac, M. Hajibabaei & L.H. Rieseberg (2012): Environmental DNA. *Mol Ecol.* 21(8):1789-93.

Thomsen, P.F. & E. Willerslev (2015): Environmental DNA – An emerging tool in conservation for monitoring past and present biodiversity. *Biological Conservation* 183:4-18.

Thomsen, P.F., J. Kielgast, L.L. Iversen, C. Wiuf, M. Rasmussen, M.T. Gilbert, L. Orlando & E. Willerslev (2012) Monitoring endangered freshwater biodiversity using environmental DNA. *Mol. Ecol.* 21(11):2565-73.

Tranah, G., D.E. Campton, & B. May (2004): Genetic evidence for hybridization of pallid and shovelnose sturgeon. *Journal of Heredity* 95(6):474-480.

Wilcox, T.M., K.S. McKelvey, M.K. Young, S.F. Jane, W.H. Lowe, A.R. Whiteley & M.K. Schwartz (2013): Robust detection of rare species using environmental DNA: the importance of primer specificity. *PLoS One* 8(3):e59520. doi: 10.1371/journal.pone.0059520.

Zhang, X., W. Wu, L. Li, X. Ma & J. Chen (2013): Genetic variation and relationships of seven sturgeon species and ten interspecific hybrids. *Genetics Selection Evolution* 45(21).

Ærtebjerg, G., J.H. Andersen & O.S. Hansen (2003): Nutrients and Eutrophication in Danish Marine Waters. A Challenge to Science and Management. National Environmental Research Institute. 126 pp.

Appendix A: Minutes of the TSL workshop

Dato: Fredag den 18. september 2015

Sted: NIVA Danmark, Ørestand Boulevard 73, 2300 København S

Deltagere:	Jesper H. Andersen Per Andersen Dorte Bekkevold Ulrik Berggren Stine Christiansen Nikolai Friberg Martin Hesselsøe Kathe R. Jensen Steen W. Knudsen Peter B. Mortensen Peter Rask Møller Peter Stæhr Hans Erik Svart Jens Thaulow Mary Wisz	NIVA Danmark Orbicon DTU Aqua Naturstyrelsen Naturstyrelsen NIVA Danmark Amphi Consult Statens Naturhistoriske Museum Statens Naturhistoriske Museum Eurofins Statens Naturhistoriske Museum DCE-AU Naturstyrelsen NIVA Danmark DTU Aqua
------------	---	--

1. Velkomst, introduktion og formål

1.1 Nikolai Friberg og Jesper Andersen indledte med at byde velkommen til workshoppen og til NIVA Danmark. MONIS 1-projektet og baggrunden for MONIS 2-projektet blev kort introduceret.

1.2 Samtlige deltager præsenterede sig via en tour-de-table.

1.3 Det blev understreget, at formålet med workshoppen, som er en del af MONIS 2-projektet, var at drøfte processen og evt. kriterier for udarbejdelse af en dansk MSFD D2 Target Species List, som skal lægges til grund for udviklingen af op til 50 in silico-testede primers.

2. MSFD og overvågning af NIS i danske farvande

2.1 Naturstyrelsen, ved Ulrik Berggren og Hans Erik Svart, gjorde indledningsvis rede for en række forvaltningsmæssige forhold af betydning for den fremtidige nationale overvågning af ikke-hjemmehørende arter i de danske farvande, herunder EU-direktiver og –forordninger, regionale konventioner (HELCOM og OSPAR) og en række nationale tiltag m.v.

2.2 Naturstyrelsen lagde vægt på at udarbejdelsen af en antional MSFD D2 Target Species List tager udgangspunkt i eksisterende lister (internationale og nationale), herunder at der overvejende fokuseres på arter, som allerede er tilstede i de danske farvande eller hvor der er international krav til overvågning, fx. EU-forordningen.

2.3 Naturstyrelsen gjorde det klart at den af MONIS 2 udarbejdede Target Speciel List vil være et første bud på hvilke arter, som skal overvåges samt være et levende dokument som fra tid til anden og efter behov skal opdateres.

2.4 Styrelsen lagde vægt på forventningsafstemning og gjorde i den forbindelse følgende klart: Styrelsen forventer, (1) at en kriterie-baseret dansk MSFD D2 Target Species List bliver udarbejdet som en del af MONIS 2-projektet, (2) at kriterierne tager udgangspunkt i tilstedeværelse, risisci og (3) om hvorvidt de pågældende arter er relevante i en eDNA-sammenhæng. Succeskriteriet er ikke at der nødvendigvis at der bliver udviklet primere for 40-50 arter på listen, men at der blandt de arter, som kommer på, er så mange som muligt der med sikkerhed kan identificeres til art med eDNA.

2.5 Denne vigtige udmelding fra styrelsen blive kort drøftet. Der var blandt MONIS 2-partnerne stort tilfredshed med denne udmelding, specielt styrelsens tilkendegivelse af at succeskriteriet for MONIS 2 er operationalitet frem for et bestemt antalt primere.

3. Amphi Consult og eDNA: Erfaringer fra feskandsovervågningen. Hvor er vi nu og hvor går vi hen?

3.1 Martin Hesselsøe orienterede kort om Amphi Consults arbejde, specielt i relation til eDNA og projekterne vedrørende overvågning af udvalgte arter i ferskvand. Det blev herunder understreget, at MONIS 2 alene kan bidrage til udviklingen af et single species detection system.

4. DTU Aqua og eDNA-baseret overvågning m.v. af fisk

4.1 Dorte Bekkevold orienterede kort om DTU Aquas arbejde med eDNA, specielt i forhold til kommersielle fiskearter og en række af de problemstillinger, der kan knytte sig hertil, bl.a. sæsonvariationer

5. Det nationale Fiskeatlas-projekt og andre relevante SNM-aktiviteter

5.1 Peter Rask Møller orienterede kort om SNM's arbejde med overvågning af kystnære fisk, herunder det nationale Fiskeatlas-projekt m.v. I forlængelse heraf orineterede Sten W. Knudsen kort om SNM's arbejde med eDNA i forhold til udvalgte fiskearter, herunder en række at de udfordringer i forbindelse med metodikken som man fagligt må forholde sig til.

6. Første skridt mod en dansk MSFD D2 Target Species List

6.1 Indledningsvis orienterede Jesper H. Andersen om den i MONIS 2 planlagte proces i for-hold til hvilke arter som evt. skal med på en dansk MSFD D2 Target Species List.

6.2 Fyto- og zooplankton: Diskussionen blev fokuseret på disse to organismegrupper. Efter en del snak frem og tilbage blev det tilkendegivet, at (1) der endnu ikke er mange erfaringer med eDNA-baseret overvågning af disse organismegrupper, og at (2) særlig opmærksomhed som udgangspunkt bør rettes mod følgende arter: (i) *Alexandrium* sp., (ii) *Pseudochattonella* sp. (iii) *Karenia* sp., (iv) 'dræbergoplen' (*Mnenopsis leydi*) og (v) kroghalet rovdafnie (*Cervopagis pengoi*).

6.3 Undervandsvegetation: Efter en kort diskussion blev det tilkendegivet, at (1) der er meget få erfaringer med eDNA-baseret overvågning af marin undervandsvegetation, og at (2) særlig opmærksomhed som udgangspunkt bør rettes mod følgende arter: (i) Gracilariatang (*Gracilaria* sp.) (ii) Saragassotang (*Saragassum muticum*), og (iii) rødalgen *Pophyra umbilicalis* sp..

6.4 Benthiske Invertebratebrater m.v.: Efter en kort diskussion blev det tilkendegivet, at (1) der er nogle erfaringer med eDNA-baseret overvågning af marine Invertebratebrater, bl.a. i relation til Vandrammedirektivet, og at (2) særlig opmærksomhed som udgangspunkt bør rettes mod følgende arter:

(i) Stillehavssøsters (*Crassostrea gigas*), (ii) svovlorm (*Marenzelleria* sp.), (iii) søpunge (*Didemnum* sp). og (iv) Kinesisk Uldhåndskrabbe (*Eriocheir sinensis*).

6.5 Fisk: Efter en del snak frem og tilbage blev det tilkendegivet, at (1) der allerede er en del positive erfaringer med eDNA-baseret overvågning af forskellige fisk, og at (2) særlig opmærksomhed som udgangspunkt bør rettes mod følgende arter: (i) sortmundet kutling (*Neogobius melanostomus*), (ii) pukzellaks (*Oncorhynchus gorbuscha*), og (iii) en række arter af stor.

6.6 Kriterier: Forslag til mulige kriterier for udabridelse af en dansk MSFD D2 Target Species List blev kort drøftet. Der var bred tilslutning til at listen skulle være transparent og kriteriebaseret.

7. Next steps – en åben diskussion

7.1 På baggrund at workshoppen vil der blive udarbejdet et kriterie-baseret udkast til en dansk MSFD D2 Target Species List. For så mange af arterne på denne liste som muligt vil der jf. referatets punkt 2.4 og 2.5 blive udviklet primers, som kan indgå i et Single Species Detection System.

7.2 Hvordan MONIS 2 skal afrapporteres er endnu ikke aftalt. NIVA Danmark vil derfor, i dialog med Naturstyrelsen, udarbejde et format herfor. I den forbindelse tilkendegav styrelsen, at afrapportering nok bør være på engelsk af hensyn til samarbejdet i regi af HELCOM og OSPAR.

8. Opsamling

8.1 NIVA Danmark vil udarbejde et referat og et udkast til en MSFD D2 Target Species List – begge vil blive sendt til samtlige deltagere i dagens workshop. Endelig takkede NIVA Danmark alle for aktiv deltagelse i workshoppen og ønskede god weekend

Appendix B: Draft Technical Guidance Report

1



Titel: Teknisk anvisning for indsamling af marine vandprøver og analyse for 'environmental DNA' (eDNA)

Dokumenttype: Teknisk anvisning	TA. nr.: X01	Version: 1	Oprettet: 01.11.2015
Forfattere:	Gyldig fra: 01.05.2016		
- Steen W.Knudsen, Amphi Consult ApS/Afdeling for Evolutionary Genomics, Statens Naturhistoriske Museum, Københavns Universitet	Sider: 16		
- Martin Hesselsøe, Amphi Consult Aps	Sidst ændret: 0x.0x.201x		
- Peter Rask Møller, Afdeling for Evolutionary Genomics, Statens Naturhistoriske Museum, Københavns Universitet			
- Jesper Andersen, NIVA Danmark			
TA henvisninger	V07 – N12 – A02		

0 Indhold

1 Indledning	2
2 Metode.....	2
Del I. Indsamling af vand	3
Del II. Ekstraktion af eDNA fra det indsamlede vand	4
Del III. In vivo detektering af eDNA ved kvantitativ PCR.....	4
2.1 Tid, sted og periode.....	6
2.2 Udstyr	6
2.3 Procedure.....	7
2.4 Tjekliste	11
2.5 Vedligehold af instrumenter.....	11
2.6 Særlige forholdsregler - faldgruber	12
3 Databehandling	12
3.1 Beregninger.....	12
3.2 Data og koder.....	13
4 Kvalitetssikring	14
4.1 Kvalitetssikring af metode	14
4.2 Kvalitetssikring af data og dataaflevering	14
5 Referencer	14
6 Oversigt over versionsændringer	17

2



1 Indledning

Denne tekniske anvisning er udarbejdet for indsamling af marine vandprøver med henblik på detektering af miljø-DNA ('environmental DNA' eller 'eDNA'), som beskrevet i flere aktuelle studier (f.eks. Ficetola et al., 2008; Goldberg et al., 2011; Olson et al., 2012; Thomsen et al., 2012a,b; Takahara et al., 2012; Pilliod et al., 2013; Taberlet et al., 2012, 2013; Deiner et al., 2015; Sigsgaard et al., 2015).

Den tekniske anvisning forudsætter for selve eDNA-analysen at der forudgående er designet artsspecifikke primere og prøber – gennem *in silico* design (dvs. design gennem analyse af nukleotidsekvens data) – for den pågældende art, der ønskes detekteret og at eDNA i den indsamlede vandprøve forekommer i tilstrækkeligt høje koncentrationer til at kunne detekteres. Anvisningen er under stadig udvikling ved afdelingen for Evolutionary Genomics ved Statens Naturhistoriske Museum, Københavns Universitet, og bør derfor betragtes som foreløbig. Det skyldes bl.a. at det ikke er endeligt fastlagt uafklaret hvorledes eDNA bedst kan indsamles, opbevares og detekteres, da det i høj grad afhænger af både vandkvalitet, indsamling og opbevaring af vandprøve, ekstraktion af DNA og brugerens erfaring med både taxonomi og bioinformatik (jf. Deiner et al., 2015; Renshaw et al., 2015; Thomsen og Willerslev, 2015; Taberlet et al., 2012; Sigsgaard et al., 2015). Erfaringer fra de først års overvågnign vil således medvirke til en løbende udvikling af anvisningen.

Denne anvisning bør som udgangspunkt kun følges såfremt indledende valideringstest (også kaldet *in vitro* test) er blevet udført for de benyttede primer- og probe systemer på DNA ekstraheret fra væv fra både den eftersøgte art, samt på DNA ekstraheret fra væv fra tætbeslægtede arter. Indledende *in vitro* test er påkrævet for ikke at risikere at fejlagtige konklusioner baseres på grundlag af falske positive signaler fra analysen.

2 Metode

Fremgangsmåden er delt i tre: (Del I) indsamling af vand, (Del II) ekstraktion af eDNA fra det indsamlede vand og (Del III) detektering af artsspecifik eDNA vha. kvantitative PCR (qPCR). Bemærk at mange studier benytter forskellige metoder for indsamling og ekstraktion af eDNA (jf. Ficetola et al., 2008; Takahara et al., 2012, 2013; Tréguier et al., 2014; Deiner et al., 2015). For at sikre at resultater er sammenlignelige på tværs af prøver, indsamling og arter, er det vigtigt at fremgangsmåden bliver standardiseret fra start.

Indsamling af vand (Del I) og ekstraktion af eDNA fra det indsamlede vand (Del II) kan som udgangspunkt gennemføres allerede nu. Detektering

3



artsspecifik eDNA vha. qPCR (Del III) forudsætter at både forudgående *in silico* design og grundig *in vitro* test er udført.

Del I. Indsamling af vand

Udbryttet af eDNA er umiddelbart koblet med mængden af indsamlet vand – jo mere vand, jo mere eDNA (Taberlet et al., 2012; Goldberg et al., 2015). Ved filtrering af vand anbefales det derfor, at så meget vand som muligt filtreres igennem en filterenhed. Alkoholfældning er et alternativ til filtrering – se bl. a. studier af Ficotela et al. (2008), Tréguier et al. (2014), Biggs et al. (2015), Deiner et al. (2015) for nærmere beskrivelse af denne fremgangsmåde. Eftersom både Amphi Consult og Statens Naturhistoriske Museum har bedst erfaring med filtrering, bygger denne anvisning udelukkende på filtrering, men det står brugeren frit for om alkoholfældning skal benyttes som et alternativ eller supplement til filtrering.

Den indsamlede filtrerede vandprøve skal p.t. opbevares i månedsvis eller årevis og skal derfor opbevares ved -80° C.

Filterenheden skal være steril inert og have en effektiv porestørrelse på 0.2–0.5 µm (se også Goldberg et al. 2011; Thomsen et al., 2012b). Filtrert placeres i en integreret filterholder, som tillader læk-tæt samling med tryksat filtreringsenhed (f.eks. Amphiltrator, se www.amphi-consult.dk) ved et effektivt overtryk på minimum 1 bar atm (101 kPa), men gerne op imod 3 bar (304 kPa). Indsamles vandprøven fra en større beholder, er det nødvendigt at fjerne potentiel kontaminerende nukleinsyrer fra evt. tidlige vandprøvetagning. Nukleinsyrer (dvs. DNA) kan nedbrydes ved brug oxiderende reagenser ved vask med fortyndet brintoverflit eller fortyndet blegemiddel. Før vandprøven indsamles med filterenheden er det nødvendigt at fjerne overskydende oxiderende reagenser gennem gentagne vaske med havvand fra den pågældende prøveindsamlingslokalitet. Dette sikrer at filteret kun indsamler eDNA fra den pågældende indsamlingslokalitet. Amphi Consult er leveringsdygtig i udstyr der tillader tryksat filtrering med brug af engangsplastic der minimerer risikoen for kryds-kontamination uden brug af oxiderende reagenser eller andre kemikalier (se mere på www.amphi-consult.dk).

For at sikre validering af resultatet er det nødvendigt med flere replikater. Som minimum skal der indsamles tre filterprøve-replikater per lokalitet, men flere replikater kan med fordel indsamles. Vandindsamling, filtertype og præservering af filter skal standardiseres fra start for at sikre sammenlignelighed på tværs af både prøvetagningstidspunkt samt prøvetagningslokalitet.

4



Del II. Ekstraktion af eDNA fra det indsamlede vand

Ekstraktion af eDNA fra filtrerede vandprøver er baseret på DNeasy Blood & tissue kit (Qiagen), eftersom mange eDNA studier netop benytter dette ekstraktionskit (f.eks. Thomsen et al., 2012a,b; Takahara et al., 2013; Biggs et al., 2014; Tréguier et al., 2014; Sigsgaard et al., 2015). Amphi Consult og afdelingen for Evolutionary Genomics ved Københavns Universitet har sammenholdt DNeasy Blood & tissue kit (Qiagen) med NucleoSpin Soil (Macherey-Nagel). umiddelbart var DNeasy Blood & tissue kit (Qiagen) en anelse bedre til at opnense eDNA end NucleoSpin Soil (Macherey-Nagel) for marine vandprøver. Men dette resultat er stadig under analyse. Deiner et al. (2015) fandt også DNeasy Blood & tissue kit (Qiagen) gav bedst resultat.

Ekstraktionsprotokollen herunder kan også benyttes for alkoholfældede vandprøver (se også Ficotela et al. (2008) og Thomsen et al. (2012a) for detaljer).

Del III. In vivo detektering af eDNA ved kvantitativ PCR

Før kvantitativ detektion af eDNA kan indledes (*in vivo* test), er det strengt nødvendigt at grundige *in silico* og *in vitro* test er blevet udført og har sikret at primer- probe systemet ikke giver falske positive signaler pga. uønsket amplifikation af eDNA fra tætbeslægtede arter. Detaljeret *in silico* design kan ikke udelukke uspecifik uønsket amplifikation. Kun en detaljeret og grundig *in vitro* test kan sikre at fejlagtige konklusioner ikke drages på grundlag af falsk positiv amplifikation. Se også studierne af Taberlet et al. (2012), Wilcox et al. (2013) og Thomsen & Willerslev (2015).

Eftersom detektion af eDNA er baseret på et højt antal amplifikationscykler (>40 cykler) med 'annealing' og 'extension', kan primer og probe systemet detektere meget lave koncentrationer eDNA. Men ved >40 amplifikationscykler er der stor risiko for at få uønsket amplifikation fra tæt-beslægtede arter, der forekommer i samme habitat som den eftersøgte art (Wilcox et al., 2013). Vær opmærksom på at det ikke nødvendigvis er muligt at udlede fra et falsk positivt resultat fra qPCR, hvilken anden tætbeslægtet art, der har resulteret i uønsket amplifikation. Ønsket amplifikation kan desværre ikke skelnes fra uønsket amplifikation. Ydermere, skal brugeren være opmærksom på om *in silico* designet af primer- og probe systemet er baseret på ufuldstændig sammenholdning af nukleotid-sekvens data fra 'the National Center for Biotechnology Information (NCBI) GenBank. Arter der ikke var inkluderet i det indledende *in silico* design, kan heller ikke udelukkes som grundlag for mulig falsk positiv amplifikation, se også studiet af Wilcox et al. (2013).

En standard fortyndingsrække kan forberedes, hvis en kvantificering af eDNA i ekstraktionen fra filtervandprøven er ønsket (Ellison et al., 2006;

5



Bustin et al., 2009; Turner et al., 2015). En standard fortyndingsrække kan tilberedes ved en indledende PCR på DNA ekstraheret fra den eftersøgte organisme og med de artsspecifikke primere der senere hen skal bruges for det efterfølgende qPCR setup. PCR-produktet kan derpå oprenses og koncentrationen på det oprensede PCR-produkt kan estimeres vha. spektrofotometri, Nanodrop, Qubit e.l. Eftersom nukleotid-sekvensen på PCR-produktet er kendt, kan molekylevægten beregnes, og antallet af PCR-produkter bestemmes som antal kopier per volumen. Dette oprensede PCR-produkt kan derpå fortyndes til en standard fortyndingsrække indledningsvis med 10^8 kopier per μL og i ti-fold fortyndinger (dvs. 10^7 kopier/ μL , 10^6 kopier/ μL osv. ned til 1 kopi/ μL).

Den findes mange forskellige typer preproducerede mastermix for real-time PCR. Alle qPCR set ups udført ved afdelingen for Evolutionary Genomics ved Statens Naturhistoriske Museum over perioden 2012-2016 er baseret på TaqMan Environmental Master Mix 2.0 (Life Technologies), som også er foretrukket i mange studier på eDNA (Thomsen et al., 2012a,b; Tréguier et al., 2014; Sigsgaard et al., 2015; Turner et al., 2015). Derfor er denne protokol også baseret på TaqMan Environmental Master Mix 2.0 (Life Technologies).

Krydkontamination mellem forskellige qPCR kan forebygges ved brug af dUTP i alle qPCR produkter og addering af uracil DNA glycosylase (UDGase) til qPCR mix, med et indledende opvarmningstrin i qPCR programmet der inaktiverer UDGase (se Sigma Aldrich hjemmeside for yderligere detaljer om uracil DNA glycosylase). TaqMan Environmental Master Mix 2.0 (Life Technologies) indeholder ikke UDGase, og må derfor tilsettes separat. Tilsættes UDGase skal mængden af ddH₂O per reaktion justeres tilsvarende så total volumen forbliver 25 μL per reaktion.

Reaktioner til qPCR forberedes i volumen mellem 25 μL til 100 μL . Her er 25 μL total volumen reaktioner anbefalet. Reaktionerne kan køres på en 'Stratagene Mx3005P quantitative PCR' maskine, eller på en tilsvarende 'real-time' qPCR maskine, der er i stand til at detektere de fluorescerende prober, der benyttes – dvs. maskinen skal kunne detektere FAM og ROX farve.

Skal forekomsten af eDNA vurderes med 95% konfidens-niveau, vil det være påkrævet at analysere minimum 19 positive qPCR-replikat-prøver ud af 20 qPCR-replikater-prøver per filtreret vandprøve, da $19/20 * 100\%$ er lig med 95%. Dette kan hurtigt resultere i uforholdsvis mæssigt mange replikat-prøver, hvorfor en 'presence/absence' detektering uden mulighed for fastsættelse af konfidens-niveau umiddelbart anbefales. Se også studiet af Takahara et al. (2013) for flere detaljer om eDNA-detektering af akvatisk invasive arter. Ønskes blot 'presence/absence' detektering, skal

6



minimum fire qPCR-replikater analyseres per filter-vandprøve, men det anbefales kraftigt at analysere flere end blot fire qPCR-replikat-prøver per filtreret vandprøve.

2.1 Tid, sted og periode

For kystvande skal vandprøver indsamles fra midten af den opblandede vandsøje, hvor fordelingen af eDNA antages som værende ensartet. Undgå vandindsamling nær bunden eller i sedimentet (Treguier et al., 2014; Turner et al., 2015). For lagdelte åbne farvande skal der indsamles vandprøver fra både over og under springlaget. For både kystvande og åbne farvande skal indsamlingen skal foregå på de stationer, der indgår i overvågningsprogrammet. Indsamlingerne skal ske to gange årligt, hhv. sent forår/tidlig sommer og sen sommer/tidlig efterår.

2.2 Udstyr

Følgende udstyr skal bruges i forbindelse med prøvetagningen:

- Nukleinsyrefri vandhenter. F.eks. en spand først overskyldet med 5% hydrogen peroxide, og derpå overskyldet med 96% ethanol, og til sidst skyldet med vand fra den lokalitet, der skal samles ind fra.
- Sterile inerte filterenheder med effektiv porestørrelse på 0,2-0,5 µm
- Filterholder, som tillader læk-tæt samling med tryksat filtreringshæd (f.eks. Amphiltrator - se www.amphi-consult.dk)
- Præserverende buffer f.eks. alkohol (96%), eller EDTA, eller LongMire Buffer (Longmire et al. 1997)
- Termometer. Skal rengøres og renses for nukleinsyrer ligesom vandhenter skal mellem hver prøvetagning.

Følgende udstyr skal anvendes til ekstraktionen:

- DNeasy Blood & tissue kit (Qiagen)
- ATL-buffer (komponent i Qiagen DNeasy Blood & tissue kit)
- Protein kinase K (komponent i Qiagen DNeasy Blood & tissue kit)

Følgende udstyr skal anvendes til selve in vivo detektering af eDNA ved kvantitativ PCR

- Rotor eller mixer, der kan modstå opvarmning til mellem 50°-80° C i varmeskab.
- Varmeskab for inkubation ved temperaturer mellem 20°-80° C.
- Pipetter, i størrelser: P10, P20, P200 og P1000
- Sterile pipettespidser med filterbarriere. Spidserne skal matche størrelsesklasserne på pipetterne



- Spektrofotometer, Nanodrop, Qubit e.l. for mål af koncentration af DNA
- TaqMan Environmental Master Mix 2.0 (Life Technologies)
- dUTP (komponent i TaqMan Environmental Master Mix 2.0)
- Uracil DNA glycosylase (UDGasa)
- Stratagene Mx3005P quantitative PCR eller tilsvarende "real-time" qPCR-maskine, der kan detektere FAM og ROX-farve
- Bord-centrifuge, eller mini-centrifuge med maksimal hastighed på minimum 2000 rpm. Med plads til eppendorf-rør i 1,5-2,0 mL volumen
- Stor centrifuge med maksimal hastighed på minimum 14000 rpm. Med plads til eppendorf-rør i 1,5-2,0 mL volumen
- MX pro software for stratagene Mx3005P qPCR-maskine
- Plastrør for qPCR med plads til 100 µL reaktioner. Plastrør skal være i strips af otte.
- Klare plastlåg i strips af otte til plastrør. Plastlågen skal være flade, og tilpasset fluorescens detektering i qPCR.
- Dobbelts distilleret RNase og DNase frit vand
- Sterile, RNase og DNase fri eppendorf rør i volumen 1,5-2,0 mL
- Primere og prober som angivet for de pågældende systemer

2.3 Procedure

I de følgende tre afsnit gennemgås proceduren for indsamling og filtrering af vand, ekstraktion af eDNA fra filteret vandprøve, samt detektering af eDNA via en qPCR maskine og bearbejdning og analyse af detekteret eDNA data. Bemærk at analyse af data forudsætter kendskab til både metoder i kvantitativ PCR, samt begrænsninger for detektering af eDNA som blandt andet er beskrevet af Wilcox et al. (2013).

Fremgangsmåde for indsamling af vand:

Vand indhentes fra midten af vandsøjlen vha. en nukleinsyrefri vandhenter eller en nukleinsyrefri spand. En pumpe kan også bruges, men skal i såfald kunne renses for rester af nukleinsyrer fra tidligere indsamlingslokaliteter så krydskontamination mellem indsamlingslokaliteter udelukkes. Amphi Consult kan vejlede i indsamling af vand, og tiltag for minimering af krydskontamination (se mere på www.amphi-consult.dk).

Indsamlet vand filtreres derpå gennem en steril nukleinsyrefri filtreringsenhed.

- 1) Benyt et steril filter med en effektiv porestørrelse på 0,2-0,5 µm som tillader filtration ved overtryk og stort filtreringsvolumen. Filtrer mellem 0,5 L og 1,5 L (Thomsen et al., 2012b). Mere end 1,5 L vand kan også filtreres så længe filteret ikke tilstoppes. Følg instruktion-



erne for den pågældende filterenhed. Man kan med fordel benytte udstyr, der integrerer både overtryk og filterenhed for at optimere filtrering i felten og laboratoriet (f.eks. en 'Amphiltrator', se mere på www.amphi-consult.dk).

- 2) Vandprøven indsamlas fra midten af vandsøjlen, hvor fordelingen af eDNA antages som værende ensartet. Undgå vandindsamling nær bunden eller i sedimentet (Treguier et al., 2014; Turner et al., 2015).
- 3) Notér det filtrerede vandvolumen. Notér temperatur på vandet, samt dato for indsamling, og hvem der har indsamlet vandprøven.
- 4) Tøm filterenheden for evt. rest af vandprøve, så filterenheden er så tør som mulig inden filterenheden opbevares på køl og/eller med præserveringsbuffer (se Renshaw et al., 2015 for anbefalinger af typer af præserveringsbuffer).
- 5) Mærk filterprøven med unik nummerkode, for senere korrekt identifikation. Notér dato, indsamlingslokalitet, type af præseverende buffer der blev tilført filteret, opbevaringstemperatur og hvem der har indsamlet prøven.
- 6) Hold filterenheden kold (helst under -15°C, og som minimum under 0°C), indtil ekstraktion af DNA kan påbegyndes.
- 7) Overvej indsamling af flere replikater.

Fremgangsmåde for ekstraktion fra filtrerede vandprøver ved brug af DNeasy Blood & tissue kit (Qiagen):

- 8) For filtre opbevaret i buffer, må filteret først fjernes fra buffer-væsken. Filteret skal udtørres under sterile forhold, evt. i et 'negative flow-hood' i 1-2 timer. Ubehandlede filter kan bruges direkte i næste trin.
- 9) ATL buffer og proteinase K tilsættes et rør med det udtørrede filter, som anbefalet i protokollen for DNeasy Blood & tissue kit (Qiagen). Mængden af ATL og proteinase K kan evt. justeres jf. med Thomsen et al. (2012b). Filter med ATL og proteinase K inkuberes på rotor eller mixer i varmeskab ved 56°C for minimum 2 timer så filtratet lyseres fuldstændigt.
- 10) Væsken med lyseret filtrat kan så overføres med pipette til 'spin columns' i DNeasy Blood & tissue kit (Qiagen), og oprenses ved at følge protokollen for DNeasy Blood & tissue kit (Qiagen).
- 11) For at sikre relativt høje koncentrationer af eDNA bør den sidste eluering ske i maksimalt 200 µL elueringsbuffer.
- 12) For langvarig opbevaring bør koncentrationen på det oprensedte eDNA måles vha. spektrofotometri, Nanodrop, Qubit e.l.
- 13) Ekstraheret eDNA opbevares koldt (under -15° C) til alle tider, bortset fra når det skal bruges i qPCR set ups.

9



Fremgangsmåde for detektering af artsspecifik eDNA vha. qPCR:

- 14) Som minimum skal fire replikater per prøve for hver ekstraheret filtreret vandprøve analyseres for en 'presence/absence' detektering uden mulighed for fastsættelse af konfidens-niveau. Bemærk at det kræver minimum 19 positive replikater ud af 20 replikater for at sikre 95% konfidens-niveau for tilstedeværelse af det eftersøgte eDNA.
- 15) Individuelle reaktioner klargøres til 25 µL total volumen, der består af 10 µL of TaqMan, Environmental Master Mix 2.0 (Life Technologies), 9 µL ddH₂O, 3 µL ekstraheret DNA, 1 µL af hver primer (10 µM) og 1 µL probe (2.5 µM). Mere end 3µL ekstraheret DNA kan med fordel tilsvættes for forøget chance for eDNA detektion, i så fald skal mængden af ddH₂O nedjusteres tilsvarende per reaktion. F.eks. hvis der i stedet benyttes 5µL ekstraheret DNA, skal der kun tilføres 7 µL of ddH₂O per reaktion. Som ved ordinær PCR set up, bør alle reagenser mixes og spinnes efter optøning (omkring 5-10 s ved 1000 rpm) i en lille bord-centrifuge, for at sikre at alle reagenser der tilføres er i de tiltænkte koncentrationer.
- 16) Minimum fire reaktioner skal tilsidesættes for 'non-target control' (NTC) test, for at sikre at alle reagenser er fri for kontamination.
- 17) 'Positive target control' (PTC) skal inkluderes i to til fire reaktioner. Inkluderes en standard fortyndningsrække kan den samtidig tjene som PTC. Ellers kan en PTC forberedes ved at til sætte en fortyndning af DNA ekstraheret fra væv fra den eftersøgte organisme.
- 18) En standard fortyndningsrække forberedes i ti-fold fortyndinger - dvs. i 10⁷ kopier/µL, 10⁶ kopier/µL, 10⁵ kopier/µL, osv. ned til 100 kopier/µL. Hver ti-fold fortynding skal testes i minimum tre replikater. Jf. også med manualen til MxPro softwaren for Stratagene Mx3005P qPCR maskinen for fremstilling af standard fortyndningsrækker. Bemærk at en standard fortyndningsrække betragtes som unik per qPCR set up, og ikke kan overføres til lignende eller andre qPCR set ups. Se også anbefalinger fremført af Bustin et al. (2009) og Turner et al. (2015).
- 19) Programmet for qPCR skal være: 5 minutter ved 50° C, 10 minutter ved 95° C, og derpå 50 cykler med 95° C i 30 s og 60° C for 1 minut. Med 'end-point collection' af fluorescerens data under 60° C trinnet.
- 20) Da proberne er forsynet med BHQ1 og FAM-farve og TaqMan, Environmental Master Mix 2.0 (Life Technologies) indeholder ROX-farve sættes qPCR maskinen til at detektere FAM med ROX som reference.
- 21) Resultatet fra qPCR kørslen kvantificeres ved brug standard kurven udregnet fra standard fortyndningsrækken. Eftersom det oprindeligt

10



filtrerede vandvolumen er kendt, og volumen af tilsat elueringsbuff-
er er kendt, og volumen ekstraheret eDNA i den efterfølgende qPCR
er kendt, kan mængden af eftersøgt eDNA i den oprindelige
vandprøve beregnes, og i sidste ende benyttes som et vejledende
mål for den relative forekomst af den eftersøgte organisme på den
pågældende lokalitet. At benytte forekomsten af eDNA i den fil-
trerede vandprøve som mål for forekomsten af den eftersøgte or-
ganisme, vil kræve at der er indsamlet bestandsstørrelsesvurderin-
ger for den eftersøgte art med konventionelle metoder samtidig med
filtervandprøve er blevet indsamlet, og at disse samtidige indsam-
linger af filter og bestandsstørrelsesvurderinger er blevet gjort
løbende igennem en længere årrække. Idet organismer må for-
ventes at udskille eDNA med forskellig rate (se også Thomsen et al.,
2012a,b), og få deres eDNA nedbrudt med forskellig rate med varia-
tion indenfor både arter, populationer, indsamlingstidspunkt og
indsamlingslokalitet, er det ikke muligt at bestemme bestandsstør-
relsen ud fra eDNA-mængden.

- 22) For kvantificering af kopier af det eftersøgte eDNA i den ekstrahe-
rede filtrerede vandprøve er det nødvendigt at skelne mellem 'limit
of Detection' (LOD) og 'level of quantification' (LOQ). For at fastslå
LOD og LOQ, må negative replikater i standard fortyndingsrækken
først fjernes, da de ellers kan påvirke beregning af standardkurven.
Dette kaldes også 'exclusion by sample' jf. studiet af Ellison et al.
(2006). LOD er defineret som den laveste standard med positiv Ct-
værdi (Cycle threshold) (se Bustin et al., 2009), hvor minimum én
ud af flere qPCR replikater returnerer positive amplifikation - dette
er som regel omkring den laveste koncentration af standard fortynd-
ingsrækken, med kun én kopi per reaktion. Prøver med Ct-værdier
over LOD bør betragtes som værende negative.
- 23) LOQ er defineret som laveste reproducerbare standard med positiv
Ct-værdi, dvs. der hvor hver qPCR-replikat for den samme reaktion
resulterer i positiv amplifikation - dette er som regel omkring 5-10
kopier per reaktion. Prøver med Ct-værdier over LOQ kan betragtes
som værende positive (dvs. indenfor LOD), men kun mulige at
kvantificere som mindre end den mindste reproducerbare standard
(f.eks. < 5 eller 10 kopier per reaktion).
- 24) Efter LOD og LOQ er fastsat, kan mængden af eftersøgt eDNA per
replikat i qPCR reaktionerne fra de ekstraherede filtrerede
vandprøver kvantificeres forudsat Ct-værdierne er under LOQ.
- 25) Koncentration af eftersøgt eDNA i den oprindelige filtervandprøve
kan derfor beregnes ved hjælp af følgende formel: $A_e = C_{qpcr} * F_e$
 $* V_{wf}$. Hvor A_e er antallet af eDNA-kopier per volumen filtreret
vand, C_{qpcr} er koncentrationen af kopier af det eftersøgte eDNA i
qPCR-reaktionen, F_e er brøkdelen af det eluerede ekstraherede fil-
trat benyttet i qPCR-reaktionen og V_{wf} er volumen af vand filtreret.

11



2.4 Tjekliste

Ved vandindsamling:

- Vandhenter rengjort forinden
- Udstyr for vandindsamling klargjort
- Minimum 0,5 L vand er filtreret
- Notater omkring vandprøve er optegnet
- Filteret opbevaret i præserveringsbuffer, og hvilken buffer der i såfald er benyttet.
- Slut-filtrat er opbevaret ved < -15° C
- Hver filterprøve er mærket med unik nummerkode

Ved ekstraktion:

- Ekstraktionskit er klargjort forinden ekstraktion påbegyndes
- Yderligere remedier for ekstraktion er klargjort inden ekstraktionerne påbegyndes
- Notater for detaljer om ekstraktionerne er optegnet
- Hver ekstraktion er mærket med unik nummerkode
- Ekstraheret filterprøve er opbevaret ved <-15° C.

Ved qPCR analyse:

- Alle remedier og reagenser er klargjort
- Standardfortyndingrække er klargjort forinden
- Primere og prober er fortyndet til de korrekt 'working solutions' inden qPCR reaktionen sættes op
- Tilfredsstillende in silico test og tilfredsstillende in vitro test er blevet gennemført inden test bliver forsøgt på filtrerede vandprøver. Således at der ikke er risiko for detekttering af falske positive
- Notater er optegnet omkring indstillingen for qPCR opsætningen og for hvilke vandprøver der analyseres.
- qPCR analyse er kørt, og resulterende data-fil gemt under unikt filnavn.
- Ved 'presence/absence' studier er tilstedeværelse/fravær af den eftersøgt organisme noteret, og ved eDNA koncentrations-kvantificerings-forsøg er eDNA koncentrationer målt for hvert replikat i qPCR analysen for hver filterenhed.

2.5 Vedligehold af instrumenter

De fleste komponenter såsom plasticrør, Qiagen ekstraktionkit, filtrer, pipettespidser m.v. er engangsbrug. Benyttes en 'Amphiltrator' (www.amphi-consult.dk) for vandinsamling bør denne vedligeholdes som

12



specificeret i manual og protokol. Pipetter bør vedligeholdes, kalibreres og renses i hht. til vejledninger for disse.

For vedligeholdelse af qPCR maskinen henvises også til manualen og softwaren for dette apparat.

Primere og prober, samt ekstraktioner fra filtere og forberedte fortyndindingsrækker bør opbevares ved temperaturer under -15° C, for at sikre lang holdbarhed.

For brug og opbevaring af TaqMan EM Master Mix 2.0 henvises til vejledningen for dette produkt.

2.6 Særlige forholdsregler - faldgruber

For at kunne reproducere resultater og sammenholde mål for eDNA er det yderst vigtigt at både indsamling, opbevaring, ekstraktion og qPCR kvantificering fastsættes fra start. Afgivelser eller ændringer i protokollen kan influere mængden af eDNA kraftigt, og gøre sammenligninger umulige.

For sikker detektering er det påkrævet at have et stærkt kendskab til litteraturen omkring sporing vha. eDNA, og til taxonomien for den eftersøgte organisme såfremt *in silico* designet ikke er baseret alle tætbeslægtede arter til den eftersøgte art. Derudover forudsættes det også at brugeren har en god forståelse for både det forudgående *in silico* design, og *in vitro* test, og et kendskab til hvordan en real-time PCR maskine fungerer og aflæses korrekt.

Falske positive signaler kan ikke umiddelbart skelnes fra sande positive. Der gives derfor ingen umiddelbar hurtig protokol for sikker eDNA-detektering af arter, men i stedet en række af vejledende retningslinjer der sammen med grundige overvejelser kan sikre korrekt detektering af eDNA.

3 Databehandling

Beskrivelse af dataflow herunder opmærksomhedspunkter.

3.1 Beregninger

Angivelse af hvordan beregninger med de indsamlede data skal foregå. Det drejer sig ikke om, de beregninger, hvor data skal indgå, men fx beregning af indeksværdi til breddevariationen og hvilken dybde blandingsprøverne til søkemi skal tages i.

13



3.2 Data og koder

Liste over de data, der skal indlæses i database og angivelse af hvilken database.

For hvert filter skal følgende detaljer noteres. Forkortelseskoder for disse detaljer er angivet i parentes:

Indsamlingslokalitet, breddegrad og længdegrad (lok_pos):

- Filtertype benyttet (Fil_Typ)
- Temperatur på vandprøve inden filtrering (Temp_Inds)
- Dato for indsamling (Dato_Inds)
- Navn på hvem der har indsamlet prøven (Navn_Inds)
- Dybdestrata for den indsamlede prøve (Dyb_Str)
- Volumen vand filtreret (Vwf)
- Unikt nummer for den pågældende prøve, for at sikre at replikat-prøver kan skelnes fra hinanden (U_Pr_Nr)
- Typen af præserveringsbuffer benyttet for at konservere filterprøven (Pr_Buff)
- Temperatur hvorunder filterprøven efterfølgende er opbevaret (Temp_F_Opb)

For hver ekstraktion af filter skal følgende noteres. Forkortelseskoder for disse detaljer er angivet i parentes:

- Unikt nummer for den pågældende prøve, skal være identisk med nummeret tildelt under filtrering (U_Pr_Nr)
- Dato for ekstraktion (Dato_eks)
- Ekstraktionskit benyttet (Eks_Kit)
- Volumen elueringsbuffer benyttet under ekstraktionen (Vol_EB)
- Temperatur hvorunder det ekstraherede filtrat opbevares (Temp_Ex_Opb)
- Målt koncentration på det ekstraherede eDNA (Cex_eDNA)

For hver kvantitativ PCR setup skal følgende data noteres. Forkortelseskoder for disse detaljer er angivet i parentes:

- Unikt fortløbende nummer tildelt pladen med alle qPCR reaktioner (Q_Nr)
- Filhavn tildelt pladen med qPCR reaktionerne. Et unikt nummer der også skal indeholde det unikke fortløbende nummer tildelt pladen qPCR-reaktionerne (QPI_No).
- Dato for qPCR (Dato_qpcr)
- Volumen på qPCR reaktioner (vol_qpcr_rxn)
- Eftersøgt art (Eft_art)

14



- Primer og probe system benyttet (prim_prob_sys)
- Antallet af reaktioner (Nr_rxn)
- Antallet af NTC reaktioner (NTC_rxn)
- Antallet af replikater for hver filtrat-ekstraktion (Repl_unk_rxn)
- Antallet af replikater for hver standard-fortyndingsrække (Repl_SD_rxn)
- Hvilke ekstraktioner af hvilke filtrater der er forsøgt analyseret i den pågældende qPCR, dvs. hvilke 'U_Pr_Nr' der er testet.
- Volumen af filtrat fra hver 'U_Pr_Nr'-esktraktion der er forsøgt testet per reaktionsrør (Vol_EksFil)
- Hvilken type real-time kvantitativ PCR maskine der er benyttet (Type_qpcr_mask)
- Hvem der har foretaget qPCR analysen (Resp_qPCR)

Eventuelt en liste med koder, der skal anvendes herunder hvilke standatkoder, der skal anvendes.

Hvis denne beskrivelse er den samme i mere end én TA, bliver dette beskrevet i en særskilt "databehandlings-TA". Afsnittet skal altid være i TA'en, men evt. kun med henvisninger til den "databehandlings-TA", der beskriver, hvordan data skal behandles.

4 Kvalitetssikring

4.1 Kvalitetssikring af metode

Beskrivelse af hvordan metoden kvalitetssikres.

4.2 Kvalitetssikring af data og dataaflevering

Beskrivelse af hvordan data kvalitetssikres.

5 Referencer

Bustin SA, Benes V, Garson JA, Hellemans J, Huggett J, Kubista M, Mueller R, Nolan T, Pfaffl MW, Shipley GL, Vandesompele J, Wittwer CT. (2009). The MIQE guidelines: minimum information for publication of quantitative real-time PCR experiments. Clinical Chemistry 55 (4): 611-622.

Deiner, K., Walser, J.-C., Mächler, E., Altermatt, F., 2015. Choice of capture and extraction methods affect detection of freshwater biodiversity from environmental DNA. Biol. Conserv. 183, 53–63.

15



Ellison, S.L.R., English, C.A., Burns, M.J., Keer, J.T., 2006. Routes to improving the reliability of low level DNA analysis using real-time PCR. *BMC Biotechnol.* 6, 33.

Ficetola, G.F., Miaud, C., Pompanon, F., Taberlet, P., 2008. Species detection using environmental DNA from water samples. *Biol. Lett.* 4, 423–425.

Goldberg, C.S., Pilliod, D.S., Arkle, R.S., Waits, L.P., 2011. Molecular detection of vertebrates in stream water: a demonstration using Rocky Mountain Tailed Frogs and Idaho Giant Salamanders. *PLoS ONE* 6, e22746.

Goldberg, C.S., Sepulveda, A., Ray, A., Baumgardt, J., Waits, L.P., 2013. Environmental DNA as a new method for early detection of New Zealand mudsnails (*Potamopyrgus antipodarum*). *Freshwater Sci.* 32, 792–800.

Longmire, J.L., Maltbie, M., Baker, R.J., 1997. Use of 'lysis buffer' in DNA isolation and its implications for museum collections. *Museum of Texas Tech University*, 163, 1–3.

Sigsgaard, E.E., Carl, H., Møller, P.R., Thomsen, P.F. (2015). Monitoring the near-extinct European weather loach in Denmark based on environmental DNA from water samples, *Biological Conservation* 183: 46-52.

Takahara, T., Minamoto, T., Yamanaka, H., Doi, H., Kawabata, Z., 2012. Estimation of fish biomass using environmental DNA. *PLoS ONE* 7, e35868. Takahara, T., Minamoto, T., Doi, H., 2013. Using environmental DNA to estimate the distribution of an invasive fish species in ponds. *PLoS ONE* 8, e56584.

Takahara, T., Minamoto, T., Doi, H., 2014. Effects of sample processing on the detection rate of environmental DNA from the Common Carp (*Cyprinus carpio*). *Biol. Conserv.*

Thomsen, P., Kielgast, J., Iversen, L.L., Wiuf, C., Rasmussen, M., Gilbert, M.T.P., Orlando, L., Willerslev, E., (2012a). Monitoring endangered freshwater biodiversity using environmental DNA. *Mol. Ecol.* 21, 2565–2573.

Thomsen, P.F., Kielgast, J., Iversen, L.L., Møller, P.R., Rasmussen, M. & Willerslev, E. (2012b) Detection of a diverse marine fish fauna using environmental DNA from seawater samples. *PLoS One*, 7, e41732.

Treguier, A., Paillisson, J.-M., Dejean, T., Valentini, A., Schlaepfer, M., Roussel, J.-M., (2014). Environmental DNA surveillance for invertebrate

16



species: advantages and technical limitations to detect invasive crayfish *Procambarus clarkii* in freshwater ponds. J. Appl. Ecol. 51, 871–879.

Wilcox, T.M., McKelvey, K.S., Young, M.K., Jane, S.F., Lowe, W.H., Whiteley, A.R., Schwartz, M.K., 2013. Robust detection of rare species using environmental DNA: the importance of primer specificity. PLoS ONE 8, e59520.

UDKAST

17



6 Oversigt over versionsændringer

Appendix C: *In silico* developed primers

The following pages describe species-specific primers / probe assays for each of the 50 species on the Target Species List.

The information presented includes:

1. Species number (following Table 2)
2. Species name
3. Species-specific primer/probe assay for the species in question with:
 - a. target gene
 - b. product size, melting temperature
 - c. primer/probe length
 - d. GC ratio (%)
4. Number of mismatches between primer and probe region in closely related non-target species

Species 1: Neogobius melanostomus

Full name: *Neogobius melanostomus* Pallas, 1814.

The genus *Neogobius* comprises four species and belongs to the family Gobiidae (Stepien & Nielson 2013; Froese & Pauly 2015).

An assay for *Neogobius melanostomus* has already been published and validated based on the mitochondrial cytochrome oxidase 1 (CO1) region and tested on digital droplet PCR (Nathan *et al.* 2014; Nathan *et al.* 2015). Annealing temperature used for this assay was 60 °C; the annealing temperatures in the table S1.1 below are the theoretical annealing temperature range calculated by Geneious version 9.0.4.

- Neo_Mel_CO1_F01: 5'-CTTCTRGCCCTCCTCTGGWGTG-3'
- Neo_Mel_CO1_R01: 5'-CCCWAGAATTGASGARATKCCGG-3'
- Neo_Mel_CO1_P01: 5'-FAM-FCAGGCAACTTRGCACATGCAG-BHQ-1-3'

Table S1.1: Species-specific primer/probe assay for *Neogobius melanostomus* with target gene, product size, melting temperature, primer/probe length, GC ratio (%), and number of mismatches between primer and probe region in closely related non-target species.

Species	Gene	Product size	Temp	Length	GC
<i>Neogobius melanostomus</i>	CO1	150 bp			
Neo_Mel_CO1_F01	CTTCTRGCCCTCCTCTGGWGTG		59.6 -62.8	22	54 -59
Neo_Mel_CO1_R01	CCCWAGAATTGASGARATKCCGG		58.9 -63.9	23	47-56
Neo_Mel_CO1_P01	CAGGCAACTTRGCACATGCAG		60.1 - 62.9	21	52 -57
Related species	F01	R01	P01		
<i>Neogobius melanostomus</i>	0	0	0		
<i>Neogobius fluviatilis</i>	1	1	0		
<i>Ponticola kessleri</i>	2	0	1		
<i>Ponticola ratan</i>	2	0	0		
<i>Ponticola rhodionii</i>	2	1	2		

Due to both the large degeneracy within the assays and better overall coverage on related species on the mitochondrial cytochrome b region a secondary assay was developed:

- Neo_Mel_Cytb_F01: 5'- TGCTTTAACCTCTCGCCC-3'
- Neo_Mel_Cytb_R01: 5'-GGAGTGACGAGGGGGTTG-3'
- Neo_Mel_Cytb_P01: 5'-FAM-TCGGAGACCCAGACAACCTCATCCCAG-BHQ-1-3'

Beside the BLAST search of the assay components other native Gobiidae species were also included in the alignments to improve and evaluate the assay, see table S1.2 on the next page.

Table S1.2: Species-specific primer/probe assay for *Neogobius melanostomus* with target gene, product size, melting temperature, primer/probe length, GC ratio (%), and number of mismatches between primer and probe region in closely related non-target species.

Species	Gene	Product size	Temp	Length	GC
<i>Neogobius melanostomus</i>	Cytb	84 bp			
Neo_Mel_Cytb_F01		TGCTTTAACCTCTCTCGCCC	59.8	20	55
Neo_Mel_Cytb_R01		GGAGTGACGAGGGGGTTTG	60.0	19	63
Neo_Mel_Cytb_P01		TCGGAGACCCAGACAACITCATCCCAG	67.5	27	56
Related species	F01	R01	P01		
<i>Neogobius fluviatilis</i>	3	1	4		
<i>Neogobius pallasi</i>	4	2	3		
<i>Babka gymnotrachelus</i>	4	3	3		
<i>Ponticola kessleri</i>	6	4	3		
<i>Ponticola ratan</i>	7	4	3		
<i>Ponticola platyrostris</i>	9	4	5		
<i>Ponticola rhodionis</i>	6	3	4		
<i>Gobius niger</i>	7	3	4		
<i>Pomatoschistus minutus</i>	8	4	6		
	4	6	4		

Species 2: Crassostrea gigas

Full name: *Crassostrea gigas* (Thunberg, 1793)

The genus *Crassostrea* comprises around 40 species worldwide (WoRMS Editorial Board 2015, Coan and Valentich-Scott 2012). The National Center for Biotechnology Information (NCBI) GenBank (Nov-2015) has sequences from 23 species of *Crassostrea* represented. The greatest diversity of *Crassostrea* on the NCBI GenBank database is found in sequence fragments of mitochondrial cytochrome oxidase 1 (mtDNA-CO1). The closest related species to *Crassostrea gigas* appears to be *Crassostrea angulata* found off Portugal (Batista *et al.* 2005) and *Crassostrea sikamea* off China (Wang *et al.* 2013). Also, *Crassostrea gigas* and *C. angulata* are known to hybridize (Batista *et al.* 2005).

Other mtDNA regions currently unavailable on NCBI, such as the mtDNA control region, might hold more variation and make it possible to distinguish between *C. gigas* and *C. angulata* in short stretches of mtDNA (80–110 bp long).

As the detection system recommended here is based on the mtDNA-CO1 fragment, this primer-probe system designed for *Crassostrea gigas* (Table S2) is likely to be unable to distinguish between *C. gigas* and *C. angulata* and *C. sikamea*, but should be able to distinguish *C. gigas*, *C. angulata*, *C. sikamea* from any other species of *Crassostrea*.

The following primers and probe are recommended as an assay for detecting eDNA from *Crassostrea gigas*, *C. angulata* and *C. sikamea*:

- Cra_gig_CO1_F05: 5'-AAGCCTTCACCTTGCTGGTA-3',
- Cra_gig_CO1_R05: 5'-CTAGTAAATGGCCCCAACAA-3',
- Cra_gig_CO1_P05: 5'-FAM-GCTCTATTTCAGGTCAATTAAATTCA-BHQ-1-3'

Table S2: Species specific primer/probe assay for *Crassostrea gigas* with, target gene, product size, melting temperature, primer/probe length, GC ratio (%), and number of mismatches between primer and probe region in closely related non-target species.

Species	Gene	Product size	Temp	Length	GC
<i>Crassostrea gigas</i>	CO1	95 bp			
Cra_gig_CO1_F05		AAGCCTTCACCTTGCTGGTA	59.9	20	50
Cra_gig_CO1_R05		CTAGTAAATGGCCCCAACAA	59.8	20	50
Cra_gig_CO1_P05		GCTCTATTTCAGGTCAATTAAATTTC	60.2	26	20

Related species	Forward	Reverse	Probe
<i>Crassostrea angulata</i>	1	0	1
<i>Crassostrea ariakensis</i>	5	6	2
<i>Crassostrea belcheri</i>	5	2	3
<i>Crassostrea brasiliiana</i>	5	4	2
<i>Crassostrea columbiensis</i>	5	8	5
<i>Crassostrea gasar</i>	5	4	2
<i>Crassostrea gigas</i>	0	0	0
<i>Crassostrea gryphoides</i>	7	5	3
<i>Crassostrea hongkongensis</i>	4	2	3
<i>Crassostrea iredalei</i>	5	2	3
<i>Crassostrea madrasensis</i>	5	2	3
<i>Crassostrea nippona</i>	4	4	3
<i>Crassostrea rhizophorae</i>	7	3	3
<i>Crassostrea sikamea</i>	4	3	1
<i>Crassostrea sikamea</i>	4	3	1
<i>Crassostrea virginica</i>	6	4	3
<i>Saccostrea cucullata</i>	7	6	6

Species 3: Crepidula fornicate

Full name: *Crepidula fornicate* Linnaeus, 1758.

The genus *Crepidula* comprises 43 marine species and belongs to the family Calyptidae (WoRMS 2015). The CO1 region is the most suitable region for the assay as it has the best coverage of both target and non-target sister species, with CO1 sequence data available for 25 out of 43 species. The regions 16S, 18S and 28S were also assessed.

- Cre_For_CO1_F01: 5'- TCGGGGGATTGGTAATTGGT -3'
- Cre_For_CO1_R01: 5'-CGGCCGAGGATAGCAATAGT -3'
- Cre_For_CO1_P01: 5'-FAM-AGGTGCTCCTGATATAGCTTTCCCTCGA-BHQ-1-3'

Based on the data available at NCBI the assay is specific, but specificity towards sister-species potentially occurring in Danish territorial waters cannot be excluded.

Table S3: Species-specific primer/probe assay for *Crepidula fornicate* with target gene, product size, melting temperature, primer/probe length, GC ratio (%), and number of mismatches between primer and probe region in closely related non-target species.

Species	Gene	Product size	Temp	Length	GC
<i>Crepidula fornicate</i>	CO1	126			
Cre_For_CO1_F01	TCGGGGGATTGGTAATTGGT		59.4	21	47.6
Cre_For_CO1_R01	CGGCCGAGGATAGCAATAGT		59.4	20	55
Cre_For_CO1_P01	AGGTGCTCCTGATATAGCTTTCCCTCGA		65.1	28	46.4

Related species	F01	R01	P01
<i>Crepidula adunca</i>	4	6	4
<i>Crepidula arenata</i>	5	6	4
<i>Crepidula argentina</i>	7	7	3
<i>Crepidula onyx</i>	8	8	8
<i>Crepidula protea</i>	7	7	3
<i>Crepidula procellana</i>	5	5	3
<i>Crepidula complanata</i>	4	5	4
<i>Crepidula coquimbensis</i>	4	6	2
<i>Crepidula incurva</i>	7	8	6
<i>Crepidula marginalis</i>	5	6	2
<i>Crepidula philippiana</i>	5	5	2
<i>Crepidula cf. Perforans</i>	5	6	2
<i>Crepidula lessoni</i>	4	6	2
<i>Crepidula maculosa</i>	4	8	5
<i>Crepidula naticarum</i>	4	8	1
<i>Crepidula excavata</i>	4	8	4
<i>Crepidula atrasolea</i>	6	8	3
<i>Crepidula depressa</i>	7	7	2
<i>Crepidula plana</i>	5	6	6
<i>Crepidula cf. aplphysioides</i>	4	6	2
<i>Crepidula navicula</i>	5	6	2
<i>Crepidula williamsi</i>	3	5	3
<i>Crepidula convexa</i>	5	8	3
<i>Crepidula unguiformis</i>	5	7	5
<i>Crepidula fimbriata</i>	5	8	6

Species 4: Teredo navalis

Full name: *Teredo navalis* Linnaeus, 1758.

The genus *Teredo* comprises 15 marine species. The genus belongs to the family Teredinidae which contains three subfamilies and includes a total of 25 marine species (WoRMS 2015).

Presently, is *T. navalis* only represented by two sequences covering the 28S and 18S segment in the NCBI database. Only one (*T. clappi*) out of the 14 sister-species within the *Teredo* genus has available sequence data for the same regions. A comparison of the sequences reveals limited differentiation between *T. clappi* and *T. navalis* which potentially could lead to lack of separation between these two species. *Teredo clappi*, however, is only found in the western Atlantic, and will therefore not be a problem for specific detection of *T. navalis* in Danish waters. Available data from several related species in other genera were also included in the alignments to improve the specificity of the assay. The following primers and probe are designed to target *T. navalis*:

- Ter_Nav_28S_F01: 5'-TGGATCCGCTACAACGATCG-3'
- Ter_Nav_28S_R01: 5'-CCGGAGAAAGTGCACCCTG-3'
- Ter_Nav_28S_P01: 5'-FAM-CGCACCGGGAGACGTCTCGTTCTCG-BHQ-1-3'

Table S4: Species-specific primer/probe assay for *Teredo navalis* with target gene, product size, melting temperature, primer/probe length, GC ratio (%), and number of mismatches between primer and probe region in closely related non-target species.

Species	Gene	Product size	Temp	Length	GC
<i>Teredo navalis</i>	28S	124			
Ter_Nav_28S_F01	TGGATCCGCTACAACGATCG		60.0	20	55
Ter_Nav_28S_R01	CCGGAGAAAGTGCACCCTG		60.7	19	63
Ter_Nav_28S_P01	CGCACCGGGAGACGTCTCGTTCTCG		70.6	25	68

Related species	F01	R01	P01
<i>Teredo clappi</i>	0	0	1
<i>Dicyathifer manni</i>	6	1	7
<i>Neoteredo reynei</i>	7	0	7
<i>Teredothyra dominicensis</i>	8	0	5
<i>Lyrodus massa</i>	2	0	1
<i>Lyrodus pedicellatus</i>	2	0	2

Species 5: Karenia mikimotoi

Full name: *Karenia mikimotoi* (Miyake & Kominami ex Oda) (G. Hansen & Ø. Moestrup 2000).

The genus *Karenia* comprises 9 species and an assay system has already been published for the target species within this genus (Yuan *et al.* 2012). The assay was designed for the Internal Transcribed Spacer 18S sequence:

- KMF: 5'-CITTGTTGTAACCTGTTGCTTGT-3'
- KMR: 5'-TCAGCGGGTTGCTTACCT-3'
- KMP: 5'-FAM-ACCTGTCCTCCTGTCTGCCACTTCATTGT-BHQ-1-3'

The probe was initially designed as a TaqMan probe (Yuan *et al.* 2012), however, in this report all probes are reported as BHQ-1-probes.

The primers and probe that have been designed show strong species specificity to *Karenia mikimotoi* compared to the other eight “sister” species, as shown in table S5.

Table S5: Species specific primer/probe assay for *Karenia mikimotoi* with, target gene, product size, melting temperature, primer/probe length, GC ratio (%), and number of mismatches between primer and probe region in closely related non-target species

Species	Gene	Product size	Temp	Length	GC
<i>Karenia mikimotoi</i>	ITS	112 bp			
KMF	CITTGTTGTAACCTGTTGCTTGT		62.5	25	40
KMR	TCAGCGGGTTGCTTACCT		57.5	19	53
KMP	ACCTGTCCTCCTGTCTGCCACTTCATTGT		72.1	30	50
Related species	KMF	KMR	KMP		
<i>K. asterichroma</i>	18	11	18		
<i>K. bidigitata</i>	3+7 gasps	1	8		
<i>K. brevis</i>	17	11	17		
<i>K. brevisulcata</i>	19	11	17		
<i>K. cristata</i>	17	11	14		
<i>K. papilionacea</i>	16	11	16		
<i>K. selliformis</i>	7 gaps	1	10		
<i>K. umbella</i>	18	11	15		

Species 6: Oncorhynchus mykiss

Full name: *Oncorhynchus mykiss* (Walbaum, 1792).

The genus *Oncorhynchus* belongs to the family Salmonidae and comprises 15 species (Froese and Pauly, 2015)

Occurrences of hybridization between *O. mykiss* and *Salmo salar* have been reported. Since the mitochondrial genome of these hybrids is maternally inherited the assay will be unable to distinguish hybrids from normal individuals in cases where the mitochondrion genome originates from *O. mykiss* (Wang *et al.* 2015). *Oncorhynchus mykiss* showed high similarity to *O. gilae* and *O. Chrysogaster* both with regard to the CO1 and Cytochrome b sequences. However, as the latter two species are of Pacific Ocean origin and neither occurs in Danish territorial waters, they do not pose a challenge for false positives. The following primers and probe are designed to target *O. mykiss*:

- Onc_Myk_Cytb_F01: 5'-ACCTCCAGCCATCTCTCAGT-3'
- Onc_Myk_Cytb_R01: 5'-AGGACGGGGAGGGAAAGTAA-3'
- Onc_Myk_Cytb_P01: 5'-FAM-TGAGCCGTGCTAGTTACTGCTGTCCTT-BHQ-1-3'

Table S6: Species-specific primer/probe assay for *Oncorhynchus mykiss* with target gene, product size, melting temperature, primer/probe length, GC ratio (%), and number of mismatches between primer and probe region in closely related non-target species.

Species	Gene	Product size	Temp	Length	GC
<i>Oncorhynchus mykiss</i>	Cytb	90 bp			
Onc_Myk_Cytb_F01	ACCTCCAGCCATCTCTCAGT		60.0	20	55
Onc_Myk_Cytb_R01	AGGACGGGGAGGGAAAGTAA		59.9	20	55
Onc_Myk_Cytb_P01	TGAGCCGTGCTAGTTACTGCTGTCCTT		67.6	27	52
Related species	F01	R01	P01		
<i>Oncorhynchus gorbuscha</i>	3	3	4		
<i>Oncorhynchus mykiss</i> ×					
<i>Salmo salar</i> _KP218514	0	0	0		
<i>Oncorhynchus mykiss</i> ×					
<i>Salmo salar</i> _NC_026537	0	0	0		
<i>Oncorhynchus gilae</i>	0	0	0		
<i>Oncorhynchus chrysogaster</i>	0	0	0		
<i>Oncorhynchus keta</i>	3	5	5		
<i>Oncorhynchus clarkii</i>	3	1	2		
<i>Oncorhynchus nerka</i>	2	3	4		
<i>Oncorhynchus tshawytscha</i>	3	3	3		
<i>Oncorhynchus kisutch</i>	3	4	3		
<i>Oncorhynchus masou</i>	4	3	3		
<i>Salmo trutta</i>	3	4	6		
<i>Salmo salar</i>	2	3	5		

Species 7: Mnemiopsis leidyi

Full name: *Mnemiopsis leidyi* Agassiz, 1865

The species *Mnemiopsis leidyi* is the only species within the genus *Mnemiopsis*. Within the nearest species under the Order *Lobata* the 18S sequence could not be used to differentiate between *Mnemiopsis leidyi* and non-target species. Therefore, to ensure sufficient specificity of the assay designed towards *Mnemiopsis leidyi*, a Blast search was conducted using several CO1 sequence available through the NCBI database. An eDNA primer/probe assay was designed based upon the alignment of in total 35 sequences from different non-target species. The assay showed a high degree of mismatches to non-target species (table S7):

- Mne_lei_CO1_F01: 5'-CTCATGTTTGTCAATTITAATTAAC-3'
- Mne_lei_CO1_R01: 5'-CTAAACAAACCTAATAAACCTATAC-3'
- Mne_lei_CO1_P01: 5'-FAM-AATGTACCTTTCTTATCCTGGCTAAAT-BHQ-1-3'

Table S7: Species specific primer/probe assay for *Mnemiopsis leidyi* with, target gene, product size, melting temperature, primer/probe length, GC ratio (%), and number of mismatches between primer and probe region in closely related non-target species

Species	Gene	Product Size	Temp	Length	GC
<i>Mnemiopsis leidyi</i>	CO1	114 bp			
Mne_lei_CO1_F01		CTCATGTTTGTCAATTITAATTAAC	56.8	26	23
Mne_lei_CO1_R01		CTAAACAAACCTAATAAACCTATAC	56.6	24	29
Mne_lei_CO1_P01		AATGTACCTTTCTTATCCTGGCTAAAT	63.9	30	30

Related species*	Forward	Reverse	Probe
<i>Cotesia salebrusa</i>	15	9	14
<i>Cotesia salebrusa voucher Cs_III_3</i>	15	9	14
<i>Cotesia salebrusa voucher Cs_II_1</i>	15	9	14
<i>Cotesia salebrusa voucher Cs_IV_1</i>	15	9	14
<i>Cotesia salebrusa voucher Cs_VII_8</i>	15	9	14
<i>Cotesia marginiventris</i>	15	9	15
<i>Cotesia griffini</i>	15	8	15
<i>Cotesia melitaearum</i>	15	7	14
<i>Aloeides arida</i>	15	8	14
<i>Ampedus ainu</i>	15	-	15
<i>Aphonopelma mojave</i>	13	10	15
<i>Callicore astarte</i>	15	9	14
<i>Catoblepia amphirhoe</i>	15	8	13
<i>Conogethes punctiferalis</i>	14	9	15
<i>Baryscapus servadeii</i>	14	8	16
<i>Hiatella</i> sp. L HML-2015	13	10	17
<i>Idarnes</i> sp. ex <i>Ficus goldmanii</i>	17	5	13
<i>Ischyropsalis adamii</i>	15	9	10
<i>Limenitis helmanni</i>	16	11	12
<i>Limoniscus atricolor</i>	16	-	10
<i>Lanxoblatta emarginata</i>	15	6	13
<i>Mazocraeoides gomialisae</i>	13	7	17
<i>Osmia juxta</i>	14	8	12
<i>Phenacoccus baccharidis</i>	17	10	18
<i>Principapillatus hitoyensis</i>	15	9	16
<i>Pauropsalta infuscata</i>	16	10	15
<i>Pterolophia rigida</i>	15	9	15
<i>Panorpodes paradoxus</i>	14	8	15

<i>Psepholax leoninus</i>	16	9	16
<i>Psepholax mastersii</i>	16	9	16
<i>Spermophagus brevipes</i>	16	9	14
<i>Spermophagus cornutus</i>	15	10	16
<i>Torymus beneficus</i>	14	9	14
<i>Telenomus rowani</i>	15	-	6
<i>Yranea lathonoides</i>	15	9	16

* not necessarily related but show strong sequence similarity to target species

Species 8: Eriocheir sinensis

Full name: *Eriocheir sinensis* Milne-Edwards, 1853

According to WoRMS the genus *Eriocheir* comprises four marine species. The genus belongs to the family Varunidae which contains six subfamilies and 36 genera (WoRMS 2015).

Sequence information is available for both *E. sinensis* and its sister-species. Beside a BLAST search of the assay, other native decapod species were included in the evaluation of the assay, see Table S8. Due to limited variation between *E. sinensis* and *E. Japonica* the assay cannot separate these two species. However, both *E. Japonica* and *E. ogasawaraensis* only occur in the western Pacific (Yamasaki *et al.* 2006; Komai *et al.* 2006; WoRMS 2015) and are therefore not expected to result in false positives for samples collected in Danish territorial waters. The following primers and probe are designed to target *E. sinensis*:

- Eri_Sin_Cytb_F01: 5'-TCGGTACCGACCTAGTACAA-3'
- Eri_Sin_Cytb_R01: 5'-AGAAAATGCTGATGCTACTAAAGGT-3'
- Eri_Sin_Cytb_P01: 5'-FAM-TGAGGAGGGTTCTGTTGATAATGCCAC-BHQ-1-3'

Table S8: Species-specific primer/probe assay for *Eriocheir sinensis* with target gene, product size, melting temperature, primer/probe length, GC ratio (%), and number of mismatches between primer and probe region in closely related non-target species.

Species	Gene	Product size	Temp	Length	GC
<i>Heterosigma akashiwo</i>	Cytb	113 bp			
<i>Eri_Sin_Cytb_F01</i>	TCGGTACCGACCTAGTACAA		57.2	20	50
<i>Eri_Sin_Cytb_R01</i>	AGAAAATGCTGATGCTACTAAAGGT		58.8	25	36
<i>Eri_Sin_Cytb_P01</i>	TGAGGAGGGTTCTGTTGATAATGCCAC		65.3	29	45

Related species	F01	R01	P01
<i>Eriocheir ogasawaraensis</i>	1	0	3
<i>Eriocheir japonica</i>	0	0	0
<i>Eriocheir hepuensis</i>	2	0	1
<i>Paralithodes camtschaticus</i>	8	10	6
<i>Nephrops norvegicus</i>	5	11	5
<i>Homarus gammarus</i>	5	10	5
<i>Homarus americanus</i>	5	9	5

Species 9: Rhithropanopeus harrisii

Full name: *Rhithropanopeus harrisii* (Gould, 1841)

The north American mud crab (*R. harrisii*) is a monotypic species for *Rhithropanopeus* (Türkay 2001, WoRMS Editorial Board 2015). The family Panopeidae holds 27 genera, and 22 of these are represented by various sequences on the NCBI database. A comparative study on mitochondrial haplotypes found in *R. harrisii* in Eastern Denmark (Aagaard 2015) and a quick comparison of cytochrome oxidase 1 (CO1) sequences from NCBI infers a close relationship between *Rhithropanopeus*, *Eurypanopeus* and *Hexapanopeus*.

For the CO1 gene in the mt-genome, there appear to be five haplotypes of *R. harrisii* present in Danish seas (Aagaard 2015). The primer-probe system recommended here is assumed to be able to differentiate between *Rhithropanopeus*, *Eurypanopeus* and *Hexapanopeus*. A European checklist (Türkay 2001) lists only *Dyspanopeus sayi* (Smith, 1869), *Panopeus africanus* A. Milne-Edwards, 1867, and *Rhithropanopeus harrisii* from the family Panopeidae in European coastal seas. This limits the number of potential non-target species that might be co-occurring with the target-species (*R. harrisii*), and thus reduces the risk of obtaining false positives from other representatives of Panopeidae. The primers and probes recommended here are matched against CO1 sequences from multiple representatives of Panopeidae (Table S9).

- Rhi_har_CO1_F01: 5'-CCACCATCCCTTACACTCCT-3',
- Rhi_har_CO1_R01: 5'-CTCCTGCATGAGCAATAGCA-3',
- Rhi_har_CO1_P01: 5'-FAM-GAAAGAGGAGTTGGAACAGGATGAAC-TG-BHQ-1-3',

Table S9: Species specific primer/probe assay for *Rhithropanopeus harrisii* with, target gene, product size, melting temperature, primer/probe length, GC ratio (%), and number of mismatches between primer and probe region in closely related non-target species.

Species	Gene	Product size	Temp	Length	GC
<i>Rhithropanopeus harrisii</i>	CO1	106 bp			
Rhi_har_CO1_F01		CCACCATCCCTTACACTCCT	58.9	20	55
Rhi_har_CO1_R01		CTCCTGCATGAGCAATAGCA	60.1	20	50
Rhi_har_CO1_P01		GAAAGAGGAGTTGGAACAGGATGAAC-TG	67.2	28	46

Related species	Forward	Reverse	Probe
<i>Acantholobulus bermudensis</i>	6	1	0
<i>Acantholobulus pacificus</i>	3	2	1
<i>Acantholobulus schmitti</i>	2	2	1
<i>Chasmophora macrophthalmia</i>	4	1	4
<i>Cyrtoplax panamensis</i>	4	6	4
<i>Dyspanopeus sayi</i>	2	4	1
<i>Dyspanopeus texanus</i>	2	3	1
<i>Eurypanopeus abbreviatus</i>	4	4	1
<i>Eurypanopeus depressus</i>	2	2	1
<i>Eurypanopeus dissimilis</i>	2	4	2
<i>Eurypanopeus ovatus</i>	3	4	1
<i>Eurypanopeus planissimus</i>	4	6	2
<i>Eurypanopeus planus</i>	4	4	4
<i>Eurypanopeus turgidus</i>	2	2	1
<i>Eurytium affine</i>	3	5	2
<i>Eurytium albidigitum</i>	3	4	2
<i>Eurytium tristani</i>	2	5	1
<i>Glyptoplax pugnax</i>	2	2	2
<i>Glyptoplax smithii</i>	3	3	2
<i>Hexapanopeus angustifrons</i>	3	4	3

<i>Hexapanopeus caribbaeus</i>	2	3	2
<i>Hexapanopeus lobipes</i>	5	2	5
<i>Hexapanopeus paulensis</i>	2	3	4
<i>Lophopanopeus bellus</i>	2	2	4
<i>Lophoxanthus lamellipes</i>	4	4	6
<i>Malacoplax californiensis</i>	5	3	1
<i>Metopocarcinus concavatus</i>	3	3	2
<i>Neopanope packardii</i>	2	5	3
<i>Panopeus africanus</i>	2	4	1
<i>Panopeus americanus</i>	2	3	1
<i>Panopeus austrobesus</i>	3	5	1
<i>Panopeus chilensis</i>	2	4	1
<i>Panopeus hartii</i>	2	3	1
<i>Panopeus herbstii</i>	2	4	1
<i>Panopeus lacustris</i>	3	5	1
<i>Panopeus obesus</i>	2	5	1
<i>Panopeus occidentalis</i>	2	4	1
<i>Panopeus purpureus</i>	3	5	1
<i>Panopeus rugosus</i>	3	5	1
<i>Panopeus simpsoni</i>	2	3	1
<i>Panoplax depressa</i>	3	4	1
<i>Rhithropanopeus harrisi</i>	0	0	0
<i>Robertrella mystica</i>	5	2	5
<i>Tetraplax quadridentata</i>	3	4	2
<i>Tetraxanthus rathbunae</i>	1	4	1
<i>Thalassoplax angusta</i>	5	2	5

Species 10: Prorocentrum minimum

Full name: *Prorocentrum minimum* (Pavillard) Schiller, 1933

Prorocentrum minimum belongs to the genus *Prorocentrum* that comprises 85 species. Primer/probe assay was designed after Scorzetti *et al.* (2009). The authors had selected the D1D2 region of the 28S sequence, which commonly selected for assay design for algae species (e.g., Dittami *et al.* 2013). The only modification to the otherwise good assay was the design of a new reverse primer to potentially provide better species-specificity to *Prorocentrum minimum*. In addition three extra bases were added to the probe to increase temperature and make it longer than the two primers:

- Pro_min_28S_F01: 5'-GGGTAGGTAGCTCGTCTA-3'
 - Pro_min_28S_R01: 5'-CAGGTCCCCGCCAGT-3'
 - Pro_min_28S_P01 (PI3*): 5'-GCAGTGTCCITGGCATTTCTGGAAG-3'
- * original probe name without AAG at the end (Scorzetti *et al.* 2009)

Besides the perfect sequence similarity to *P. balticum* the number of base pair mismatch to closest related species, where it was possible to align sequences, was high (Table S10). In total 34 species within the genus *Prorocentrum* had been sequenced for the gene region of interest in the assay design. Please note that for the probe the number of mismatches to non-target sister species was not recorded due to 11 extra bases in several of these non-target species. However, mismatch in the direct comparable sequence was high (6–12 bp) for all, except *P. balticum* (0 bp).

Table S10: Species specific primer/probe assay for *Prorocentrum minimum* with, target gene, product size, melting temperature, primer/probe length, GC ratio (%), and number of mismatches between primer and probe region in closely related non-target species (- = missing nucleotide possibly due to gap insertion for alignment in BioEdit (Hall 1999)). ND = not done due to large gap insertion. Problematic species are marked in bold

Species	Gene	Size	Temp	Length	GC
<i>Prorocentrum minimum</i>	28S	134 bp			
Pro_min_28S_F01	GGGTAGGTAGCTCGTCTA		60.5	20	55
Pro_min_28S_R01	CAGGTCCCCGCCAGT		58.4	16	75
Pro_min_28S_P01	GCAGTGTCCITGGCATTTCTGGAAG		66.9	24	54
Related species	Forward	Reverse	Probe		
<i>Prorocentrum arabianum</i>	11	9	ND		
<i>Prorocentrum arenarium</i>	10	11	ND		
<i>Prorocentrum balticum</i>	0	0	ND		
<i>Prorocentrum bimaculatum</i>	10	12	ND		
<i>Prorocentrum concavum</i>	11	9	ND		
<i>Prorocentrum consutum</i>	9	12	ND		
<i>Prorocentrum dentatum</i>	2	5	ND		
<i>Prorocentrum donghaiense</i>	2	5	ND		
<i>Prorocentrum elegans</i>	4	-	ND		
<i>Prorocentrum emarginatum</i>	11	-	ND		
<i>Prorocentrum</i> <i>emarginatum/fukuyoi</i>	10	8	ND		
<i>Prorocentrum faustiae</i>	11	9	ND		
<i>Prorocentrum foveolatum</i>	6	14	ND		
<i>Prorocentrum fukuyoi</i>	11	10	ND		
<i>Prorocentrum gracile</i>	4	-	ND		
<i>Prorocentrum hoffmannianum</i>	11	11	ND		
<i>Prorocentrum levnis</i>	9	13	ND		
<i>Prorocentrum lima</i>	11	11	ND		

<i>Prorocentrum mexicanum</i>	4	8	ND
<i>Prorocentrum micans</i>	7	11	ND
<i>Prorocentrum panamense</i>	9	13	ND
<i>Prorocentrum playfairi</i>	6	-	ND
<i>Prorocentrum rhathymum</i>	4	8	ND
<i>Prorocentrum sculptile</i>	4	8	ND
<i>Prorocentrum sigmoides</i>	4	12	ND
<i>Prorocentrum sp. CCMP1724</i>	11	9	ND
<i>Prorocentrum sp. IRTA002</i>	6	13	ND
<i>Prorocentrum sp. SM19</i>	9	10	ND
<i>Prorocentrum sp. Y8</i>	11	9	ND
<i>Prorocentrum texanum</i>	6	11	ND
<i>Prorocentrum triestinum</i>	7	11	ND
<i>Prorocentrum compressum</i>	6	9	ND

Species 11: *Gracilaria vermiculophylla*

Full name: *Gracilaria vermiculophylla* (Ohmi) Papenfuss (1967)

The genus *Gracilaria* contains 99 species and varietates. The most available sequence from these species was the *rbcL* gene. For some species this was the only sequences gene. However, for the following *Gracilaria* species the *rbcL* gene was not available: *G. abbottiana*, *G. brasiliensis*, *G. cearensis*, *G. conseroides*, *G. crassa*, *G. dawsonii*, *G. dentata*, *G. epihippisora*, *G. ferox*, *G. foliifera* var. *angustissima*, *G. aff. lacinulata*, *G. aff. mammillaris*, *G. manilaensis*, and *G. aff. tepecensis*.

Sequence information from the following species could not be aligned:

G. cliftonii, *G. corticata*, *G. corticata* var. *corticata*, *G. delibis*, *G. denticulate*, *G. euchaeumoides*, *G. fergusonii*, *G. foliifera*, *G. gigas*, *G. millardentii*, and *G. robusta*.

The following species showed identical sequences to *Gracilaria vermiculophylla*:

G. asiatica (AY769260.1, AY769261.1), *G. tenuistipita* (AY049312.1) and *G. tenuistipitata*, (AY049313.1, AY049314.1, AY049324.1). Some confusion remains around *G. textorii* since HQ880647.1 shows on differentiation to target sequence, whereas DQ095814.1 (shown in table S11.1) shows decent differentiation.

Based on the sequences available in NCBI and that could be aligned the following assay was designed:

- Gra_ver_rbcL_F01: 5'-CTTCCAGGGGCCAGCA-3'
- Gra_ver_rbcL_R01: 5'-AAAGGACGACCAAACCTTATCC-3'
- Gra_ver_rbcL_P01: 5'-FAM-GGACTGGTTGTAGAACGTGAACGTAT-BHQ-1-3'

The assays base pair differentiation to related species was 3.7, 2.1, and 4.0 on average for the F01, R01, and P01 primers and probe, respectively. The exact differences are listed in table S11.1. Please note here that the assay could not be designed to not target *Gracilaria asiatica*.

Table S11.1: Species specific primer/probe assay for *Gracilaria vermiculophylla* with, target gene, product size, melting temperature, primer/probe length, GC ratio (%), and number of mismatches between primer and probe region in closely related non-target species. Problematic species are marked in bold

Species	Gene	Product size	Temp	Length	GC
<i>Gracilaria vermiculophylla</i>	rbcL	67 bp			
Gra_ver_rbcL_F01	CTTCCAGGGGCCAGCA		57.3	17	65
Gra_ver_rbcL_R01	AAAGGACGACCAAACCTTATCC		57.5	21	43
Gra_ver_rbcL_P01	GGACTGGTTGTAGAACGTGAACGTAT		66.2	26	46

Related species	F01	R01	P01
<i>Gracilaria apiculata</i>	4	2	4
<i>Gracilaria arcuata</i>	3	1	5
<i>Gracilaria armata</i>	4	3	4
<i>Gracilaria asiatica</i>	0	0	0
<i>Gracilaria babae</i>	3	1	4
<i>Gracilaria beckeri</i>	3	0	7
<i>Gracilaria birdiae</i>	4	3	3
<i>Gracilaria blodgettii</i>	6	2	3
<i>Gracilaria bursa-pastoris</i>	3	3	5
<i>Gracilaria cacalia</i>	3	1	4
<i>Gracilaria canaliculata</i>	3	1	4
<i>Gracilaria capensis</i>	3	1	5
<i>Gracilaria caudata</i>	4	2	2

<i>Gracilaria cervicornis</i>	4	2	4
<i>Gracilaria changii</i>	6	2	2
<i>Gracilaria chilensis</i>	4	2	4
<i>Gracilaria chonae</i>	4	3	5
<i>Gracilaria conferta</i>	6	3	4
<i>Gracilaria cornea</i>	4	3	3
<i>Gracilaria coronopifolia</i>	4	4	5
<i>Gracilaria crassissima</i>	4	3	3
<i>Gracilaria cuneata</i>	4	2	3
<i>Gracilaria cuneifolia</i>	4	2	5
<i>Gracilaria curtissiae</i>	4	2	3
<i>Gracilaria cylindrica</i>	4	1	4
<i>Gracilaria damaecornis</i>	4	3	4
<i>Gracilaria domingensis</i>	4	2	4
<i>Gracilaria dotyi</i>	3	4	5
<i>Gracilaria dura</i>	4	2	3
<i>Gracilaria edulis</i>	4	2	4
<i>Gracilaria eucheumatoides</i>	4	1	3
<i>Gracilaria firma</i>	6	2	3
<i>Gracilaria flabelliforme</i>	4	3	4
<i>Gracilaria gracilis</i>	4	3	3
<i>Gracilaria galetensis</i>	4	3	5
<i>Gracilaria bayi</i>	4	2	5
<i>Gracilaria huangii</i>	3	1	4
<i>Gracilaria incurvata</i>	3	2	5
<i>Gracilaria intermedia</i>	3	1	5
<i>Gracilaria irregularis</i>	5	2	2
<i>Gracilaria isabellana</i>	4	1	4
<i>Gracilaria lacinulata</i>	4	1	5
<i>Gracilaria longa</i>	3	2	4
<i>Gracilaria mammillaris</i>	4	4	5
<i>Gracilaria multipartita</i>	2	2	4
<i>Gracilaria occidentalis</i>	4	3	4
<i>Gracilaria oliveirarum</i>	4	3	5
<i>Gracilaria ornata</i>	4	3	4
<i>Gracilaria pacifica</i>	4	2	4
<i>Gracilaria parrispora</i>	4	3	5
<i>Gracilaria punctata</i>	4	2	3
<i>Gracilaria rhodymenoides</i>	4	2	5
<i>Gracilaria salicornia</i>	3	1	4
<i>Gracilaria secundata</i>	4	2	4
<i>Gracilaria smithsoniensis</i>	4	3	5
<i>Gracilaria spinulosa</i>	6	1	5
<i>Gracilaria stipitata</i>	3	1	5
<i>Gracilaria taiwanensis</i>	4	3	4
<i>Gracilaria tenuistipitata</i>	4	1	4
<i>Gracilaria textorii</i>	3	2	4
<i>Gracilaria tikvahiae</i>	4	2	5
<i>Gracilaria truncata</i>	4	2	4
<i>Gracilaria urvilleii</i>	4	1	2
<i>Gracilaria usneoides</i>	4	3	3
<i>Gracilaria venezuelensis</i>	3	1	5
<i>Gracilaria viellardii</i>	4	3	4
<i>Gracilaria yonesigueana</i>	3	2	5

Due to relatively large proportions of missing data for *rbcL* gene, an attempt to construct an assay for CO1 was also undertaken. However, a smaller proportion of the species have been sequenced for CO1 compared to *rbcL* (table S11.1 vs. table S11.2). The assay for *Gracilaria vermiculophylla* based on CO1:

- Gra_ver_CO1_F02: 5'-CGTCAGCAGTTGTAGAAGTC-3'
- Gra_ver_CO1_R02: 5'-AGAGGCTCCTGATATGTGC-3'
- Gra_ver_CO1_P02: 5'-FAM-CTCAGGCAGCTGTAGATTAGCT-BHQ-1-3'

Despite the lower number of sequences species within the genus *Gracilaria* the designed assay showed on average higher degrees of differentiation to sister species compared to *rbcL* (please note that the list of sister species is not directly comparable between tables S11.1 and S11.2). Averages were F02 2.9, R02 4.5, and P02 6.4. However, *Gracilaria corticata* and *Gracilaria foliifera* showed none or only one mismatch to the designed assay. In general these two species showed very little mismatch in the sequence and not in positions that would make the design of an assay possible.

Table S11.2: Species specific primer/probe assay for *Gracilaria vermiculophylla* with, target gene, product size, melting temperature, primer/probe length, GC ratio (%), and number of mismatches between primer and probe region in closely related non-target species. Problematic species are marked in bold

Species	Gene	Product size	Temp	Length	GC
<i>Gracilaria vermiculophylla</i>	CO1	109 bp			
Gra_ver_CO1_F02	CGTCAGCAGTTGTAGAAGTC		58.4	20	50
Gra_ver_CO1_R02	AGAGGCTCCTGATATGTGC		57.5	19	53
Gra_ver_CO1_P02	CTCAGGCAGCTGTAGATTAGCT		67.4	25	67.4
Related species	F02	R02	P02		
<i>Gracilaria corticata</i>	1	0	0		
<i>Gracilaria foliifera</i>	0	0	0		
<i>Gracilaria flabelliformis</i> subsp. <i>simplex</i>	4	4	5		
<i>Gracilaria arcuata</i>	2	5	8		
<i>Gracilaria manilaensis</i>	4	4	8		
<i>Gracilaria salicornia</i>	1	6	5		
<i>Gracilaria smithsoniensis</i>	5	4	5		
<i>Gracilaria bayi</i>	4	6	6		
<i>Gracilaria dentata</i>	4	3	7		
<i>Gracilaria damaecornis</i>	4	4	7		
<i>Gracilaria curtissiae</i>	4	6	6		
<i>Gracilaria cervicornis</i>	5	3	8		
<i>Gracilaria birdiae</i>	4	5	5		
<i>Gracilaria gracilis</i>	3	4	7		
<i>Gracilaria chouae</i>	6	5	4		
<i>Gracilaria textorii</i>	5	4	8		
<i>Gracilaria cornea</i>	5	5	5		
<i>Gracilaria yonashigueana</i>	4	5	7		
<i>Gracilaria isabellana</i>	4	4	6		
<i>Gracilaria ferox</i>	5	3	7		
<i>Gracilaria domingensis</i>	5	3	8		
<i>Gracilaria cuneata</i>	4	5	8		
<i>Gracilaria caudata</i>	5	3	7		
<i>Gracilaria pacifica</i>	4	4	8		
<i>Gracilaria tenuistipitata</i> var. <i>liui</i>	4	6	6		
<i>Gracilaria chouae</i>	6	5	4		
<i>Gracilaria blodgettii</i>	6	4	6		
<i>Gracilaria salicornia</i>	2	6	5		

Species 12: Sargassum muticum

Full name: *Sargassum muticum* (Yendo) Fensholt, 1955

Within the Genus *Sargassum* there are 121 species and variates. Due to the large number of species within this genus it was very challenging to identify a gene with enough variation between *Sargassum muticum* and the remaining members. Despite large availability of genes within this genus it was not possible to design any assay using sequence information from the following genes *rbcL*, 28S, *psaA*, ITS, and CO1. After extensive search the gene region COX3 was identified as a good alternative for designing an assay.

Several possibilities to position elements of an assay were identified within the published sequences for COX3 showing a good amount of mismatches to non-target species. However, the majority of these showed the potential for secondary structures or self-annealing, and could therefore not with confidence be used to construct an assay. Despite this it was possible to design an assay for *Sargassum muticum* from the COX3 gene, however with relatively modest numbers of mismatch to non-target species, as represented in table S12.

- Sar_mut_COX3_F01: 5'-TGCACCTTCGCTATGTCC-3'
- Sar_mut_COX3_R01: 5'-GGTACCAATAATAACATGGAAAC-3'
- Sar_mut_COX3_P01: 5'-FAM-CAGTGTATGGTTCTGTITTTATGGC-BHQ-1-3'

In the event that the assay should not be specific enough to *Sargassum muticum* an alternative primer pair without probe was designed. A probe could not be design due to formation of secondary structures preventing annealing to target strand. Despite the lack of a probe the primer pair should be successful due to a higher degree of mismatch to non-target species (not shown) compared to assay system 1 (above):

- Sar_mut_COX3_F02: 5'-TTATTTACTTCTTCCTCGCGCT-3'
57.6°C, 25 bp long, and GC% = 28.
- Sar_mut_COX3_R02: 5'-CAAATATTATTGTAATGAGCAAACCTT-3'
56.8°C, 26 bp long, and GC% = 23.
Product length including primers, 104 bp

Table S12: Species specific primer/probe assay for *Sargassum muticum* with, target gene, product size, melting temperature, primer/probe length, GC ratio (%), and number of mismatches between primer and probe region in closely related non-target species. Problematic species are marked in bold

Species	Gene	Product size	Temp	Length	GC
<i>Sargassum muticum</i>	COX3	84 bp			
Sar_mut_COX3_F01	TGCACCTTCGCTATGTCC		57.5	19	53
Sar_mut_COX3_R01	GGTACCAATAATAACATGGAAAC		57.6	23	35
Sar_mut_COX3_P01	CAGTGTATGGTTCTGTITTTATGGC		65.1	30	33

Related species	F02	R02	P02
<i>Sargassum aquifolium</i>	4	6	2
<i>Sargassum autumnale</i>	3	6	4
<i>Sargassum binderi</i>	4	6	2
<i>Sargassum brandegeei</i>	3	5	3
<i>Sargassum carpophyllum</i>	3	5	2
<i>Sargassum cf. cymosum</i>	3	5	2
<i>Sargassum cf. latifolium</i>	3	6	2
<i>Sargassum cf. polyceratum</i>	3	5	2
<i>Sargassum cf. yamamotoi</i>	4	5	4
<i>Sargassum coreanum</i>	4	5	4
<i>Sargassum cymosum</i>	3	5	2

<i>Sargassum cymosum</i> var. <i>scabriuscum</i>	3	5	2
<i>Sargassum echinocarpum</i>	4	6	2
<i>Sargassum elegans</i>	3	5	2
<i>Sargassum fallax</i>	3	6	4
<i>Sargassum filicinum</i>	1	3	1
<i>Sargassum filipendula</i>	3	5	2
<i>Sargassum fluitans</i>	3	5	2
<i>Sargassum fulvellum</i>	1	1	1
<i>Sargassum fusiforme</i>	4	6	2
<i>Sargassum giganteum</i>	3	5	2
<i>Sargassum hemiphyllum</i>	2	5	1
<i>Sargassum herporhizum</i>	3	5	3
<i>Sargassum horneri</i>	1	3	1
<i>Sargassum horridum</i>	3	5	3
<i>Sargassum howeanum</i>	3	5	2
<i>Sargassum hystrix</i>	3	5	2
<i>Sargassum ilicifolium</i>	3	5	2
<i>Sargassum ilicifolium</i> var. <i>acaraeocarpum</i>	3	5	1
<i>Sargassum johnstonii</i>	3	5	4
<i>Sargassum lajaeanum</i>	3	5	3
<i>Sargassum linearifolium</i>	3	5	2
<i>Sargassum macrocarpum</i>	3	6	4
<i>Sargassum marginatum</i>	3	5	2
<i>Sargassum micracanthum</i>	3	6	4
<i>Sargassum miyabei</i>	1	1	0
<i>Sargassum natans</i>	3	5	2
<i>Sargassum nigrifolium</i>	3	6	4
<i>Sargassum obovatum</i>	3	6	2
<i>Sargassum obovatum</i> var. <i>pfeifferae</i>	3	5	2
<i>Sargassum obovatum</i> var. <i>robillardii</i>	2	6	2
<i>Sargassum obtusifolium</i>	3	5	2
<i>Sargassum pacificum</i>	3	5	2
<i>Sargassum paradoxum</i>	3	6	4
<i>Sargassum patens</i>	3	6	3
<i>Sargassum pilularium</i>	3	5	2

Species 13: *Cordylophora caspia*

Full name: *Cordylophora caspia* (Pallas, 1771)

The genus *Cordylophora* comprises three valid species *Cordylophora caspia* (Pallas, 1771), *Cordylophora japonica* Itô, 1951 and *Cordylophora solangiae* Redier, 1967 (WoRMS Editorial Board 2015). Only nucleotide sequence data for *Cordylophora caspia* is available on NCBI GenBank. The comparison between sequences from closely related non-target species is therefore based on nucleotide sequence data from other representatives of the family Oceaniidae. Without sequence information available from *C. japonica* and *C. solangiae* it is currently not possible to determine whether the recommended primer-probe system for detection of eDNA from *Cordylophora caspia* will be able to differentiate between *C. japonica* and *C. solangiae* and *Cordylophora caspia*.

The recommended primer-probe system for *Cordylophora caspia* should be able to detect eDNA originating from the mtDNA cytochrome oxidase 1 gene (Table S13.1).

- Cor_cas_CO1_F01: 5'-TCATCTGTACAAGCACATTCTGG-3',
- Cor_cas_CO1_R01: 5'-TTGAAGAAGCTCCTGCACAGT-3',
- Cor_cas_CO1_P01: 5'-FAM-CCTCTGTAGACATGGCTATTTAGTC-BHQ1-3',

Table S13.1: Species specific primer/probe assay for *Cordylophora caspia* with, target gene, product size, melting temperature, primer/probe length, GC ratio (%), and number of mismatches between primer and probe region in closely related non-target species.

Species	Gene	Product size	Temp	Length	GC
<i>Cordylophora caspia</i>	CO1	76 bp			
Cor_cas_CO1_F01	TCATCTGTACAAGCACATTCTGG		60.2	23	43
Cor_cas_CO1_R01	TTGAAGAAGCTCCTGCACAGT		60.2	21	48
Cor_cas_CO1_P01	CCTCTGTAGACATGGCTATTTAGTC		59.2	28	39
Related species	Forward	Reverse	Probe		
<i>Cordylophora</i> sp. NFR1	0	0	0		
<i>Cordylophora</i> sp. NFR9	4	2	6		
<i>Turritopsis lata</i>	9	4	7		
<i>Turritopsis nutricula</i>	9	4	7		
<i>Turritopsis rubra</i>	6	3	6		

An additional primer-probe system for eDNA from the nuclear 28S region is recommended in case the first recommended system turns out to be unable to distinguish *C. japonica* and *C. solangiae* from *Cordylophora caspia* (Table S13.2):

- Cor_cas_28S_F02: 5'-ACTGGACAGAGGAGGAGTCG-3',
- Cor_cas_28S_R02: 5'-CGACGACCAACAGTGACAAG-3',
- Cor_cas_28S_P02: 5'-FAM-ACATGCTCTTGGGCTGGCCTCT-BHQ1-3',

Table S13.2: Species specific primer/probe assay for *Cordylophora caspia* with, target gene, product size, melting temperature, primer/probe length, GC ratio (%), and number of mismatches between primer and probe region in closely related non-target species.

Species	Gene	Product size	Temp	Length	GC
<i>Cordylophora caspia</i>	28S	70 bp			
Cor_cas_28S_F02	ACTGGACAGAGGAGGAGTCG		59.4	20	60
Cor_cas_28S_R02	CGACGACCAACAGTGACAAG		60.4	20	55
Cor_cas_28S_P02	ACATGCTCTTGGGCTGGCCTCT		70.1	24	54

Related species	Forward	Reverse	Probe
<i>Turritopsis dohrnii</i>	9	9	5
<i>Turritopsis lata</i>	9	9	5
<i>Turritopsis rubra</i>	9	9	4

Species 14: *Styela clava*

Full name: *Styela clava* Herdman, 1881

The genus *Styela* comprises more than 200 species worldwide (WoRMS Editorial Board 2015), and around 18 species of *Styela* can be found in European seas (Monniot 2001), including *Styela clava*. A phylogenetic study on the families Pyuridae and Styelidae (Pérez-Portela *et al.* 2009) found a close relationship between *Styela montereyensis*, *S. gibbsii*, *S. plicata* and *Styela clava*. *Styela plicata* is occurring in the Atlantic and Mediterraen Sea, and could therefore lead to false positive eDNA detection.

Because very few species of the 244 species of *Styela* (WoRMS Editorial Board 2015) have nucleotide sequence data deposited on the NCBI GenBank database (Nov-2015), it is currently impossible to rule out that the suggested primer and probe systems for detection of mitochondrial eDNA fragments from *Styela clava* potentially could give rise to false positive eDNA detection because of nucleotide similarity with other co-occurring species of *Styela*.

In an attempt to minimize this risk of obtaining false positive eDNA detection due to other co-occurring non-sequenced species of *Styela*, a second primer-probe system for eDNA detection from mitochondrial cytochrome oxidase 1 gene fragments of *S. clava* is here recommended used together with the first recommended systems that targets eDNA from the mitochondrial cytochrome b region. Hopefully, the combination of these two primer-probe systems used together will reduce the number of false positives that might arise due to similarity in mitochondrial gene regions from the non-sequenced closely related species of *Styela*. Because there appears to limited variation among mitochondrial haplotypes among *S. clava* (Goldstien *et al.* 2011), there might be a risk of not detecting haplotypes that are deviating from the most common haplotype among *S. clava* (Goldstien *et al.* 2011). Here two different species-specific systems are recommended for targeting eDNA fragments from mitochondrial cytochrome oxidase 1 from *S. clava* (Table S14.1, Table S14.2).

- Sty_cla_Cytb_F01: 5'-GAGTITTGAGGGGCAACTG-3',
- Sty_cla_Cytb_R01: 5'-TGGCCAATGAAAACCTACC-3',
- Sty_cla_Cytb_P01: 5'-FAM-TGAGTGTAAATCCCTTITTATGGGGTGG-BHQ-1-3',

Table S14.1: Species-specific primer/probe assay for *Styela clava* with target gene, product size, melting temperature, primer/probe length, GC ratio (%), and number of mismatches between primer and probe region in closely related non-target species.

Species	Gene	Product size	Temp	Length	GC
<i>Styela clava</i>	Cytb	102 bp			
Sty_cla_Cytb_F01	GAGTITTGAGGGGCAACTG		59.7	20	50
Sty_cla_Cytb_R01	TGGCCAATGAAAACCTACC		60.0	20	50
Sty_cla_Cytb_P01	TGAGTGTAAATCCCTTITTATGGGGTGG		67.5	27	44

Related species	Forward	Reverse	Probe
<i>Botrylloides leachii</i>	5	12	6
<i>Botrylloides nigrum</i>	4	8	6
<i>Botrylloides pizoni</i>	4	9	7
<i>Botrylloides violaceus</i>	4	9	10
<i>Botryllus planus</i>	4	9	11
<i>Cnemidocarpa finmarkiensis</i>	2	8	9
<i>Dendrodoa grossularia</i>	3	9	6
<i>Metandrocarpa taylori</i>	4	8	12
<i>Polycarpa mytiligera</i>	4	7	6
<i>Polycarpa papillata</i>	2	7	8
<i>Styela clava</i>	0	0	0
<i>Styela gibbsii</i>	6	9	8

<i>Styela montereyensis</i>	4	8	7
<i>Styela plicata</i>	4	7	8
<i>Symplegma viride</i>	4	7	8

- Sty_cla_CO1_F02: 5'-GTAATTGTTACGGCCCATGC-3',
- Sty_cla_CO1_R02: 5'-CGGACTTCCCAACATCAAAG-3',
- Sty_cla_CO1_P02: 5'-FAM-TGCCGGTAATAATTAGAAGATTAGGA-BHQ1-3',

Table S14.2: Species-specific primer/probe assay for *Styela clava* with target gene, product size, melting temperature, primer/probe length, GC ratio (%), and number of mismatches between primer and probe region in closely related non-target species.

Species	Gene	Product size		Temp	Length	GC
<i>Styela clava</i>	CO1	105 bp				
Sty_cla_CO1_F02		GTAATTGTTACGGCCCATGC		60.2	20	50
Sty_cla_CO1_R02		CGGACTTCCCAACATCAAAG		60.5	20	50
Sty_cla_CO1_P02		TGCCGGTAATAATTAGAAGATTAGGA		61.5	27	33
Related species	Forward	Reverse	Probe			
<i>Boltenia ovifera</i>	5	7	8			
<i>Botrylloides fuscus</i>	4	8	7			
<i>Botrylloides leachii</i>	4	6	10			
<i>Botrylloides nigrum</i>	5	3	5			
<i>Botrylloides perspicuus</i>	1	5	8			
<i>Botrylloides pizzoni</i>	1	5	8			
<i>Botrylloides violaceus</i>	2	5	8			
<i>Botryllus schlosseri</i>	5	4	6			
<i>Botryllus tyrensis</i>	4	5	6			
<i>Dendrodoa grossularia</i>	4	6	6			
<i>Distomus variolosus</i>	4	8	6			
<i>Halocynthia papillosa</i>	4	8	9			
<i>Halocynthia pyriformis</i>	3	9	5			
<i>Herdmania grandis</i>	5	6	9			
<i>Microcosmus claudicans</i>	4	3	5			
<i>Microcosmus polymorphus</i>	1	5	10			
<i>Microcosmus squamiger</i>	4	5	7			
<i>Polyandrocarpa zorritensis</i>	3	5	9			
<i>Polycarpa aurata</i>	4	4	9			
<i>Polycarpa pomaria</i>	5	5	9			
<i>Polycarpa tenera</i>	5	5	9			
<i>Polyzoa opuntia</i>	5	6	10			
<i>Pyura australis</i>	5	5	9			
<i>Pyura dura</i>	3	3	7			
<i>Pyura gibbosa</i>	5	4	9			
<i>Pyura praefutialis</i>	5	4	7			
<i>Pyura spinifera</i>	3	2	9			
<i>Pyura squamulosa</i>	5	3	7			
<i>Styela canopus</i>	3	5	8			
<i>Styela clava</i>	0	0	0			
<i>Styela gibbsii</i>	3	5	6			
<i>Styela montereyensis</i>	3	5	5			
<i>Styela plicata</i>	5	5	7			
<i>Symplegma rubra</i>	3	10	7			

Species 15: Dreissena polymorpha

Full name: *Dreissena polymorpha* (Pallas, 1771)

According to the Worms database (WoRMS Editorial Board 2015) the genus *Dreissena* comprises 11 species: *Dreissena anatolica* Locard, 1893, *D. bourguignati* Locard, 1883, *D. bugensis* (Andrusov, 1897), *D. caputlacus* Schütt, 1993, *D. carinata* (Dunker, 1853), *D. caspia* Eichwald, 1855, *D. elata* Andrusov, 1897, *D. gallandi* Locard, 1893, *D. polymorpha* (Pallas, 1771), *D. rostriformis* (Deshayes, 1838) and *D. sioffi* Locard, 1893. From the NCBI GenBank database five species of *Dreissena* have nucleotide sequence information deposited, including *D. polymorpha*. *Dreissena polymorpha* is native to the Caspian Sea (Kantor *et al.* 2010).

The nucleotide sequence comparisons made here are made with the five species of *Dreissena* represented on NCBI GenBank: *Dreissena blanca*, *D. caputlacus*, *D. polymorpha*, *D. presbensis*, and *D. rostriformis*. The nucleotide sequence comparisons have been made with representatives of Dreissenidae and the recommended primer-probe systems may very well be unable to differentiate between *D. polymorpha* and other species of *Dreissena* not represented on NCBI GenBank. Additional sampling and sequencing of other species of *Dreissena* would be required to verify whether these suggested primer- probe system are species-specific (Table S15.1, Table S15.2, Table S15.3).

- Dre_pol_CO1_F01: 5'-CACCACTCATGGGCTTGTAA-3',
- Dre_pol_CO1_R01: 5'-ACCAATCAATTTCGAATCC-3',
- Dre_pol_CO1_P01: 5'-FAM-TTTTCTAGTAATACTATAATGGGG-BHQ-1-3',

Table S15.1: Species-specific primer/probe assay for *Dreissena polymorpha* with target gene, product size, melting temperature, primer/probe length, GC ratio (%), and number of mismatches between primer and probe region in closely related non-target species.

Species	Gene	Product size	Temp	Length	GC
<i>Dreissena polymorpha polymorpha</i>	CO1	75 bp			
Dre_pol_CO1_F01		CACCACTCATGGGCTTGTAA	59.6	20	50
Dre_pol_CO1_R01		ACCAATCAATTTCGAATCC	58.7	20	40
Dre_pol_CO1_P01		TTTTCTAGTAATACTATAATGGGG	57.3	29	28

Related species	Forward	Reverse	Probe
<i>Dreissena blanca</i>	4	2	4
<i>Dreissena polymorpha anatolica</i>	0	0	1
<i>Dreissena polymorpha gallandi</i>	0	0	0
<i>Dreissena polymorpha polymorpha</i>	0	0	0
<i>Dreissena presbensis</i>	3	3	4

- Dre_pol_CO1_F02: 5'-TCAGCTTTAGGGAAGGAGGA-3',
- Dre_pol_CO1_R02: 5'-GCAGGGCCTGAATGTCCTAT-3',
- Dre_pol_CO1_P02: 5'-FAM-GGGGTGGTTGAACCTTATATCCTCCT-BHQ-1-3',

Table S15.2: Species-specific primer/probe assay for *Dreissena polymorpha* with target gene, product size, melting temperature, primer/probe length, GC ratio (%), and number of mismatches between primer and probe region in closely related non-target species.

Species	Gene	Product size	Temp	Length	GC
<i>Dreissena polymorpha polymorpha</i>	CO1	83 bp			
Dre_pol_CO1_F02		TCAGCTTTAGGGAAGGAGGA	60.3	21	48
Dre_pol_CO1_R02		GCAGGGCCTGAATGTCCTAT	61.4	20	55
Dre_pol_CO1_P02		GGGGGTGGTTGAACCTTATATCCTCCT	68.6	27	52

Related species	Forward	Reverse	Probe
<i>Congeria jalisci</i>	5	3	7
<i>Congeria kusceri</i>	4	3	6

<i>Congeria mulaomerovici</i>	5	4	6
<i>Dreissena blinci</i>	3	3	4
<i>Dreissena caputlacus</i>	6	2	5
<i>Dreissena polymorpha</i>	0	1	0
<i>Dreissena polymorpha anatolica</i>	0	1	0
<i>Dreissena polymorpha anatolica</i>	0	1	0
<i>Dreissena polymorpha gallandi</i>	0	1	1
<i>Dreissena polymorpha polymorpha</i>	0	0	0
<i>Dreissena presbensis</i>	2	3	3
<i>Dreissena rostriformis</i>	5	4	5
<i>Dreissena rostriformis bugensis</i>	5	3	5
<i>Mytilopsis sallei</i>	3	4	3

- Dre_pol_CO1_F03: 5'-GGTCCCTITGTTCTGCACAT-3',
- Dre_pol_CO1_R03: 5'-AAAGCCCCAGCTAATACAGG-3',
- Dre_pol_CO1_P03: 5'-FAM-TGGTGTGACCAGATTCCTTTAATCTT-BHQ-1-3',

Table S15.3: Species-specific primer/probe assay for *Dreissena polymorpha* with target gene, product size, melting temperature, primer/probe length, GC ratio (%), and number of mismatches between primer and probe region in closely related non-target species.

Species	Gene	Product size	Temp	Length	GC
<i>Dreissena polymorpha polymorpha</i>	CO1	78 bp			
Dre_pol_CO1_F03		GGTCCCTITGTTCTGCACAT	60.0	20	50
Dre_pol_CO1_R03		AAAGCCCCAGCTAATACAGG	58.4	20	50
Dre_pol_CO1_P03		TGGTGTGACCAGATTCCTTTAATCTT	64.1	27	37
Related species	Forward	Reverse	Probe		
<i>Congeria jalzici</i>	5	3	6		
<i>Congeria kusceri</i>	6	3	7		
<i>Congeria mulaomerovici</i>	4	3	7		
<i>Dreissena blinci</i>	3	3	6		
<i>Dreissena caputlacus</i>	4	4	5		
<i>Dreissena polymorpha</i>	0	1	0		
<i>Dreissena polymorpha anatolica</i>	1	0	0		
<i>Dreissena polymorpha anatolica</i>	1	0	0		
<i>Dreissena polymorpha gallandi</i>	0	0	0		
<i>Dreissena polymorpha polymorpha</i>	0	0	0		
<i>Dreissena presbensis</i>	3	3	6		
<i>Dreissena rostriformis</i>	4	5	5		
<i>Dreissena rostriformis bugensis</i>	0	0	0		
<i>Mytilopsis sallei</i>	0	0	0		

Acipenser

It was not always possible to design species-specific primer-probe system. For example the sturgeons in the genus *Acipenser* comprises 17 species. *Acipenser baerii*, *A. gueldenstadtii*, *A. ruthenus* and *A. stellatus* are introduced in the Baltic whereas *A. sturio* and *A. oxyrinchus* are considered as native (Zhang *et al.* 2013, Froese & Pauly 2015). Hybridization is common among species of sturgeons (Tranah *et al.* 2004, Zhang *et al.* 2013) including *A. ruthenus*, *A. baerii*, *A. gueldenstaedtii* and *A. stellatus* (Zhang *et al.* 2013). This makes it very hard to develop functional a species-specific assay. Because species of *Acipenser* hybridize regularly it is considered to be impossible to develop a generally applicable single primer-probe assay that will be able to detect eDNA from all the various mitochondrial haplotypes in the different species of *Acipenser*. Instead primer-probe assays specific for unique haplotypes from the different targeted species of *Acipenser* are recommended. The endangered and native *A. sturio* and the *A. oxyrinchus* do not appear to hybridize with *A. baerii*, *A. gueldenstadtii*, *A. ruthenus* and *A. stellatus*, and the recommended eDNA detection system should therefore be able to distinguish native species (*A. oxyrinchus* and *A. sturio*) from introduced species (*A. baerii*, *A. gueldenstadtii*, *A. ruthenus* and *A. stellatus*).

Species 16: Acipenser baerii

Full name: *Acipenser baerii* Brandt, 1869

The recommended system targeting eDNA fragments from *A. baerii* might not be able to detect all haplotype variants of *A. baerii*, and might also return false positive amplification for other species of *Acipenser* carrying an *A. baerii* haplotype variant.

- Aci_bae_CR_F01: 5'-AGAGCCGAACATTCTTGTCTG-3',
- Aci_bae_CR_R01: 5'-TGGACCTGAAATAGGAACCAG-3',
- Aci_bae_CR_P01: 5'-FAM-TGAAGGACAGTAATTGTAGAGTTCA-BHQ-1-3',

Table S16: Species-specific primer/probe assay for *Acipenser baerii* with target gene, product size, melting temperature, primer/probe length, GC ratio (%), and number of mismatches between primer and probe region in closely related non-target species.

Species	Gene	Product size	Temp	Length	GC
<i>Acipenser baerii</i>	CR	112 bp			
Aci_bae_CR_F01		AGAGCCGAACATTCTTGTCTG	59.5	21	48
Aci_bae_CR_R01		TGGACCTGAAATAGGAACCAG	59.0	21	48
Aci_bae_CR_P01		TGAAGGACAGTAATTGTAGAGTTCA	58.5	26	35
Related species	Forward	Reverse	Probe		
<i>Acipenser baerii baicalensis</i>	13	11	18		
<i>Acipenser brevirostrum</i>	14	10	18		
<i>Acipenser dabryanus</i>	11	3	7		
<i>Acipenser fulvescens</i>	2	0	4		
<i>Acipenser gueldenstaedtii</i>	0	0	4		
<i>Acipenser medirostris</i>	4	0	6		
<i>Acipenser mikadoi</i>	13	12	17		
<i>Acipenser naccarii</i>	0	0	4		
<i>Acipenser nudiventris</i>	6	1	9		
<i>Acipenser oxyrinchus</i>	18	6	20		
<i>Acipenser oxyrinchus desotoi</i>	14	12	16		
<i>Acipenser persicus</i>	0	1	5		
<i>Acipenser ruthenus</i>	6	0	3		
<i>Acipenser schrenckii</i>	6	0	4		
<i>Acipenser sinensis</i>	13	12	17		
<i>Acipenser stellatus</i>	13	11	17		
<i>Acipenser stellatus donensis</i>	9	12	17		
<i>Acipenser stellatus stellatus</i>	9	12	17		
<i>Acipenser sturio</i>	14	10	16		
<i>Acipenser transmontanus</i>	6	0	4		
<i>Polyodon spathula</i>	7	0	6		
<i>Psephurus gladius</i>	9	1	9		
<i>Scaphirhynchus albus</i>	7	1	9		
<i>Scaphirhynchus platorynchus</i>	7	1	9		
<i>Scaphirhynchus suttkusi</i>	7	1	9		

Species 17: Acipenser gueldenstaedtii

Full name: *Acipenser gueldenstaedtii* Brandt & Ratzeburg, 1833

For *Acipenser gueldenstaedtii* a species-specific assay for one haplotype of *A. gueldenstaedtii* is recommended. Note that this might not cover all haplotypes of *A. gueldenstaedtii*, and might also yield false positives from hybrids of *Acipenser*, including *A. baerii* (Table S17).

- Aci_gue_CR_F01: 5'-GAGCAGTCGTGAATGTTCCA-3',
- Aci_gue_CR_R01: 5'-CGAGAACCCCATCAACATT-3',
- Aci_gue_CR_P01: 5'-FAM-ATCTTTCTGAAGGAGCCCTGCATGT-BHQ-1-3',

Table S17: Species-specific primer/probe assay for *Acipenser gueldenstaedtii* with target gene, product size, melting temperature, primer/probe length, GC ratio (%), and number of mismatches between primer and probe region in closely related non-target species.

Species	Gene	Product size	Temp	Length	GC
<i>Acipenser gueldenstaedtii</i>	CR	112 bp			
Aci_gue_CR_F01		GAGCAGTCGTGAATGTTCCA	59.5	20	48
Aci_gue_CR_R01		CGAGAACCCCATCAACATT	59.0	20	48
Aci_gue_CR_P01		ATCTTTCTGAAGGAGCCCTGCATGT	65.0	26	38
Related species	Forward	Reverse	Probe		
<i>Acipenser baerii</i>	0	0	0		
<i>Acipenser naccarii</i>	5	1	4		
<i>Acipenser persicus</i>	4	2	2		
<i>Acipenser ruthenus</i>	2	2	7		
<i>Acipenser stellatus</i>	6	3	14		
<i>Huso huso</i>	4	1	4		

Species 18: Acipenser ruthenus

Full name: *Acipenser ruthenus* Linnaeus, 1758

For *Acipenser ruthenus* a species-specific assay for one haplotype of *A. ruthenus* is recommended (Table S18). Note that this might not cover all haplotypes of *A. ruthenus*, and might also yield false positives from hybrids of *Acipenser*.

- Aci_rut_CR_F01: 5'-TGCCGAGAACCATATCAACA-3',
- Aci_rut_CR_R01: 5'-GACCGTGAATGTTCCACCTT-3',
- Aci_rut_CR_P01: 5'-FAM-ATITGGGTTCATTCACTGACATGC-BHQ-1-3',

Table S18: Species-specific primer/probe assay for *Acipenser ruthenus* with target gene, product size, melting temperature, primer/probe length, GC ratio (%), and number of mismatches between primer and probe region in closely related non-target species.

Species	Gene	Product size		Temp	Length	GC
<i>Acipenser ruthenus</i>	CR	112 bp				
Aci_rut_CR_F01		TGCCGAGAACCATATCAACA		60.1	20	45
Aci_rut_CR_R01		GACCGTGAATGTTCCACCTT		59.8	20	50
Aci_rut_CR_P01		ATTGGGTTCATTCACTGACATGC		67.9	26	42
Related species		Forward	Reverse	Probe		
<i>Acipenser baerii</i>		2	5	3		
<i>Acipenser baerii baicalensis</i>		11	14	20		
<i>Acipenser baerii stenorhynchus</i>		11	13	20		
<i>Acipenser brevirostrum</i>		13	13	8		
<i>Acipenser dabryanus</i>		4	7	2		
<i>Acipenser fulvescens</i>		3	6	4		
<i>Acipenser gueldenstaedtii</i>		4	7	3		
<i>Acipenser gueldenstaedtii colchicus</i>		13	13	18		
<i>Acipenser gueldenstaedtii gueldenstaedtii</i>		13	13	19		
<i>Acipenser medirostris</i>		4	6	2		
<i>Acipenser mikadoi</i>		10	12	8		
<i>Acipenser naccarii</i>		4	7	3		
<i>Acipenser nudiventris</i>		4	5	2		
<i>Acipenser oxyrinchus desotoi</i>		12	13	7		
<i>Acipenser persicus</i>		3	7	3		
<i>Acipenser persicus persicus</i>		13	13	19		
<i>Acipenser ruthenus</i>		0	0	0		
<i>Acipenser schrenckii</i>		5	5	2		
<i>Acipenser sinensis</i>		4	7	2		
<i>Acipenserstellatus</i>		12	13	8		
<i>Acipenserstellatus donensis</i>		11	13	20		
<i>Acipenserstellatus ponticus</i>		12	12	20		
<i>Acipenserstellatusstellatus</i>		12	13	20		
<i>Acipensersturio</i>		9	14	8		
<i>Acipensertransmontanus</i>		4	5	2		
<i>Huso huso</i>		12	12	18		
<i>Huso huso caspicus</i>		12	12	18		
<i>Huso huso maeoticus</i>		12	12	18		
<i>Polyodon spathula</i>		4	9	5		
<i>Scaphirhynchus albus</i>		2	8	3		
<i>Scaphirhynchus platorynchus</i>		2	7	3		
<i>Scaphirhynchus suttkusi</i>		2	7	3		

Species 19: Acipenser stellatus

Full name: *Acipenser stellatus* Pallas, 1771

For *Acipenser stellatus* a species-specific assay for one haplotype of *A. stellatus* is recommended (Table S19). Note that this might not cover all haplotypes of *A. stellatus*, and might also yield false positives from hybrids of *Acipenser*.

- Aci_ste_CR_F03: 5'-GTGTGAGGGAGAAAAGACCA-3',
- Aci_ste_CR_R03: 5'-CACCACATACAAACCTGCGTTA-3',
- Aci_ste_CR_P03: 5'-FAM-GGCCTCGTGAAAACACTCCTCATGT-BHQ-1-3'

Table S19: Species-specific primer/probe assay for *Acipenser stellatus* with target gene, product size, melting temperature, primer/probe length, GC ratio (%), and number of mismatches between primer and probe region in closely related non-target species.

Species	Gene	Product size	Temp	Length	GC
<i>Acipenser stellatus stellatus</i>	CR	70 bp			
Aci_ste_CR_F03		GTGTGAGGGAGAAAAGACCA	60.1	20	50
Aci_ste_CR_R03		CACCACATACAAACCTGCGTTA	60.8	22	45
Aci_ste_CR_P03		GGCCTCGTGAAAACACTCCTCATGT	68.6	25	52
Related species	Forward	Reverse	Probe		
<i>Acipenser baerii</i>	8	13	3		
<i>Acipenser brevirostrum</i>	13	17	18		
<i>Acipenser fulvescens</i>	17	15	13		
<i>Acipenser fulvescens</i>	17	15	13		
<i>Acipenser gueldenstaedtii</i>	13	17	18		
<i>Acipenser medirostris</i>	17	16	17		
<i>Acipenser medirostris</i>	17	16	17		
<i>Acipenser mikadoi</i>	13	17	18		
<i>Acipenser naccarii</i>	17	15	17		
<i>Acipenser naccarii</i>	17	15	17		
<i>Acipenser naccarii</i>	17	15	17		
<i>Acipenser nudiventris</i>	12	16	19		
<i>Acipenser oxyrinchus</i>	17	16	15		
<i>Acipenser persicus</i>	17	15	15		
<i>Acipenser persicus persicus</i>	8	10	7		
<i>Acipenser ruthenus</i>	17	16	17		
<i>Acipenser schrenckii</i>	13	19	18		
<i>Acipenser sinensis</i>	13	17	19		
<i>Acipenser stellatus</i>	17	16	16		
<i>Acipenser stellatus stellatus</i>	0	0	0		
<i>Acipenser sturio</i>	13	17	18		
<i>Acipenser transmontanus</i>	13	17	18		
<i>Huso huso</i>	8	9	5		

Species 20: *Huso huso*

Full name: *Huso huso* (Linnaeus, 1758)

The genus *Huso* comprises two species *H. huso* (Linnaeus, 1758) and *H. dauricus* (Georgi, 1775). For *Huso huso* a species-specific assay is recommended (Table S20) that should be able to distinguish between *Huso* and *Acipenser*. Further validation tests would be required to test whether this system can distinguish between *H. huso* and *H. dauricus*. *Huso huso* is native to Eurasia, the Caspian Sea and Black Sea and *H. dauricus* is native to Eastern Asia, and the recommended system will likely detect both invasive species (i.e. *H. huso* and *H. dauricus*), but note that the system might be unable to distinguish between these two species.

- Hus_hus_Cytb_F01: 5'-TCCGATCCATCCCAAATAAA-3',
- Hus_hus_Cytb_R01: 5'-TTCCCCGTTGTTAGAGGTG-3',
- Hus_hus_Cytb_P01: 5'-FAM-GGCGGGGTACTAGCCCTCTATTCTC-BHQ1-3',

Table S20: Species-specific primer/probe assay for *Huso huso* with target gene, product size, melting temperature, primer/probe length, GC ratio (%), and number of mismatches between primer and probe region in closely related non-target species.

Species	Gene	Product size	Temp	Length	GC
<i>Huso huso</i>	Cytb	102 bp			
Hus_hus_Cytb_F01	TCCGATCCATCCCAAATAAA		60.1	20	40
Hus_hus_Cytb_R01	TTCCCCGTTGTTAGAGGTG		60.0	20	50
Hus_hus_Cytb_P01	GGCGGGGTACTAGCCCTCTATTCTC		67.8	26	58
Related species	Forward	Reverse	Probe		
<i>Acipenser baerii</i>	3	0	2		
<i>Acipenser brevirostrum</i>	2	1	2		
<i>Acipenser dabryanus</i>	2	0	1		
<i>Acipenser dabryanus</i>	2	0	1		
<i>Acipenser fulvescens</i>	3	0	2		
<i>Acipenser gueldenstaedtii</i>	3	0	2		
<i>Acipenser medirostris</i>	1	1	1		
<i>Acipenser mikadoi</i>	1	1	0		
<i>Acipenser naccarii</i>	3	0	2		
<i>Acipenser nudiventris</i>	2	0	3		
<i>Acipenser oxyrinchus</i>	3	1	1		
<i>Acipenser persicus</i>	3	0	2		
<i>Acipenser ruthenus</i>	3	1	4		
<i>Acipenser schrenckii</i>	1	0	1		
<i>Acipenser sinensis</i>	2	0	1		
<i>Acipenser stellatus</i>	2	1	3		
<i>Acipenser sturio</i>	3	2	1		
<i>Acipenser transmontanus</i>	2	0	2		
<i>Huso huso</i>	0	0	0		
<i>Polyodon spathula</i>	3	1	5		
<i>Psephurus gladius</i>	4	0	3		

Species 21: Oncorhynchus gorbuscha

Full name: *Oncorhynchus gorbuscha* (Walbaum, 1861)

The genus *Oncorhynchus* belongs to the family Salmonidae and contains 15 species (Froese & Pauly 2015).

The following primers and probes are designed to target *Oncorhynchus gorbuscha*:

- Onc_Gor_Cytb_F01: 5'-CAGGAGCATCCGTCGACTT-3'
- Onc_Gor_Cytb_R01: 5'-CTGAGAGATTGCCGGTGGTT-3'
- Onc_Gor_Cytb_P01: 5'-FAM-GCTGGAATCTCATCAATTAGGGGCCA-BHQ-1-3'

Both *O. gorbuscha* and its sister-species are well represented by sequence material. Only for the single sister-species *O. rhodurus* is sequence not available. However, according Froese and Pauly (2015) this species is endemic to lake Biwa, Chûzenji and Ashinoko in Japan and is therefore not considered to pose a threat of false positives. Additional to the BLAST search of the assay, two native Danish salmonid species were included in the comparison to strengthen the precision of the assay, see Table S21.

Table S21: Species-specific primer/probe assay for *Oncorhynchus gorbuscha* with target gene, product size, melting temperature, primer/probe length, GC ratio (%), and number of mismatches between primer and probe region in closely related non-target species.

Species	Gene	Product size	Temp	Length	GC
<i>Oncorhynchus mykiss</i>	CytB	119 bp			
Onc_Gor_Cytb_F01	CAGGAGCATCCGTCGACTT		59.5	19	58
Onc_Gor_Cytb_R01	CTGAGAGATTGCCGGTGGTT		60.0	20	55
Onc_Gor_Cytb_P01	GCTGGAATCTCATCAATTAGGGGCCA		65.3	28	46
Related species	F01	R01	P01		
<i>Oncorhynchus mykiss</i>	4	3	2		
<i>Oncorhynchus mykiss</i> x					
<i>Salmo salar</i> _KP218514	4	3	2		
<i>Oncorhynchus mykiss</i> x					
<i>Salmo salar</i> _NC_026537	4	3	2		
<i>Oncorhynchus gilae</i>	4	3	3		
<i>Oncorhynchus chrysoaster</i>	4	3	2		
<i>Oncorhynchus keta</i>	2	4	2		
<i>Oncorhynchus clarkii</i>	4	3	4		
<i>Oncorhynchus nerka</i>	4	4	3		
<i>Oncorhynchus tshawytscha</i>	5	2	4		
<i>Oncorhynchus kisutch</i>	5	4	4		
<i>Oncorhynchus masoni</i>	4	5	3		
<i>Salmo trutta</i>	2	5	4		
<i>Salmo salar</i>	2	3	5		

Species 22: Salvinus fontinalis

Full name: *Salvinus fontinalis* Mitchill, 1814.

The genus *Salvinus* belongs to the family Salmonidae and is comprised of nine marine species (WoRMS 2015). Sequences for target and sister-species were well represented in the NCBI database.

An assay for *S. fontinalis* has already been published and validated by Jane *et al.* (2014) based on the cytochrome b region. The assay utilizes a TaqMan MGB probe:

- Sal_For_Cytb_F01: 5'-CCATGAGGGCAAATATCCTTCTGA-3'
- Sal_For_Cytb_R01: 5'-TCATTGTACAAGGGCACCTCCTA-3'
- Sal_For_Cytb_P01: 5'-FAM-CTCCTCTCTGCTGTACCC-MGB-NFQ-3'

The annealing temperature utilized for this assay is 60 °C, whereas the values in table S22.1 are the temperatures calculated in Geneious version 9.0.4. Due to a low number of mismatches between *S. alpinus* and *S. malma* and the utilization of a TaqMan MGB probe, a secondary assay was also designed to improve precision.

Table S22.1: Species-specific primer/probe assay for *Salvinus fontinalis* with target gene, product size, melting temperature, primer/probe length, GC ratio (%), and number of mismatches between primer and probe region in closely related non-target species.

Species	Gene	Product size	Temp	Length	GC
<i>Salvinus fontinalis</i>	Cytb	90bp			
Sal_For_Cytb_F01		CCATGAGGGCAAATATCCTTCTGA	60.4	24	45.8
Sal_For_Cytb_R01		TCATTGTACAAGGGCACCTCCTA	61.1	23	47.8
Sal_For_Cytb_P01		CTCCTCTCTGCTGTACCC*	55.7	18	61.1

*TaqMan MGB probe

Related species	F01	R01	P01
<i>Salvelinus albus</i>	2	0	3
<i>Salvelinus malma</i>	1	0	3
<i>Salvelinus taranetzii</i>	2	0	3
<i>Salvelinus alpinus</i>	0	0	3
<i>Salvelinus confluentus</i>	2	0	3
<i>Salvelinus namaycush</i>	2	1	4
<i>Salvelinus leucomaenis</i>	1	1	4
<i>Salvelinus malma malma</i> ×	1	0	4
<i>Salvelinus leucomaenis</i>			
<i>Salvelinus levanidovi</i>	4	1	4

*MGB probe

The following primers and probe are designed to target *Salvelinus fontinalis*:

- Sal_For_Cytb_F01: 5'-AGGGCAAATATCCTTCTGAGGG-3'
- Sal_For_Cytb_R01: 5'-GGCACCTCCTACATAGGGTAC-3'
- Sal_For_Cytb_P01: 5'-FAM-GCCACTGTAATTACAAACCTCCTCTGC-BHQ-1-3'

Table S22.2: Species-specific primer/probe assay for *Salvinus fontinalis* with target gene, product size, melting temperature, primer/probe length, GC ratio (%), and number of mismatches between primer and probe region in closely related non-target species.

Species	Gene	Product size	Temp	Length	GC
<i>Salvinus fontinalis</i>	Cytb	73 bp			
Sal_For_Cytb_F02		AGGGCAAATATCCTTCTGAGGG	59.6	22	50
Sal_For_Cytb_R02		GGCACCTCCTACATAGGGTAC	59	21	57.1
Sal_For_Cytb_P02		GCCACTGTAATTACAAACCTCCTCTGC	65.5	29	48.3
Related species	F02	R02	P02		
<i>Salvelinus albus</i>	3	2	3		
<i>Salvelinus malma</i>	2	2	3		
<i>Salvelinus tanganetzi</i>	3	2	3		
<i>Salvelinus alpinus</i>	2	2	3		
<i>Salvelinus confluentus</i>	3	2	3		
<i>Salvelinus namaycush</i>	3	2	3		
<i>Salvelinus leucomaenis</i>	3	2	3		
<i>Salvelinus malma malma</i> ×	2	3	2		
<i>Salvelinus leucomaenis</i>					
<i>Salvelinus levanidovii</i>	5	3	2		

Species 23: Colpomenia peregrine

Full name: *Colpomenia peregrine* (Sauvageau) Hamel 1937

The genus *Colpomenia* contains 14 species and both *rbcL* and *psaA* genes have been sequenced, however not the *matK* gene. The *psaA* gene does not show strong representation among the species in the genus and could therefore not be used for the design of a species specific assay for *Colpomenia peregrine*. The *rbcL* gene on the other hand has been more widely sequenced, however was found not useful for assay design. In addition, the COX3 gene was represented at a significant scale. After looking through the three genes COX3 was identified as the only gene where it was possible to design an assay containing enough differentiation in primer/probe sequences to non-target species, as seen in table S23:

- Col_per_COX3_F01: 5' GCAAGCTTGTGAATATGCTAATG-3'
- Col_per_COX3_R01: 5'-CAGCTAAAAATATTGTACCGATTA-3'
- Col_per_COX3_P01: 5'-FAM-TTCAGTTTTACATGGCTACAGGCITC-BHQ-1-3'

Table S23: Species specific primer/probe assay for *Colpomenia peregrina* with, target gene, product size, melting temperature, primer/probe length, GC ratio (%), and number of mismatches between primer and probe region in closely related non-target species

Species	Gene	Product size	Temp	Length	GC
<i>Colpomenia peregrina</i>	COX3	116 bp			
Col_per_COX3_F01		GCAAGCTTGTGAATATGCTAATG	57.6	23	35
Col_per_COX3_R01		CAGCTAAAAATATTGTACCGATTA	56.6	24	29
Col_per_COX3_P01		TTCAGTTTTACATGGCTACAGGCITC	65.6	28	39
Related species*		Forward	Reverse	Probe	
<i>Colpomenia bullosa</i>		7	6	4	
<i>Colpomenia claytoniae</i>		1	3	4	
<i>Colpomenia durvillei</i>		7	3	4	
<i>Colpomenia ecuticulata</i>		6	5	5	
<i>Colpomenia expansa</i>		1	6	5	
<i>Colpomenia phaeodactyla</i>		8	4	4	
<i>Colpomenia ramosa</i>		7	5	7	
<i>Colpomenia sinuosa</i>		4	6	6	
<i>Colpomenia tuberculata</i>		7	5	7	
<i>Colpomenia sp. CCC-2009a</i>		7	5	4	
<i>Colpomenia sp. KML-2014</i>		8	4	4	
<i>Petalonia zosterifolia</i>		7	5	2	
<i>Scytosiphon lomentaria</i>		6	5	6	
<i>Scytosiphon sp. CCC-2009a</i>		6	5	6	
<i>Desmarestia menziesii</i>		9	5	4	
<i>Padina pavonica</i>		8	7	5	

* not necessarily related but show strong sequence similarity to target species

Species 24: Ensis directus

Full name: *Ensis directus* (Conrad, 1843)

Ensis americanus (Gould, 1870) is considered a junior synonym of *E. directus* (Cosal 2009). The genus *Ensis* comprises around 30 species (WoRMS Editorial Board 2015) and on the NCBI Genbank nucleotide sequences are available from nine species of *Ensis*, including *E. directus*. Since the NCBI GenBank database does not have all known species of *Ensis* represented, false positives from closely related non-target species is a risk. The primers and probe suggested here will only be able to differentiate between species of *Ensis* that are represented by nucleotide sequences on the NCBI Genbank database.

A population study on *E. directus* and *E. terranovensis* Vierna & Martínez-Lage, 2012 found subtle differences in the genetic variation (Vierna *et al.* 2012).

For detection of eDNA from cytochrome oxidase 1 in the mitochondrial genome from *Ensis directus* the following primers and probes are recommended (Table S24.1 and Table S24.2):

- Ens_dir_CO1_F01: 5'- GTTGTTGGTAGCGGGATTG -3',
- Ens_dir_CO1_R01: 5'- TCCCATAATCCATTGAAGGAC -3',
- Ens_dir_CO1_P01: 5'-FAM- GCCGGATGAACCTATTATCCGCCTTA -BHQ-1-3'

Table S24.1: Species specific primer/probe assay for *E. directus* with, target gene, product size, melting temperature, primer/probe length, GC ratio (%), and number of mismatches between primer and probe region in closely related non-target species.

Species	Gene	Product size	Temp	Length	GC
<i>Ensis directus</i>	mtDNA-CO1	92 bp			
Ens_dir_CO1_F01	GTTGTTGGTAGCGGGATTG		60.4	20	50
Ens_dir_CO1_R01	TCCCATAATCCATTGAAGGAC		58.7	21	43
Ens_dir_CO1_P01	GCCGGATGAACCTATTATCCGCCTTA		68.3	27	44
Related species	Forward	Reverse	Probe		
<i>Ensis ensis</i>	4	3	3		
<i>Ensis goreensis</i>	4	4	5		
<i>Ensis macha</i>	3	1	3		
<i>Ensis magnus</i>	3	3	5		
<i>Ensis minor</i>	3	2	4		
<i>Ensis minor</i>	3	2	4		
<i>Ensis silique</i>	3	4	2		
<i>Ensis terranovensis</i>	2	0	2		
<i>Phaxas pellucidus</i>	4	3	2		

An additional system for detection of eDNA from cytochrome oxidase 1 in the mitochondrial genome from *Ensis directus* is also recommended. In case the first one is unspecific:

- Ens_dir_CO1_F02: 5'- CATTCCCTCGATTAAACAATTGTA-3',
- Ens_dir_CO1_R02: 5'- CAATCCCGCTACCAACAAAC -3',
- Ens_dir_CO1_P02: 5'-FAM- TTGACTTITACCTTGTGCTTGTITTT-BHQ-1-3'

Table S24.2: Species specific primer/probe assay for *E. directus* with, target gene, product size, melting

temperature, primer/probe length, GC ratio (%), and number of mismatches between primer and probe region in closely related non-target species.

Species	Gene	Product size	Temp	Length	GC
<i>Ensis directus</i>	mtDNA-CO1	90 bp			
Ens_dir_CO1_F02	CATTTCCTCGATTAAACAATTTGA		59.4	24	29
Ens_dir_CO1_R02	CAATCCCGCTACCAACAAAC		60.4	20	50
Ens_dir_CO1_P02	TTGACTTTTACCTTGTGCTTTGTTTT		62.4	27	30
<hr/>					
Related species	Forward	Reverse	Probe		
<i>Ensis ensis</i>	5	4	4		
<i>Ensis goreensis</i>	5	4	7		
<i>Ensis macha</i>	5	3	4		
<i>Ensis magnus</i>	4	3	8		
<i>Ensis minor</i>	2	3	1		
<i>Ensis minor</i>	2	3	1		
<i>Ensis siliqua</i>	4	3	7		
<i>Ensis terranovensis</i>	1	2	2		
<i>Phaxas pellucidus</i>	3	4	4		

These two eDNA detection systems targeting *E. directus* might not be able to differentiate between *E. directus* and *Ensis terranovensis*. But as both *E. directus* and *Ensis terranovensis* are native inhabitants in the North Western Atlantic and both are considered introduced in the North Eastern Atlantic, this inability to differentiate between these two closely related species might be considered irrelevant for the purpose of monitoring the invasive *E. directus*-*E. terranovensis* species complex in European seas.

Species 25: *Dasya baillouviana*

Full name: *Dasya baillouviana* (Gmelin) Montagne, 1841

The genus *Dasya* represents 19 different species. For 12 of these species the *rbcL* gene has been sequenced, and due to the indicated species specificity of this gene for eDNA assay design for aquatic plants (Scriven *et al.* 2015), of available sequences genes for *Dasya baillouviana* and relevant non-target species, this gene was selected. For better assay design a subset of various species showing sequence similarity through a BLAST search were included in the sequence alignment (Table S25). Assay design in *rbcL* was not trivial and base pair mismatch to non-target species was maybe not as high as one could desire. Despite this a species specific eDNA primer/probe assay was designed for *Dasya baillouviana*:

- Das_bai_rbcL_F01: 5'-CAAACITTAAC TGCA TCTATC ATC-3'
- Das_bai_rbcL_R01: 5'-AATCCTGTTGCAGGACCTTG-3'
- Das_bai_rbcL_P01: 5'-FAM-CGTATCCCAGTAGCTTATCTAAAAACATT-BHQ-1-3'

Table S25: Species specific primer/probe assay for *Dasya baillouviana* with, target gene, product size, melting temperature, primer/probe length, GC ratio (%), and number of mismatches between primer and probe region in closely related non-target species

Species	Gene	Product size	Temp	Length	GC
<i>Dasya baillouviana</i>	<i>rbcL</i>	113 bp			
Das_bai_rbcL_F01		CAAACITTAAC TGCA TCTATC ATC	57.6	23	35
Das_bai_rbcL_R01		AATCCTGTTGCAGGACCTTG	58.4	20	50
Das_bai_rbcL_P01		CGTATCCCAGTAGCTTATCTAAAAACATT	64.6	29	34
Related species*	Forward	Reverse	Probe		
<i>Dasya anastomosans</i>	4	1	5		
<i>Dasya collabens</i>	4	2	3		
<i>Dasya elongata</i>	4	1	6		
<i>Dasya enomotoi</i>	2	3	4		
<i>Dasya hutchinsiae</i>	4	2	4		
<i>Dasya iridescens</i>	2	3	4		
<i>Dasya ocellata</i>	3	1	2		
<i>Dasya sessilis</i>	1	0	4		
<i>Dasya</i> sp. GENT HV TZ13	4	1	5		
<i>Dasya</i> sp. KS-2014	5	2	2		
<i>Dasya</i> sp. L SGAD0712164	4	2	4		
<i>Dasyaceae</i> sp. GH04	6	2	1		
<i>Heterosiphonia japonica</i>	4	2	3		
<i>Dipterocladia pinnatifolia</i>	4	2	6		
<i>Eupogodon planus</i>	3	2	3		
<i>Rhodoptilum plumosum</i>	3	0	1		
<i>Zellera tanallina</i>	4	1	4		
<i>Chauviella coriifolia</i>	4	2	3		
<i>Branchioglossum woodii</i>	4	2	3		
<i>Branchioglossum bipinnatifidum</i>	5	3	3		
<i>Grinnellia americana</i>	5	2	2		
<i>Chauviella coriifolia</i>	5	2	4		
<i>Phitymorphora amansioides</i>	5	3	4		
<i>Phitymorphora linearis</i>	6	3	3		
<i>Hypoglossum hypoglossoides</i>	4	3	4		
<i>Hypoglossum</i> sp. 1	4	2	4		
<i>Polyneuropsis stolonifera</i>	5	4	3		
<i>Paraglossum crassinervium</i>	4	2	2		
<i>Bartoniella crenata</i>	6	2	1		

* not necessarily related but show strong sequence similarity to target species

Species 26: Heterosiphonia japonica

Full name: *Heterosiphonia japonica* Yendo, 1920

The genus *Heterosiphonia* comprises 10 species. Despite relatively good representation of genes for assay design for *Heterosiphonia japonica* not all species within the genus were represented in all genes. Best representation and assay design was obtained using the *rbcL* gene:

- Het_jap_rbcL_F01: 5'- CAACAGGTATTATTGTAGAACGC-3'
- Het_jap_rbcL_R01: 5'-TAATTITACCAAGATAGACCTAGC-3'
- Het_jap_rbcL_P01: 5'-FAM-ATAAGTTGGCCGTCCTTCCTAGGA-BHQ-1-3'

Mismatch to the genus *Dasya*, within the family *Dayaceae* (within which the genus *Heterosiphonia* also belongs) was for *Dasya collabens* not possible to design, whereas another species on this list, *Dasya baillouiana*, did show any mismatches in the sequence regions of the assay to *Heterosiphonia japonica* (Table S26).

Table S26: Species specific primer/probe assay for *Heterosiphonia japonica* with, target gene, product size, melting temperature, primer/probe length, GC ratio (%), and number of mismatches between primer and probe region in closely related non-target species. Problematic species are marked in bold

Species	Gene	Product size	Temp	Length	GC
<i>Heterosiphonia japonica</i>	<i>rbcL</i>	99 bp			
Het_jap_rbcL_F01		CAACAGGTATTATTGTAGAACGC	59.2	23	39
Het_jap_rbcL_R01		TAATTITACCAAGATAGACCTAGC	58.3	24	33
Het_jap_rbcL_P01		ATAAGTTGGCCGTCCTTCCTAGGA	66.2	26	46

Related species*	Forward	Reverse	Probe
<i>Heterosiphonia gibbesii</i>	4	3	8
<i>Heterosiphonia plumosa</i>	3	4	10
<i>Heterosiphonia pulchra</i>	3	5	8
<i>Nannochloropsis gaditana</i>	6	4	7
<i>Achnanthidium saprophilum</i>	3	4	12
<i>Nitzschia amphibia</i>	1	4	11
<i>Coscinodiscus radiatus</i>	3	4	9
<i>Polysiphonia virgata</i>	5	5	8
<i>Polysiphonia denudata</i>	4	5	7
<i>Aulacoseira granulata</i>	5	4	9
<i>Rhodella violacea</i>	7	4	6
<i>Caloglossa rotundata</i>	4	4	4
<i>Eupogonon spinellus</i>	2	3	4
<i>Eupogonon planus</i>	2	5	3
<i>Rhodymenia corallina</i>	5	4	3
<i>Dasya sp. L SGAD0712164</i>	1	2	2
<i>Dasya collabens</i>	0	0	0
<i>Dasya baillouiana</i>	5	4	5

* not necessarily related but show strong sequence similarity to target species

Species 27: Spartina anglica

Full name: *Spartina anglica*, C. E. Hubbard

According to the NCBI database the genus *Spartina* consists of 17 species. Within the genus sequence information was available for the *matK* gene. However, alignment of the sequences revealed insufficient numbers of mismatches among the species, and it was therefore not possible to design a useful eDNA primer/probe assay for *Spartina anglica* using sequence information from this gene. Alignments of available *rbcL* sequence were all almost identical and therefore also not usable for assay design. Having tested the more recognised genes for eDNA primer/probe design (Scriven *et al.* 2015) alignment of available sequences for the *trnT-trnL* and *trnL-trnF* intergenic spacer region was tried, however also unsuccessful.

According to the UniProt consortium (The UniProt Consortium 2007) the species *Spartina anglica* is a synonym of *Sporobolus anglicus* (C.E. Hubb.) P.M. Peterson & Saarela. However, despite extending the search to include information for genes sequenced for *Sporobolus anglicus* (a genus containing 157 species) it was not possible to design an eDNA primer/probe assay for either of the two synonyms.

Species 28: Heterosigma akashiwo

Full name: *Heterosigma akashiwo* (Y. Hada) Y. Hada ex Y. Hara & M. Chihara, 1987

The genus *Heterosigma* comprises only this one species. The genus *Heterosigma* belongs to the family *Chattonellaceae* containing 18 species and variates of these.

An assay for *Heterosigma akashiwo* has already been published based on the 18S sequence (Coyne *et al.* 2005). However, this assay has a product size of 376bp. Due to this a new assay was designed based on available sequence information for the *rbcL* gene obtained from the NCBI online database:

- Het_aka_rbcL_F01: 5'-CTCTTGAAGCTATGATTAGCTA-3'
- Het_aka_rbcL_R01: 5'-CACATAATTAGCGATGTCTCTT-3'
- Het_aka_rbcL_P01: 5'-FAM-CGAAGGTCGTGACTACGTITCTGAAG-BHQ-1-3'

Species specificity and assay details can be seen in table S28. Target species was compared to most related species but also species showing sequence similarity identified by blasting the *rbcL* sequence for *Heterosigma akashiwo*. *Olisthodiscus luteus* (N. Carter) and *Fibrocapsa cf. japonica* (JX067601.1) were found to have identical sequence for *rbcL* as *Heterosigma akashiwo*. According to AlgaeBase.org *Olisthodiscus luteus* and *Heterosigma akashiwo* are most likely two different species. The sequence identity may therefore be caused by wrong species identification and registration within the NCBI online database for the gene. This could also be the reason why *Fibrocapsa cf. japonica* (AB280606.1) and *Heterosigma akashiwo* have identical sequences. The sequence available for *Chattonella minima* was not of equal length compared to the other species in the alignment and an alignment between this species and *Heterosigma akashiwo* could therefore not be done. However, comparison among the other *Chattonella* species in the alignment it is very likely that *Chattonella minima* would show the same pattern of mismatches as the others if sequenced at equal length. Because of this it is believed that the designed assay will not amplify for *Chattonella minima*.

Table S28: Species specific primer/probe assay for *Heterosigma akashiwo* with, target gene, product size, melting temperature, primer/probe length, GC ratio (%), and number of mismatches between primer and probe region in closely related non-target species. Problematic species are marked in bold

Species	Gene	Product size	Temp	Length	GC
<i>Heterosigma akashiwo</i>	<i>rbcL</i>	99 bp			
Het_aka_rbcL_F01		CTCTTGAAGCTATGATTAGCTA	58.3	24	33
Het_aka_rbcL_R01		CACATAATTAGCGATGTCTCTT	57.6	23	35
Het_aka_rbcL_P01		CGAAGGTCGTGACTACGTITCTGAAG	67.9	26	50
Related species*	F01	R01	P01		
<i>Olisthodiscus luteus</i>	0	0	0		
<i>Chattonella marian</i>	7	4	5		
<i>C. minima</i>	-	-	-		
<i>C. subsalsa</i>	7	4	3		
<i>C. orata</i>	7	4	5		
<i>C. antiqua</i>	7	4	5		
<i>Fibrocapsa japonica</i>	2	5	5		
<i>Haramonas dimorpha</i>	7	4	5		
<i>Ophiocytium capitatum</i>	5	9	6		
<i>Ophiocytium majus</i>	4	8	5		
<i>Tribonema regulare</i>	5	8	5		
<i>Tribonema viride</i>	5	8	5		
<i>Tribonema intermixum</i>	5	8	5		
<i>Botrydium becharianum</i>	4	8	5		
<i>Bumilleria klebsiana</i>	5	8	5		
<i>Xanthonema muciculum</i>	7	8	4		

* not necessarily related but show strong sequence similarity to target species

Species 29: Pseudochattonella farcimen

Full name: *Pseudochattonella farcimen* (W. Eikrem, B. Edvardsen & J. Throndsen) W. Eichrem, 2009

The genus *Pseudochattonella* only contains the two species *Pseudochattonella farcimen* and *Pseudochattonella verruculosa*. The two species are more or less impossible to tell apart morphologically after fixation in lygol. Due to this a molecular method has been developed and to date serve as the best solution to tell apart the two species (Dittami *et al.* 2013). However, the designed primers (best possible) only have two mismatches each, respectively. According to the published qPCR protocol it is relatively difficult even with the molecular method to determine if the amplification in the qPCR reaction is caused by the presence of the one or the other species (Dittami *et al.* 2013):

- PfarF: 5'-GAGGTCCCGGGGAGAAATT-3'
- PfarR: 5'-GCAACTCGACTCCACTAGG-3'

No other mismatches exist within the sequence between the two primers. Therefore, no probe was designed. However, it has come to our attention that a primer/probe assay is under development and awaiting publication (Eckford-Soper & Daugbjerg (in press)). With this new assay it is expected that differentiation between the two *Pseudochattonella* species will be possible.

Table S29: Species specific primer/probe assay for *Pseudochattonella farcimen* with, target gene, product size, melting temperature, primer/probe length, and GC ratio (%)

Species	Gene	Product size	Temp	Length	GC
<i>Pseudochattonella farcimen</i>	28S	91 bp			
PfarF		GAGGTCCCGGGGAGAAATT	59.5	19	58
PfarR		GCAACTCGACTCCACTAGG	59.5	19	58

Species 30: Molgula manhattensis

Full name: *Molgula manhattensis* (De Kay, 1843)

According to WoRMS the genus *Molgula* comprises 143 marine species. The genus belongs to the family Molgulidae, which contains 15 genera (WoRMS 2015).

The most suitable genetic region was found to be 18S with the broadest interspecies coverage, representing 13 out of 143 species including the target species *M. manhattensis*. Based on the sequence data, there is a high level of species similarity, e.g. a pairwise alignment between *M. manhattensis* and *M. retortiformis* shows that only one nucleotide insertion separates the two species in a 964 nucleotide fragment. The limited interspecies variation and limited sequence material from sister-species leads to uncertainty with regards to assay specificity, depending on local species occurrences.

The following primers and probe are designed to target *M. manhattensis*:

- Mol_Man_18S_F01: 5'-CAAGGTGTGTACTGTCGGCC-3'
- Mol_Man_18S_R01: 5'-AAAGTAAACGCCTGGCCAC-3'
- Mol_Man_18S_P01: 5'-FAM-TCCGGCGGTTCTCTTAAC TGAGTGTCG-BHQ-1-3'

Table S30: Species-specific primer/probe assay for *Molgula manhattensis* with target gene, product size, melting temperature, primer/probe length, GC ratio (%), and number of mismatches between primer and probe region in closely related non-target species.

Species	Gene	Product size	Temp	Length	GC
<i>Molgula manhattensis</i>	18S	90 bp			
Mol_Man_18S_F01	CAAGGTGTGTACTGTCGGCC		60.9	20	60
Mol_Man_18S_R01	AAAGTAAACGCCTGGCCAC		59.3	20	50
Mol_Man_18S_P01	TCCGGCGGTTCTCTTAAC TGAGTGTCG		68.0	27	56
Related species	F01	R01	P01		
<i>Molgula retortiformis</i>	0	0	0		
<i>Molgula arenata</i>	0	0	2		
<i>Molgula provisionalis</i>	0	0	2		
<i>Molgula socialis</i>	0	0	2		
<i>Molgula complanata</i>	3	5	5		
<i>Molgula tectiformis</i>	4	6	4		
<i>Molgula occidentalis</i>	2	2	4		
<i>Molgula bleizi</i>	5	4	4		
<i>Molgula pugetensis</i>	5	3	4		
<i>Molgula oculata</i>	4	4	4		
<i>Molgula pacifica</i>	4	4	4		
<i>Molgula citrina</i>	5	4	5		

Species 31: Cercopagis pengoi

Full name: *Cercopagis pengoi* (Ostroumov, 1891)

Cercopagis pengoi is the only species within the genus *Cercopagis*. Within the Family *Cercopagidae* only one other genus (*Bythotrephes*) exists. Due to this, the sequence alignment search was extended to the infraorder *Onychopoda*. Within this infraorder all species sequenced for the CO1 gene were compiled for the design of the species specific eDNA primer/probe assay for *Cercopagis pengoi*. The assay was easily constructed and showed substantial mismatch to non-target species (table S31):

- Cer_pen_CO1_F01: 5'-ATTTCTTCATGGTAATGCCAGTA-3'
- Cer_pen_CO1_R01: 5'-GGGAAGGAATCAGAAACTAAGA-3'
- Cer_pen_CO1_P01: 5'-FAM-AATTGACTTGTCCCTCTGATGCTAGG-BHQ-1-3'

Table S31: Species specific primer/probe assay for *Cercopagis pengoi* with, target gene, product size, melting temperature, primer/probe length, GC ratio (%), and number of mismatches between primer and probe region in closely related non-target species

Species	Gene	Product size	Temp	Length	GC
<i>Cercopagis pengoi</i>	CO1	122 bp			
Cer_pen_CO1_F01		ATTTCTTCATGGTAATGCCAGTA	58.3	24	33
Cer_pen_CO1_R01		GGGAAGGAATCAGAAACTAAGA	58.4	22	41
Cer_pen_CO1_P01		AATTGACTTGTCCCTCTGATGCTAGG	66.2	26	46

Related species*	Forward	Reverse	Probe
<i>Bythotrephes cederstroemi</i>	9	3	6
<i>Bythotrephes longimanus</i>	8	3	6
<i>Bythotrephes</i> sp. BOLD:AAB9254	9	3	6
<i>Cornigerius maeoticus</i>	4	4	4
<i>Eavadne nordmanni</i>	4	7	7
<i>Eavadne spinifera</i>	3	3	7
<i>Pleopis polyphemoides</i>	4	6	7
<i>Pleopis</i> sp. BOLD:AAI2700	3	6	8
<i>Podon intermedius</i>	5	7	7
<i>Podon leuckartii</i>	7	4	7
<i>Podon eavadne angusta</i>	5	5	5
<i>Podon eavadne camptonyx</i>	5	5	5
<i>Podon eavadne trigona</i>	5	5	5
<i>Pseudeavadne tergestina</i>	3	5	6
<i>Polyphemus pediculus</i>	8	5	8
<i>Polyphemus</i> aff. <i>pediculus</i> 3 JRD-2015	6	5	7
<i>Polyphemus pediculus complex</i> sp. NA2	8	5	9
<i>Polyphemus pediculus complex</i> sp. NA3	6	5	7

* not necessarily related but show strong sequence similarity to target species

Species 32: Homarus americanus

Full name: *Homarus americanus*. Milne Edwards, 1837.

The genus *Homarus* comprises two marine species, *H. americanus* and *H. gammarus*. The genus belongs to the family Nephropidae which contains 16 genera with marine species (WoRMS 2015).

Both *H. americanus* and the sister-species *H. gammarus* are represented by sequence data. Additionally to the BLAST search of the assay, other decapod species were included in the alignment to improve and test the precision of the assay, see Table S32.

The following primers and probe are designed to target *H. americanus*:

- Hom_Ame_Cytb_F01: 5'-CCCTACCTTCTTGGAGACCC-3'
- Hom_Ame_Cytb_R01: 5'-CAGGTTGAATATGTGCTGGTGT-3'
- Hom_Ame_Cytb_P01: 5'-FAM-ACTTGTCGGCTAATCCACTCGT-BHQ-1-3'

Table S32: Species-specific primer/probe assay for *Homarus americanus* with target gene, product size, melting temperature, primer/probe length, GC ratio (%), and number of mismatches between primer and probe region in closely related non-target species.

Species	Gene	Product size	Temp	Length	GC
<i>Homarus americanus</i>	Cytb	73 bp			
Hom_Ame_Cytb_F01	CCCTACCTTCTTGGAGACCC		58.8	20	60
Hom_Ame_Cytb_R01	CAGGTTGAATATGTGCTGGTGT		58.9	22	46
Hom_Ame_Cytb_P01	ACTTGTCGGCTAATCCACTCGT		66	25	52
Related species	F01	R01	P01		
<i>Homarus gammarus</i>	2	1	6		
<i>Nephrops norvegicus</i>	3	2	7		
<i>Eriocheir sinensis</i>	5	6	9		
<i>Eriocheir japonica</i>	4	7	9		
<i>Eriocheir hepensis</i>	4	5	9		
<i>Paralithodes camtschaticus</i>	7	2	6		

Species 33: Paralithodes camtschaticus

Full name: *Paralithodes camtschaticus*Tilesius, 1815.

The genus *Paralithodes* belongs to the family Lithodidae which contains ten genera. The genus *Paralithodes* has five marine species (WoRMS 2015).

Paralithodes camtschaticus sequence was compared to that of its most closely related species and to common native species and other non-native decapod species (following Muus *et al.* 2006). Sequence data was available from the NCBI database for three of the five congeners; no sequence was found for *P. californiensis* and *P. rathbuni*. CO1 was found to be the most suitable genetic region with the broadest interspecies coverage.

The following primers and probe are designed to target *P. camtschaticus*:

- Par_Cam_CO1_F01: 5'-GCAGGAGCATCAGTGGATT-3'
- Par_Cam_CO1_R01: 5'-AGGTTATTCCTGTGGACGTATAT-3'
- Par_Cam_CO1_P01: 5'-FAM-TGGCTGGAGTATCTCTATTAGGGCT-BHQ-1-3'

Table S33: Species-specific primer/probe assay for *Paralithodes camtschaticus* with target gene, product size, melting temperature, primer/probe length, GC ratio (%), and number of mismatches between primer and probe region in closely related non-target species.

Species	Gene	Product size	Temp	Length	GC
<i>Paralithodes camtschaticus</i>	CO1	118bp			
Par_Cam_CO1_F01	GCAGGAGCATCAGTGGATT		58.2	20	50
Par_Cam_CO1_R01	AGGTTATTCCTGTGGACGTATAT		57.3	24	37.5
Par_Cam_CO1_P01	TGGCTGGAGTATCTCTATTAGGGCT		65.0	29	44.8
Related species	F01	R01	P01		
<i>Paralithodes platypus</i>	1	5	3		
<i>Paralithodes brevipes</i>	3	3	5		
<i>Maja squinado</i>	4	8	6		
<i>Nephrops norvegicus</i>	6	6	6		
<i>Homarus americanus</i>	5	7	6		
<i>Homarus gammarus</i>	5	7	6		
<i>Palinurus mauritanicus</i>	5	6	7		

Species 34: Didemnum vexillum

Full name: *Didemnum vexillum* Romanow, 1989 .

The genus *Didemnum* belongs to the family Didemnidae. The genus contains 236 marine species according to WoRMS.

CO1 was found to be the most suitable region due to the broadest coverage across sister-taxa. However, out of the 236 marine species within the genus *Didemnum* there are only 11 species that are represented by sequence data in the NCBI database. Furthermore, for available sequence data there is an unusually high level of intraspecies variation. Especially the species *D. candidum*, *D. granulatum* and to a lesser extent *D. vexillum*, all of which have multiple individual repository samples, exhibit more than 60 mismatches when aligned to other sequences. Species misidentification is common for this genus as there are few morphological characteristics and a high level of variability within the relevant characters (Stefaniak *et al.* 2009). Although the majority of sequence data for *D. vexillum* show similarity, the uncertainty presented by both the potential misidentification within the genus and the lack of genetic data for 225 sister-species, suggest that the specificity of any assay could be suboptimal.

Species 35: Fucus evanescens

Full name: *Fucus evanescens* C. Agardh, 1820

There are 14 species and variates within the genus *Fucus*. For the species in question, *Fucus evanescens*, only three genes were abundant enough within the genus to be considered for the task at hand. However:

- 18S not suitable to design species specific eDNA primer/probe assay.
- 23S does not provide useful sequence mismatch among *Fucus evanescens*, *F. serratus*, and *F. gardneri*. For the remaining species assay design was prohibited due to the formation of secondary structures and self-annealing of the primers or probe.
- 26S did also not provide the necessary sequence mismatch between target and non-target species. At the one area where an assay potentially could have been positioned this was prevented by the formation of secondary structures and self-annealing of the primers or probe.

Because of this no eDNA primer/probe assay could be designed for *Fucus evanescens*.

Species 36: Petricolaria pholadiformis

Full name *Petricolaria pholadiformis* (Lamarck, 1818)

The genus *Petricolaria* comprises 7 species (WoRMS Editorial Board 2015). But only *Petricolaria pholadiformis* (Lamarck, 1818) and *Petricola lapicida* (Gmelin, 1791) from Petricolidae are represented by nucleotide sequence data on the NCBI GenBank database (Nov-2015).

The recommended primers and probes will likely detect eDNA from the nuclear 28s gene in *Petricolaria pholadiformis*, but whether it will detect eDNA from 28s from other species of *Petricolaria* is not possible to say without first obtaining additional sequence information from 28s genes from other representatives of *Petricolaria* (Table S36.1).

- Pet_pho_28S_F01: 5'-AATTCAAGCCGGGTGTCTT-3',
- Pet_pho_28S_R01: 5'-AGAAAGTGCACCGGGTCTC-3',
- Pet_pho_28S_P01: 5'-FAM-ATCGTCCCTCTCGGAGGGAAGAC-BHQ-1-3'

Table S36.1: Species specific primer/probe assay for *Petricolaria pholadiformis* with, target gene, product size, melting temperature, primer/probe length, GC ratio (%), and number of mismatches between primer and probe region in closely related non-target species.

Species	Gene	Product size	Temp	Length	GC
<i>Petricolaria pholadiformis</i>	28S	104 bp			
Pet_pho_28S_F01	AATTCAAGCCGGGTGTCTT		60.9	20	45
Pet_pho_28S_R01	AGAAAGTGCACCGGGTCTC		61.6	20	55
Pet_pho_28S_P01	ATCGTCCCTCTCGGAGGGAAGAC		68.9	24	58
Related species	Forward	Reverse	Probe		
<i>Petricola lapicida</i>	4	2	6		

In case the recommended primers and probes for the 28s region is unable to differentiate between *Petricolaria pholadiformis* and other species of *Petricolaria*, a second system (Table S36.2) of primers and probes for detection of the mitochondrial cytochrome oxidase 1 gene is also recommended. But without other species of *Petricolaria* than just *Petricolaria pholadiformis* in the NCBI Gen Bank database, it is not possible to rule out that these primers and probes might return positive detection on other species of *Petricolaria*. It would require additional sequencing of correctly identified vouchered specimens of the other six species of *Petricolaria*, to ascertain whether the recommended system truly is species-specific.

- Pet_pho_CO1_F02: 5'-GATCCGGTTAACGGGTACG-3',
- Pet_pho_CO1_R02: 5'-AAAATAGTCCCCGGATGAGC-3',
- Pet_pho_CO1_P02: 5'-FAM-GCTTCAGTGTGATTATCGTATGGAA-BHQ-1-3'

Table S36.2: Species specific primer/probe assay for *Petricolaria pholadiformis* with, target gene, product size, melting temperature, primer/probe length, GC ratio (%), and number of mismatches between primer and probe region in closely related non-target species.

Species	Gene	Product size	Temp	Length	GC
<i>Petricolaria pholadiformis</i>	CO1	70 bp			
Pet_pho_CO1_F01	GATCCGGTTAACGGGTACG		59.2	20	50
Pet_pho_CO1_R01	AAAATAGTCCCCGGATGAGC		60.3	20	50
Pet_pho_CO1_P01	GCTTCAGTGTGATTATCGTATGGAA		63.5	24	37
Related species	Forward	Reverse	Probe		
<i>Petricola lapicida</i>	6	8	6		

Species 37: Percottus glenii

Full name: *Percottus glenii* Dybowski, 1877.

The genus *Percottus* includes the target species *P. glenii* only (Froese & Pauly 2015; WoRMS 2015). The genus belongs to the family Odontobutidae, containing five genera with a total of 21 species (Froese & Pauly 2015).

Sequence data were obtained for both *P. glenii* and the most closely related species from sister genera. Sequence data was available for eight out of a total of 21 species within the family Odontobutidae. Cytochrome b was found to be the most suitable region with best interspecies coverage.

The following primers and probe are designed to target *P. glenii*:

- Per_Gle_Cytb_F01: 5'-AGACAATGCAACTCTTACCCGA-3'
- Per_Gle_Cytb_R01: 5'-CTGCCGGGTTATTGAGCCT-3'
- Per_Gle_Cytb_P01: 5'-FAM-GTTCCCTTTGTTATTGCAGCATTCACTGT-BHQ-1-3'

Table S37: Species-specific primer/probe assay for *Percottus glenii* with target gene, product size, melting temperature, primer/probe length, GC ratio (%), and number of mismatches between primer and probe region in closely related non-target species.

Species	Gene	Product size	Temp	Length	GC
<i>Percottus glenii</i>	Cytb	119 bp			
Per_Gle_Cytb_F01	AGACAATGCAACTCTTACCCGA		59.7	22	46
Per_Gle_Cytb_R01	CTGCCGGGTTATTGAGCCT		60.4	20	60
Per_Gle_Cytb_P01	GTTCCCTTTGTTATTGCAGCATTCACTGT		65.0	30	40
Related species	F01	R01	P01		
<i>Odontobutis yaluensis</i>	3	4	12		
<i>Odontobutis potamophila</i>	3	7	11		
<i>Odontobutis interrupta</i>	2	4	11		
<i>Odontobutis platycephala</i>	3	3	14		
<i>Odontobutis obscura</i>	2	3	14		
<i>Odontobutis sinensis</i>	2	4	11		
<i>Micropercops cinctus</i>	2	5	13		

Species 38: Elminius modestus

Full name: *Elminius modestus* Darwin, 1854.

The genus *Elminius* includes six marine species (WoRMS 2015). The genus belongs to the family Austrobalanidae, containing two subfamilies and eight genera with a total of 23 species.

Multiple genetic regions including 12S, 16S, 18S, 28S and CO1 were investigated, but all regions have limited sequence material available for closely related species, both on a genus and a family level. Based on the data available at NCBI the assay is specific, but false positives with sister-species occurring in Danish territorial waters cannot be excluded.

Following primers and probe have been designed to target *E. modestus*:

- Aus_Mod_CO1_F01: 5'-GGAGCAGGTACCGGATGAAC-3'
- Aus_Mod_CO1_R01: 5'-AGAGATCTACTGACGCTCCTGA-3'
- Aus_Mod_CO1_P01: 5'-FAM-ACCCTCCCTGTCTAGAAATATTGCTCAC-BHQ-1-3'

Table S38: Species-specific primer/probe assay for *Elminius modestus* with target gene, product size, melting temperature, primer/probe length, GC ratio (%), and number of mismatches between primer and probe region in closely related non-target species.

Species	Gene	Product size	Temp	Length	GC
<i>Austrominius modestus</i>	CO1	76 bp			
Aus_Mod_CO1_F01	GGAGCAGGTACCGGATGAAC		60.2	20	60
Aus_Mod_CO1_R01	AGAGATCTACTGACGCTCCTGA		59.8	22	50
Aus_Mod_CO1_P01	ACCCTCCCTGTCTAGAAATATTGCTCAC		65.6	29	48
Related species	F01	R01	P01		
<i>Epopella plicata</i>	3	6	6		
<i>Austrobalanus imperator</i>	3	2	6		

Species 39: Ficopomatus enigmaticus

Full name: *Ficopomatus enigmaticus* (Fauvel, 1923)

The genus *Ficopomatus* comprises five species (Ten Hove & Kupriyanova 2009), and three of these (*Ficopomatus macrodon*, *F. miamiensis* and *F. enigmaticus*) are represented with nuclear 18s and 28s nucleotide sequence information on the NCBI database.

For detection of eDNA from 28s nuclear DNA from *Ficopomatus enigmaticus* two sets of primers and probes are recommended (Table S39.1):

- Fic_eni_28S_F01: 5'-GCAAGGTAGCTTGTGTCGCT-3',
- Fic_eni_28S_R01: 5'-GCGCACCAGACAGTCAGTAA-3',
- Fic_eni_28S_P01: 5'-FAM-CGGCGCAAGAACITATAGGTGGCTAT-BHQ-1-3'

Table S39.1: Species specific primer/probe assay for *Ficopomatus enigmaticus* with, target gene, product size, melting temperature, primer/probe length, GC ratio (%), and number of mismatches between primer and probe region in closely related non-target species.

Species	Gene	Product size	Temp	Length	GC
<i>Ficopomatus enigmaticus</i>	28S	72 bp			
Fic_eni_28S_F01		GCAAGGTAGCTTGTGTCGCT	59.5	20	55
Fic_eni_28S_R01		GCGCACCAGACAGTCAGTAA	60.1	20	55
Fic_eni_28S_P01		CGGCGCAAGAACITATAGGTGGCTAT	68.6	27	48
Related species	Forward	Reverse	Probe		
<i>Amplicaria spiculosa</i>	8	10	11		
<i>Apomatus globifer</i>	10	12	13		
<i>Bushiella abnormis</i>	9	11	9		
<i>Circeis armoricana</i>	5	13	11		
<i>Circeis spirillum</i>	4	13	10		
<i>Crucigera inconstans</i>	10	12	13		
<i>Crucigera tricornis</i>	9	13	14		
<i>Crucigera zygophora</i>	10	12	12		
<i>Ditrupa arietina</i>	5	5	13		
<i>Eulaeospira convexis</i>	8	9	12		
<i>Ficopomatus enigmaticus</i>	0	0	0		
<i>Ficopomatus macrodon</i>	4	12	11		
<i>Ficopomatus miamiensis</i>	1	3	6		
<i>Filograna implexa</i>	8	9	10		
<i>Filigranella elatensis</i>	9	10	12		
<i>Filigranula stellata</i>	9	11	10		
<i>Floriprotis sabiuraensis</i>	10	11	13		
<i>Galeolaria hystrix</i>	7	6	12		
<i>Hydroides brachyacanthus</i>	10	13	12		
<i>Hydroides elegans</i>	8	11	12		
<i>Hydroides ezoensis</i>	9	11	12		
<i>Hydroides minax</i>	9	11	12		
<i>Hydroides pseudouncinatus</i>	10	11	12		
<i>Hydroides sanctaerucis</i>	9	12	11		
<i>Hydroides trivesiculosus</i>	9	12	12		
<i>Janua pagenstecheri</i>	11	7	9		
<i>Josephella marenzelleri</i>	11	12	13		
<i>Jugaria quadrangularis</i>	11	12	10		
<i>Laminatubus alvini</i>	5	7	9		

<i>Marifugia caratica</i>	1	3	5
<i>Metalaeospira tenuis</i>	8	11	11
<i>Neodexiospira brasiliensis</i>	7	9	11
<i>Neodexiospira nipponica</i>	6	9	9
<i>Neodexiospira steueri</i>	9	10	10
<i>Neovermilia globula</i>	8	3	14
<i>Paradexiospira vitrea</i>	5	14	8
<i>Paraprotis dendrova</i>	7	10	10
<i>Pileolaria marginata</i>	12	12	11
<i>Pileolaria militaris</i>	15	12	15
<i>Placostegus tridentatus</i>	5	11	8
<i>Pomatoceros lamarckii</i>	7	7	9
<i>Pomatostegus stellatus</i>	7	9	15
<i>Protolaeospira eximia</i>	12	10	11
<i>Pseudovermilia occidentalis</i>	10	9	10
<i>Rhodopsis pusilla</i>	6	11	13
<i>Semivermilia elliptica</i>	8	11	10
<i>Serpula columbiana</i>	10	12	12
<i>Serpula uschakovi</i>	10	12	12
<i>Serpula vermicularis</i>	9	11	15
<i>Serpula vittata</i>	12	11	12
<i>Serpula watsoni</i>	9	11	13
<i>Simplaria potswaldi</i>	12	12	11
<i>Spirobranchus corniculatus</i>	4	9	7
<i>Spirobranchus lima</i>	8	10	9
<i>Spirorbis bifurcatus</i>	10	11	8
<i>Spirorbis bushi</i>	10	13	9
<i>Spirorbis corallinae</i>	10	12	10
<i>Spirorbis rupestris</i>	11	12	10
<i>Spirorbis spirorbis</i>	11	13	10
<i>Spirorbis tridentatus</i>	11	12	8
<i>Vermiliopsis pygidialis</i>	8	10	12
<i>Vinearia koehleri</i>	12	11	11

In addition a second set (Table S39.2) is also recommended to enable detection of eDNA from 18S nuclear DNA from *Ficopomatus enigmaticus*:

Fic_eni_18S_F02: 5'-TTGACGGCGCITACACATAG-3',

Fic_eni_18S_R02: 5'-ATAAGCCACCCACGGTCAC-3',

Fic_eni_18S_P02: 5'-FAM-CTCATGATTGTCGGCATCTGGCAAC-BHQ-1-3'

Table S39.2 Species specific primer/probe assay for *Ficopomatus enigmaticus* with, target gene, product size, melting temperature, primer/probe length, GC ratio (%), and number of mismatches between primer and probe region in closely related non-target species.

Species	Gene	Product size	Temp	Length	GC
<i>Ficopomatus enigmaticus</i>	18S	103 bp			
<i>Fic_eni_18S_F02</i>	TTGACGGCGCITACACATAG		59.9	20	50
<i>Fic_eni_18S_R02</i>	ATAAGCCACCCACGGTCAC		60.8	19	58
<i>Fic_eni_18S_P02</i>	CTCATGATTGTCGGCATCTGGCAAC		70.7	25	52
Related species	Forward	Reverse	Probe		
<i>Amphicorina mobilis</i>	14	8	1		
<i>Amplicaria spiculosa</i>	14	12	13		
<i>Apomatus globifer</i>	15	9	12		
<i>Apomatus voightae</i>	11	8	12		

<i>Bathyvermilia eliasoni</i>	12	5	11
<i>Bispira melanostigma</i>	14	8	1
<i>Bushiella abnormis</i>	11	9	24
<i>Calcisabella piloseta</i>	15	7	1
<i>Chitinopoma serrula</i>	12	4	11
<i>Circeis armoricana</i>	14	9	31
<i>Circeis spirillum</i>	15	8	25
<i>Crucigera inconstans</i>	11	5	1
<i>Crucigera tricornis</i>	12	5	1
<i>Crucigera zygomphora</i>	13	8	1
<i>Ditrupa arietina</i>	5	3	15
<i>Echinofabricia alata</i>	12	16	17
<i>Eulaeospira convexis</i>	13	11	2
<i>Ficopomatus enigmaticus</i>	0	0	0
<i>Ficopomatus enigmaticus</i>	0	0	0
<i>Ficopomatus macrodon</i>	3	7	13
<i>Ficopomatus miamiensis</i>	3	1	6
<i>Ficopomatus shenzhensis</i>	2	2	11
<i>Filograna implexa</i>	13	7	19
<i>Filogranella elatensis</i>	13	8	12
<i>Filogranula stellata</i>	12	4	11
<i>Floriprotis sabiuraensis</i>	11	5	1
<i>Galeolaria caespitosa</i>	6	4	15
<i>Galeolaria hystrix</i>	6	5	14
<i>Gunnarea capensis</i>	12	7	0
<i>Hyalopomatus biformis</i>	5	4	13
<i>Hyalopomatus mironovi</i>	5	3	14
<i>Hydroides brachyacanthus</i>	11	5	1
<i>Hydroides elegans</i>	12	9	2
<i>Hydroides ezoensis</i>	12	6	1
<i>Hydroides minax</i>	11	5	1
<i>Hydroides norvegicus</i>	13	3	2
<i>Hydroides novae-pomeraniae</i>	11	5	1
<i>Hydroides pseudouncinatus</i>	12	6	0
<i>Hydroides sanctaecrucis</i>	11	5	1
<i>Hydroides trivesiculosus</i>	11	5	1
<i>Hydroides tuberculatus</i>	11	5	1
<i>Janua pagenstecheri</i>	12	12	14
<i>Josephella marenzelleri</i>	12	4	14
<i>Jugaria quadrangularis</i>	11	9	24
<i>Laminatubus alvini</i>	5	4	13
<i>Manayunkia athalassia</i>	15	13	22
<i>Marisugia caratica</i>	1	2	10
<i>Metalaeospira tenuis</i>	13	11	2
<i>Metavermilia acanthophora</i>	11	7	14
<i>Neodexiospira brasiliensis</i>	13	12	21
<i>Neodexiospira nipponica</i>	13	12	21
<i>Neodexiospira steueri</i>	13	12	18
<i>Neovermilia globula</i>	13	5	0
<i>Novafabricia labrus</i>	13	13	19
<i>Paradexiospira vitrea</i>	15	9	29
<i>Paraprotis dendrova</i>	13	9	14
<i>Pileolaria marginata</i>	12	9	21
<i>Pileolaria militaris</i>	13	9	26

<i>Pomatoceros lamarckii</i>	5	3	13
<i>Pomatoceros taeniata</i>	5	3	14
<i>Pomatoceros triqueter</i>	5	3	15
<i>Pomatoleios kraussii</i>	6	3	14
<i>Pomatostegus stellatus</i>	12	4	12
<i>Protis hydrothermica</i>	13	8	17
<i>Protolaeospira capensis</i>	13	11	2
<i>Protolaeospira eximia</i>	12	11	2
<i>Protolaeospira tricostalis</i>	13	11	2
<i>Protula atypba</i>	15	7	13
<i>Protula bispiralis</i>	13	9	12
<i>Protula palliata</i>	13	9	12
<i>Protula tubularia</i>	16	9	12
<i>Pseudochitinopoma occidentalis</i>	7	3	13
<i>Pseudovermilia occidentalis</i>	15	8	24
<i>Rhodopsis pusilla</i>	15	6	23
<i>Romanchella quadricostalis</i>	13	11	2
<i>Sabella spallanzanii</i>	14	7	1
<i>Sabellaria alveolata</i>	12	7	0
<i>Selenidium neosabellariae</i>	9	16	1
<i>Selenidium sensimiae</i>	8	15	2
<i>Semivermilia elliptica</i>	15	8	24
<i>Serpula columbiana</i>	11	5	1
<i>Serpula concharum</i>	10	5	1
<i>Serpula jukesii</i>	11	5	1
<i>Serpula uschakovi</i>	12	8	1
<i>Serpula vermicularis</i>	11	5	1
<i>Serpula vittata</i>	11	6	1
<i>Serpula watsoni</i>	11	5	1
<i>Simplaria potswaldi</i>	12	9	24
<i>Spirobranchus cariniferus</i>	5	3	13
<i>Spirobranchus corniculatus</i>	5	4	15
<i>Spirobranchus latiscapus</i>	5	3	15
<i>Spirobranchus lima</i>	5	3	13
<i>Spirorbis bifurcatus</i>	13	9	24
<i>Spirorbis bushi</i>	13	11	25
<i>Spirorbis corallinae</i>	12	9	26
<i>Spirorbis rupestris</i>	12	9	26
<i>Spirorbis spirorbis</i>	12	9	25
<i>Spirorbis tridentatus</i>	12	9	26
<i>Vermiliopsis infundibulum</i>	12	8	20
<i>Vermiliopsis labiata</i>	11	8	21
<i>Vermiliopsis pygidialis</i>	8	9	16
<i>Vermiliopsis striaticeps</i>	12	8	21
<i>Vinearia koehlerii</i>	11	8	25

Because there are species of *Ficopomatus* that are not represented by nucleotide sequences on the NCBI database, the recommended primer and probe systems suggested here are may or may not be able to distinguish between *Ficopomatus enigmatus* and other non-sequenced species of *Ficopomatus*.

Species 40: Marenzelleria viridis

Full name: *Marenzelleria viridis* (Verrill, 1873)

The genus *Marenzelleria* comprises seven species (WoRMS Editorial Board 2015). In European seas two species of *Marenzelleria* are reported (Bellan 2001). On the NCBI GenBank database five species of *Marenzelleria* are represented by nucleotide sequence information. It might be that there are multiple haplotypes of (Blank & Bastrop 2009) of *Marenzelleria viridis* occurring in North European seas, and it is thus difficult to assure the recommended primer-probe system will be able to detect all haplotypes of *Marenzelleria viridis* occurring in Northern European seas, while also not returning false positives from other co-occurring species of *Marenzelleria* (Blank & Bastrop 2009, Bellan 2001). To accommodate this multiple primer-probe systems are here recommended for being able to discriminate between the various species and haplotypes of *Marenzelleria*.

For the mitochondrial 16s gene region the following primers and probes are recommended for detection of eDNA from *Marenzelleria viridis* (Table S40.1):

- Mar_vir_16S_F01: 5'-AAGCTCCTATAGCTACGCAAA-3',
- Mar_vir_16S_R01: 5'-GAAGATTAGGTCTCTCTTGTGCG-3',
- Mar_vir_16S_P01: 5'-FAM-GAACTTACCTACAGAAATTGTGTTGGGG-BHQ1-3'

Table S40.1: Species specific primer/probe assay for *Marenzelleria viridis* with, target gene, product size, melting temperature, primer/probe length, GC ratio (%), and number of mismatches between primer and probe region in closely related non-target species.

Species	Gene	Product size	Temp	Length	GC
<i>Marenzelleria viridis</i>	16S	76 bp			
Mar_vir_16S_F01		AAGCTCCTATAGCTACGCAAA	57.6	22	41
Mar_vir_16S_R01		GAAGATTAGGTCTCTCTTGTGCG	57.8	24	41
Mar_vir_16S_P01		GAACTTACCTACAGAAATTGTGTTGGGG	64.8	29	38

Related species	Forward	Reverse	Probe
<i>Laonice norgensis</i>	16	12	14
<i>Malacoceros fuliginosus</i>	15	2	10
<i>Malacoceros indicus</i>	16	13	13
<i>Malacoceros indicus</i>	16	13	13
<i>Marenzelleria arctia</i>	5	0	7
<i>Marenzelleria bastropi</i>	4	0	3
<i>Marenzelleria neglecta</i>	2	0	2
<i>Marenzelleria viridis</i>	0	0	0
<i>Marenzelleria wireni</i>	4	0	3
<i>Rhynchospio foliosa</i>	15	5	11
<i>Scolelepis eltaninae</i>	17	3	14

In case the recommended system for detection of mtDNA-16s from *M. viridis* turns out to be unspecific, a system for the mitochondrial cytochrome b gene region is also recommended for detection of eDNA from *Marenzelleria viridis* (Table S40.2):

- Mar_vir_Cytb_F02: 5'-CGATTCTTGCGTTCCATTT-3'
- Mar_vir_Cytb_R02: 5'-GCCAAGAGGGTTATTGATCC-3'
- Mar_vir_Cytb_P02: 5'-FAM-CCCTCTCAGGGGTTCACCTCTATTTC-BHQ-1-3'

Table S40.2: Species specific primer/probe assay for *Marenzelleria viridis* with, target gene, product size, melting temperature, primer/probe length, GC ratio (%), and number of mismatches between primer and probe region in closely related non-target species.

Species	Gene	Product size	Temp	Length	GC
<i>Marenzelleria viridis</i>	Cytb	102 bp			
Mar_vir_Cytb_F02		CGATTCTTTGCGTTCCATT	60.1	20	40
Mar_vir_Cytb_R02		GCCAAGAGGGTTATTGATCC	59.8	21	48
Mar_vir_Cytb_P02		CCCTCTCAGGGGTTCACCTTCTATTTC	67.9	27	52
Related species	Forward	Reverse	Probe		
<i>Boccardia proboscidea</i>	6	4	12		
<i>Boccardiella hamata</i>	4	6	14		
<i>Malacoceros fuliginosus</i>	2	6	8		
<i>Marenzelleria arctia</i>	4	4	9		
<i>Marenzelleria bastropi</i>	5	3	8		
<i>Marenzelleria neglecta</i>	1	6	5		
<i>Marenzelleria viridis</i>	0	0	0		
<i>Marenzelleria wireni</i>	2	5	6		
<i>Polydora brevipalpa</i>	5	4	14		
<i>Polydora hoplura</i>	5	5	14		
<i>Polydora websteri</i>	5	2	12		

Species 41: Ocenebra inornata

Full name: *Ocenebra inornata* Récluz, 1851

The *Ocenebra* genus comprises seven marine species. The genus belongs to the subfamily Ocenebrinae which has 28 genera, according to WoRMS.

Out of the seven marine species, sequence data was only available for *O. inornata* and *O. erinacea*. The sequence data from *O. inornata* was compared to related species and to species showing sequence similarity via sequence BLAST analysis. Two assays are designed, spanning two different regions of the CO1 segment.

The first set of primers and probe to target *O. inornata* is:

- Oce_Ino_CO1_F01: 5'-GGCTTGTGCGGAAGTGCCTTA-3'
- Oce_Ino_CO1_R01: 5'-TCATCCCCAAGCAAAGCTCC-3'
- Oce_Ino_CO1_P01: 5'-FAM-AGTCTTCTTATTGAGCTGAACTAGGGCA-BHQ-1-3'

Table S41.1: Species-specific primer/probe assay for *Ocenebra inornata* with target gene, product size, melting temperature, primer/probe length, GC ratio (%), and number of mismatches between primer and probe region in closely related non-target species.

Species	Gene	Product size	Temp	Length	GC
<i>Ocenebra inornata</i>	CO1	73 bp			
Oce_Ino_CO1_F01		GGCTTGTGCGGAAGTGCCTTA	60.3	20	55
Oce_Ino_CO1_R01		TCATCCCCAAGCAAAGCTCC	60.3	20	55
Oce_Ino_CO1_P01		AGTCTTCTTATTGAGCTGAACTAGGGCA	65.3	29	45
Related species	F01	R01	P01		
<i>Ocenebra erinacea</i>	4	7	2		
<i>Ocenebrina edwardsii</i>	4	7	1		
<i>Ocenebrina lurida</i>	1	2	1		
<i>Purpura persica</i>	7	6	0		
<i>Tiariturris spectabilis</i>	7	5	0		
<i>Turris hidalgovi</i>	4	1	0		
<i>Xenuroturris albino</i>	4	3	1		

The second set of primers and probe designed to target *O. inornata*:

- Oce_Ino_CO1_F02: 5'-ACCCCCTGCCTTACTTCCTTT-3'
- Oce_Ino_CO1_R02: 5'-ACAGATCCTCCAGCATGAGC-3'
- Oce_Ino_CO1_P02: 5'-FAM-TGGACTGTCTACCCTCCATTAGCTGGT-BHQ-1-3'

Table S41.2: Species-specific primer/probe assay for *Ocenebra inornata* with target gene, product size, melting temperature, primer/probe length, GC ratio (%), and number of mismatches between primer and probe region in closely related non-target species.

Species	Gene	Product size	Temp	Length	GC
<i>Ocenebra inornata</i>	CO1	114 bp			
Oce_Ino_CO1_F02		ACCCCCTGCCTTACTTCTTTT	59.2	21	48
Oce_Ino_CO1_R02		ACAGATCCTCCAGCATGAGC	59.5	20	55
Oce_Ino_CO1_P02		TGGACTGTCTACCCTCCATTAGCTGGT	66.4	27	52
Related species	F01	R01	P01		
<i>Ocenebra erinacea</i>	5	1	7		
<i>Ocinebrina edwardsii</i>	5	1	5		
<i>Ocinebrina lurida</i>	3	1	3		
<i>Purpura persica</i>	8	3	4		
<i>Tiariturris spectabilis</i>	7	2	6		
<i>Turris hidalgoi</i>	5	3	5		
<i>Xenuroturris albino</i>	10	5	4		

Species 42: Potamopyrgus antipodarum

Full name: *Potamopyrgus antipodarum* Gray, 1843.

The genus *Potamopyrgus* consists of 11 marine species and belongs to the family Tateidae (WoRMS 2015).

An assay for *P. antipodarum* has already been published and validated based on the cytochrome b region (Goldberg *et al.* 2013). However, sequence data on the cytochrome b region is only available for one sister-species at NCBI; hence occurrence of non-target sister species may lead to false positives.

- Pot_Ant_Cytb_F01: 5'- TGTITCAAGTGTGCTGGTTAYA-3'
- Pot_Ant_Cytb_R01: 5'-CAAATGGRGCTAGTTGATTCTT-3'
- Pot_Ant_Cytb_P01: 5'-FAM-CCTCGACCAATATGTAAAT-MGB-NFQ-3'

The annealing temperature was 60 °C in the study by Goldberg *et al.* (2013), whereas the values in Table S42 are the optimal temperatures calculated in Geneious version 9.0.4.

Table S42: Species-specific primer/probe assay for *Potamopyrgus antipodarum* with target gene, product size, melting temperature, primer/probe length, GC ratio (%), and number of mismatches between primer and probe region in closely related non-target species.

Species	Gene	Product size	Temp	Length	GC
<i>Potamopyrgus antipodarum</i>	Cytb	93 bp			
Pot_Ant_Cytb_F01		TGTITCAAGTGTGCTGGTTAYA	56.6 - 59.2	23	35 - 39
Pot_Ant_Cytb_R01		CAAATGGRGCTAGTTGATTCTT	55.2 - 57.3	23	35 - 39
Pot_Ant_Cytb_P01		CCTCGACCAATATGTAAAT*	49.5	19	36
*TaqMan MGB probe					
Related species	F01	R01	P01		
<i>Potamopyrgus estuarinus</i>	1	1	0		

Species 43: Pseudochattonella verruculosa

Full name: *Pseudochattonella verruculosa* (Y.Hara & M.Chihara) S.Tanabe-Hosoi, D.Honda, S.Fukaya, Y.Inagaki & Y.Sako

The genus *Pseudochattonella* only contains the two species *Pseudochattonella farcimen* and *Pseudochattonella verruculosa*. The two species are more or less impossible to morphologically tell apart after fixation in lygol. Due to this a molecular method has been developed and to date serve as the best solution to tell apart the two species (Dittami *et al.* 2013). However, the designed primers (best possible) only have two mismatches each, respectively. According to the published qPCR protocol it is relatively difficult even with the molecular method to determine if the amplification in the qPCR reaction is caused by the presence of the one or the other species (Dittami *et al.* 2013):

- PverF: 5'-GGTCCCGGGGAGAAGTC-3'
- PverR: 5'-AAGCAACTCGACTCCATTAGC-3'

No other mismatches exist within the sequence between the two primers. Therefore, no probe was designed. However, it has come to our attention that a primer/probe assay is under development and awaiting publication (Eckford-Soper & Daugbjerg (in press)). With this new assay it is expected that differentiation between the two *Pseudochattonella* species will possible.

Table S43: Species specific primer/probe assay for *Pseudochattonella verruculosa* with, target gene, product size, melting temperature, primer/probe length, and GC ratio (%)

Species	Gene	Product size	Temp	Length	GC
<i>Pseudochattonella verruculosa</i>	28S	91 bp			
PverF		GGTCCCGGGGAGAAGTC	59.8	17	71
PverR		AAGCAACTCGACTCCATTAGC	59.5	21	48

Species 44: Codium fragile* subsp. *fragile

Full name: *Codium fragile* subsp. *fragile* (Suringar) Hariot, 1889

Codium fragile subsp. *fragile* is as the name indicates a subspecies to *Codium fragile* together with three other subspecies. These subspecies belongs to the genus *Codium* showing a very large number of species (n137). Within the subspecies group only the sequence for the ribosomal protein S3 (*rps3*) and L16 (*rpl16*) is represented by all four subspecies:

- Cod_sub_fra_rps3_F01: 5'-AAATGTTACTCATGCTCCAACC-3'
- Cod_sub_fra_rps3_R01: 5'-CCAATAGGGCTGTAATGCC-3'
- Cod_sub_fra_rps3_P01: 5'-FAM-AAGTAAATTTCGTAAAAATCATCGCGG-BHQ-1-3'

However, the sequence showed no usable mismatches among the four subspecies (represented by *Codium fragile* subsp. *novae-zelandiae* in table S44). Despite this and the relative large size of the amplicon the *rps3* sequence provides a good basis for future eDNA work on the species. The assay was designed based on the alignment of sequence information from no less than 71 species within the genus *Codium*. However, due to simplicity only nine species with less than six mismatches have been listed in the table S44. Five species (first five in the table) were selected to represent the remaining 62 species from the comparison. These are also listed in table S44, however without mismatch information.

Table S44: Species specific primer/probe assay for *Codium fragile* subsp. *fragile* with, target gene, product size, melting temperature, primer/probe length, GC ratio (%), and number of mismatches between primer and probe region in closely related non-target species. Problematic species are marked in bold

Species	Gene	Product size	Temp	Length	GC
<i>Codium fragile</i> subsp. <i>fragile</i>	<i>rps3-rpl16</i>	138 bp			
Cod_sub_fra_rps3_F01	AAATGTTACTCATGCTCCAACC		58.4	22	41
Cod_sub_fra_rps3_R01	CCAATAGGGCTGTAATGCC		57.5	19	53
Cod_sub_fra_rps3_P01	AAGTAAATTTCGTAAAAATCATCGCGG		61.1	27	33
Related species	F01	R01	P01		
<i>Codium fragile</i> subsp. <i>novae-zelandiae</i>	0	0	0		
<i>Codium adhaerens</i>	7	6	6		
<i>Codium arabicum</i>	6	6	6		
<i>Codium barbatum</i>	8	5	4		
<i>Codium bursa</i>	8	6	2		
<i>Codium capitatum</i>	7	5	3		
<i>Codium capitulatum</i>	6	7	7		
<i>Codium crannelliae</i>	4	6	3		
<i>Codium cylindricum</i>	3	4	3		
<i>Codium decorticatum</i>	2	4	3		
<i>Codium duttieae</i>	2	4	2		
<i>Codium galeatum</i>	3	3	2		
<i>Codium minus</i>	4	8	3		
<i>Codium papenfussii</i>	3	4	2		
<i>Codium</i> sp. DML66031	1	5	3		
<i>Codium yezoense</i>	1	2	1		
<i>Codium</i> cf. <i>bursa</i> DHO2-176					
<i>Codium</i> cf. <i>dimorphum</i> C29					
<i>Codium</i> cf. <i>dimorphum</i> C77					
<i>Codium</i> cf. <i>flabellatum</i> DHO-009					
<i>Codium</i> cf. <i>fragile</i> AU5					

Codium cf. latum ASH-018
Codium cf. latum C51
Codium cf. minus DHO-015
Codium cf. minus DHO2-188
Codium cf. tenue HV608
Codium contractum
Codium convolutum
Codium coralloides
Codium dimorphum
Codium effusum
Codium geppiorum
Codium gracile
Codium hubbsii
Codium hubbsii
Codium intertextum
Codium intricatum
Codium isthmocladum
Codium isthmocladum subsp. *clavatum*
Codium laminarioides
Codium latum
Codium lucasii
Codium lucasii subsp. *capense*
Codium megalophysum
Codium muelleri
Codium ovale
Codium parvulum
Codium platylobium
Codium prostratum
Codium repens
Codium saccatum
Codium setchellii
Codium sp. 'arenicola'
Codium sp. DB2006
Codium sp. DHO-007
Codium sp. DHO2-196
Codium sp. DHO2-348
Codium sp. DML30929
Codium sp. DML30930
Codium sp. DML40218
Codium sp. DML40227
Codium sp. DML65827
Codium sp. DML65829
Codium sp. HV1061
Codium sp. HV1068
Codium spinescens
Codium spongiosum
Codium subtubulosum
Codium taylorii
Codium vermilara

Species 45: *Carassius auratus*

Full name: *Carassius auratus* (Linnaeus, 1758).

The genus *Carassius* belongs to the family Cyprinidae and includes five species (Froese & Pauly 2015). Occurrences of hybridization between *C. auratus* and other closely related species, including *Cyprinus carpio*, are reported. Since the mitochondrial genome of these hybrids is maternally inherited the assay will be unable to distinguish hybrids from normal individuals in cases where the mitochondrion genome originates from *C. auratus* (Yan *et al.* 2005).

An assay for *C. auratus* based on the cytochrome b sequence has been published and validated (Nathan *et al.* 2014). That assay consists of a primer pair with the sequences:

- Car_Aur_Cytb_F01: 5'-GCTTCTCCGTAGATAATG-3'
- Car_Aur_Cytb_R01: 5'-TTCGTGAAGAACAGTAG-3'

Testing the primers reveals limited variation between *C. gibelio* and *C. auratus*. To increase specificity a probe was tested within the amplicon, but the amplicon had no optimal binding sites. Furthermore, the most common intraspecies nucleotide polymorphisms are not represented within the primers sequence and will therefore potentially not bind to the targeted organisms DNA.

Table S45.1: Species-specific primer/probe assay for *Carassius auratus* with target gene, product size, melting temperature, primer/probe length, GC ratio (%), and number of mismatches between primer and probe region in closely related non-target species.

Species	Gene	Product size	Temp	Length	GC
<i>Carassius auratus auratus</i>	Cytb	106 bp			
Car_Aur_Cytb_F01		GCTTCTCCGTAGATAATG	49.3	18	44
Car_Aur_Cytb_R01		TTCGTGAAGAACAGTAG	49.2	18	39

Related species	F01	R01	P01
<i>C. auratus</i>	1	1	-
<i>C. gibelio</i>	1	1	-
<i>C. carassius</i>	2	2	-
<i>Cyprinus carpio</i>	2	2	-
<i>C. auratus</i> x <i>Megalobrama amblycephala</i> EU518483	1	1	-
<i>C. auratus</i> x <i>Megalobrama amblycephala</i> EU518484	1	1	-
<i>C. auratus</i> x 2 <i>Megalobrama amblycephala</i> EU518485	1	2	-
<i>C. auratus</i> x <i>Cyprinus carpio</i> NC_006136	1	1	-
<i>C. auratus</i> x <i>Cyprinus carpio</i> AY694420	1	1	-
<i>C. auratus</i> x <i>Cyprinus carpio</i> GU186888	1	1	-
<i>C. auratus</i> x <i>Cyprinus carpio</i> x <i>Carassius cuvieri</i> AY771781	1	2	-
<i>C. auratus</i> x <i>Cyprinus carpio</i> x <i>Carassius cuvieri</i> C_006387	1	2	-
<i>Cyprinus carpio</i> x <i>C. auratus</i> NC_018038	2	1	-
<i>Cyprinus carpio</i> x <i>C. auratus</i> NC_026543	3	1	-

A secondary assay was designed with the inclusion of a probe with following sequences:

- Car_Aur_Cytb_F02: 5'-GACAGCCTTGTCGGTTATGT-3'
- Car_Aur_Cytb_R02: 5'-TATGGCACGGCGGATAGAAG-3'
- Car_Aur_Cytb_P02: 5'-TCCTTTGAGGCGCTACAGTAATCACAAAC-BHQ-1-3'

A comparison between *C. auratus* and *C. gibelio* showed limited variation, which confines the ability to distinguish these two species. This is potentially due to taxonomic confusion between *C. auratus auratus* and *C. gibelio* and their complex modes of reproduction (Froese & Pauly 2015)

Table S45.2: Species-specific primer/probe assay for *Carassius auratus* with target gene, product size, melting temperature, primer/probe length, GC ratio (%), and number of mismatches between primer and probe region in closely related non-target species.

Species	Gene	Product size	Temp	Length	GC
<i>Carassius auratus auratus</i>	Cytb	90 bp			
Car_Aur_Cytb_F02	GACAGCCTTTGTCGGTTATGT		58.6	21	47
Car_Aur_Cytb_R02	TATGGCACGGCGGATAGAAG		59.7	20	55
Car_Aur_Cytb_P02	TCCTTTGAGGCGCTACAGTAATCACAAAC		65.7	30	43
Related species		F01	R01	P01	
<i>C. auratus</i>		0	0	0	
<i>C. gibelio</i>		0	1	0	
<i>C. carassius</i>		2	3	1	
<i>Cyprinus carpio</i>		4	3	1	
<i>C. auratus</i> × <i>Megalobrama amblycephala</i> EU518483		0	1	1	
<i>C. auratus</i> × <i>Megalobrama amblycephala</i> EU518484		0	0	0	
<i>C. auratus</i> × 2 <i>Megalobrama amblycephala</i> EU518485		0	3	0	
<i>C. auratus</i> × <i>Cyprinus carpio</i> NC_006136		0	0	0	
<i>C. auratus</i> × <i>Cyprinus carpio</i> AY694420		0	0	0	
<i>C. auratus</i> × <i>Cyprinus carpio</i> GU186888		0	1	0	
<i>C. auratus</i> × <i>Cyprinus carpio</i> × <i>Carassius cuvieri</i> AY771781		0	3	1	
<i>C. auratus</i> × <i>Cyprinus carpio</i> × <i>Carassius cuvieri</i> NC_006387		0	3	1	
<i>Cyprinus carpio</i> × <i>C. auratus</i> NC_018038		4	3	1	
<i>Cyprinus carpio</i> × <i>C. auratus</i> NC_026543		4	3	4	

Species 46: Cyprinus carpio

Full name: *Cyprinus carpio* Linnaeus, 1758.

The genus *Cyprinus* comprises two species that can occur in brachish waters and 24 species in total. The genus belongs to the family Cyprinidae containing 3032 species (Froese & Pauly 2015; WoRMS 2015).

Hybridization between *C. carpio* and *Carassius auratus*, is reported. The mitochondrial genome of these hybrids is maternally inherited and the assay will hence not be able to distinguish hybrids from normal individuals in cases where the mitochondrion genome originates from *C. carpio* (Yan *et al.* 2005).

Following primer pair for *C. carpio* has already been published based on the D-loop region (Turner *et al.* 2014):

- Cyp_Car_dloop_F01: 5'-GAGTGCAGGCTCAAATGTTAAA-3'
- Cyp_Car_dloop_R01: 5'-GTAAGGATAAGTTGAACTAGAGACAG-3'

A BLAST analysis and local alignment show that the primers are specific to the species within the genus *Cyprinus*. As *C. carpio* is the only species occurring in Danish territorial waters it is therefore expected that the assay is specific (Froese & Pauly 2015).

An internal probe within the amplicon was developed to increase the specificity of the assay. It has the following sequence:

- Cyp_Car_dloop_P01: 5'-FAM-AAGGTTGAACATTTCCTGTATGTGAT-BHQ-1-3'

Table S46: Species-specific primer/probe assay for *Cyprinus carpio* with target gene, product size, melting temperature, primer/probe length, GC ratio (%), and number of mismatches between primer and probe region in closely related non-target species.

Species	Gene	Product size	Temp	Length	GC
<i>Cyprinus carpio</i>	D-loop	146 bp			
Cyp_Car_dloop_F01	GAGTGCAGGCTCAAATGTTAAA		57.5	22	41
Cyp_Car_dloop_R01	GTAAGGATAAGTTGAACTAGAGACAG		56.9	26	39
Cyp_Car_dloop_P01	AAGGTTGAACATITTCCTGTATGTGAT		60.2	28	32
Related species		F01	R01	P01	
<i>Cyprinus acutidorsalis</i>		0	0	0	
<i>Cyprinus rubrofuscus</i>		0	0	0	
<i>Cyprinus pellegrini</i>		0	0	2	
<i>Cyprinus multitaeniata</i>		0	0	1	
<i>Cyprinus melanes</i>		0	0	0	
<i>Carassius gibelio</i>		4	6	1	
<i>Carassius cuvieri</i>		4	6	3	
<i>Carassius auratus auratus</i>		5	6	1	
<i>Carassius carassius</i>		10	3	2	
<i>C. auratus auratus</i> × <i>C. carpio</i>		4	7	1	
<i>C. auratus auratus</i> × <i>C. carpio</i> × <i>C. cuvieri</i>		4	6	1	

Species 47: Mya arenaria

Full name: *Mya arenaria* Linnaeus, 1758

The genus *Mya* comprises four species and the genus belongs to the family Myoidea. Within the family the CO1 gene is good represented and therefore selected for the assay design:

- Mya_are_CO1_F01: 5'-CGTGCAGGATGTCACCCG-3'
- Mya_are_CO1_R01: 5'-GCCAAAAGTGGCATTGCTAA-3'
- Mya_are_CO1_P01: 5'-FAM-CTGTAGGTCTATTGGAGTTACCAGGTIT-HBQ-1-3'

Table S47: Species specific primer/probe assay for *Mya arenaria* with, target gene, product size, melting temperature, primer/probe length, GC ratio (%), and number of mismatches between primer and probe region in closely related non-target species

Species	Gene	Product size	Temp	Length	GC
<i>Mya arenaria</i>	CO1	95 bp			
Mya_are_CO1_F01	CGTGCAGGATGTCACCCG		60.8	18	67
Mya_are_CO1_R01	CGCCAAAAGTGGCATTGCTAA		59.5	21	48
Mya_are_CO1_P01	CTGTAGGTCTATTGGAGTTACCAGGTIT		67.2	28	43
Related species*	Forward	Reverse	Probe		
<i>Mya truncata</i>	2	7	5		
<i>Varicorbula dissimilis</i>	7	5	8		
<i>Corbula amurensis</i>	7	5	6		
<i>Corbula erythrodon</i>	4	5	3		
<i>Corbula gibba</i>	9	5	12		
<i>Corbula tunicata</i>	5	5	8		
<i>Potamocorbula fasciata</i>	9	5	6		

* not necessarily related but show strong sequence similarity to target species

Species 48: *Penilia avirostris*

Full name: *Penilia avirostris* Dana, 1849

Penilia avirostris is the only species in the genus *Penilia*. However, in the Family Sididae comprise a fairly larger number of species in close relationship to *Penilia avirostris*. For a large portion of Sididae the gene sequence for CO1 was available and was found to be very divergent between the species (Table S48):

- Pen_avi_CO1_F01: 5'-AGCGGGGATTTCATCAATCC-3'
- Pen_avi_CO1_R01: 5'-CCACACAAAGAGGGGAATAC-3'
- Pen_avi_CO1_P01: 5'-FAM- CGACAATTGTAAATATACGATCTAAAGGA-BHQ-1-3'

Table S48: Species specific primer/probe assay for *Penilia avirostris* with, target gene, product size, melting temperature, primer/probe length, GC ratio (%), and number of mismatches between primer and probe region in closely related non-target species

Species	Gene	Product size		Temp	Length	GC
<i>Penilia avirostris</i>	CO1	103 bp				
Pen_avi_CO1_F01		AGCGGGGATTTCATCAATCC		58.4	20	50
Pen_avi_CO1_R01		CCACACAAAGAGGGGAATAC		58.4	20	50
Pen_avi_CO1_P01		CGACAATTGTAAATATACGATCTAAAGGA		63.3	29	31
Related species*		Forward	Reverse	Probe		
<i>Sarsilatona serricauda</i>		6	5	10		
<i>Sarsilatona cf. serricauda</i>		3	6	7		
<i>Sida crystallina americana</i>		5	8	8		
<i>Latonopsis australis</i>		6	4	6		
<i>Latonopsis cf. australis</i>		6	6	6		
<i>Diaphanosoma birgei</i>		4	5	8		
<i>Diaphanosoma brachyurum</i>		6	6	11		
<i>Diaphanosoma brevireme</i>		6	6	4		
<i>Diaphanosoma cf. amurensis</i>		6	7	8		
<i>Diaphanosoma cf. dubium</i>		7	9	9		
<i>Diaphanosoma cf. heberti</i>		6	6	10		
<i>Diaphanosoma cf. macropthalma</i>		6	4	9		
<i>Diaphanosoma cf. orientalis</i>		5	7	6		
<i>Diaphanosoma dubium</i>		6	8	11		
<i>Diaphanosoma sp. 1 MEG-2008</i>		5	9	5		
<i>Diaphanosoma sp. 1 MEG-2013</i>		5	9	5		
<i>Diaphanosoma sp. HE-091</i>		5	4	6		
<i>Diaphanosoma sp. HE-099</i>		5	4	6		
<i>Diaphanosoma sp. HE-572</i>		6	6	7		
<i>Diaphanosoma sp. HE-572.1</i>		6	6	7		
<i>Diaphanosoma sp. HE-648.1</i>		6	4	8		
<i>Diaphanosoma sp. HE-649</i>		6	4	8		
<i>Diaphanosoma sp. HE-649.1</i>		6	4	8		
<i>Diaphanosoma sp. HE-652</i>		6	4	8		
<i>Diaphanosoma sp. IW-2015-1</i>		5	4	6		
<i>Diaphanosoma sp. JRdW-2005</i>		6	6	11		
<i>Diaphanosoma spinulosum</i>		6	5	6		

* not necessarily related but show strong sequence similarity to target species

Species 49: *Diadumene lineata*

Full name: *Diadumene lineata* (Verrill, 1870)

The genus *Diadumene* comprises 11 species (WoRMS Editorial Board 2015). From the family Diadumenidae only *Diadumene cineta* Stephenson, 1925 and *Haliplanella luciae* (Verrill, 1899) appear to be recorded in European seas (van der Land & den Hartog, 2001). Unfortunately, very few species of *Diadumene* are represented with nucleotide sequence information on the NCBI GenBank database (Nov-2015). Also, very few species of both Diadumenidae and the suborder Nynantheae are represented by nucleotide sequence data on NCBI GenBank. Only *Diadumene cincta* and *Diadumene leucolena* have nucleotide sequence information deposited on the NCBI GenBank database. Without any sequence information available for *Diadumene lineata* it is currently not possible to develop a species-specific primer-probe system that only detects *Diadumene lineata* and not any other species of *Diadumene*. The primer-probe systems recommended here are therefore based on various other representatives from Nynantheae.

As only *Diadumene cincta* and *Diadumene leucolena* have nucleotide sequence information deposited on the NCBI GenBank database, the recommended primers and probes have been designed to detect eDNA from these two species instead. More universal primers that will detect eDNA from *Diadumene* spp. are currently not possible to design with the sequence information deposited on the NCBI GenBank database.

A comparison of fragments of nuclear 18s gene markers from Nynantheae unfortunately returns *Diadumene* as a polyphyletic genus, which makes a more universal primer-probe system unable to differentiate between *Diadumene* and various other genera in Nynantheae. The nuclear 18S gene fragments deposited at NCBI GenBank have the highest nucleotide diversity among different gene marker fragments deposited on NCBI, but even the most variable region of 18s from Nynantheae does not have sufficient variation to distinguish between *Diadumene* and several other genera within Nynantheae. This excludes the possibility of designing primers and probes specific for only *Diadumene* spp.

Here two primer-probe systems are recommended, that are targeting eDNA from *Diadumene cincta* and *Diadumene leucolena*. However, it is not possible to determine whether these two systems will enable detection of eDNA from any of the other nine species of *Diadumene*, or if these two systems will be able to differentiate between *Diadumene cincta* and *Diadumene leucolena* and any other species of *Diadumene* spp. Also, the limited nucleotide variation in the 18S gene fragments introduce a risk of obtaining false positive eDNA detection from other species from Nynantheae, as even the most variable region of 18S is unable to distinguish between all genera and species of Nynantheae in just a short stretch of 70 – 120 bp size DNA fragment.

For detection eDNA-fragments form 18S-nDNA from *Diadumene leucolena* the following primers and probes are recommended (Table S49.1):

- Dia_leu_18S_F01: 5'-ACTGACCGGGCTGTTCTTCT-3',
- Dia_leu_18S_R01: 5'-GTCACAAGTCCTACCGCACCA-3',
- Dia_leu_18S_P01: 5'-FAM-AAGGACTGCGTGTGCTCTAGCTGAG-BHQ1-3',

Table S49.1: Species specific primer/probe assay for *Diadumene leucolena* with, target gene, product size, melting temperature, primer/probe length, GC ratio (%), and number of mismatches between primer and probe region in closely related non-target species.

Species	Gene	Product size	Temp	Length	GC
<i>Diadumene leucolena</i>	18S	70 bp			
Dia_leu_18S_F01		CGATTCTTGCGTTCCATT	61.2	20	55
Dia_leu_18S_R01		GCCAAGAGGGTTATTGATCC	59.8	20	55
Dia_leu_18S_P01		CCCTCTCAGGGGTTCACCTCTATTTC	68.2	26	54

Related species	Forward	Reverse	Probe
<i>Actinauge richardi</i>	1	1	2
<i>Actinia equina</i>	4	3	3
<i>Actinia fragacea</i>	4	4	3
<i>Actinia tenebrosa</i>	4	3	2
<i>Actinostola crassicornis</i>	2	3	3
<i>Adamsia palliata</i>	1	4	3
<i>Alicia mirabilis</i>	0	1	1
<i>Alicia sansibarensis</i>	0	1	1
<i>Allantactis parasitica</i>	1	1	1
<i>Anemonia sulcata</i>	3	6	4
<i>Anemonia viridis</i>	2	6	3
<i>Antholoba achates</i>	4	5	9
<i>Anthopleura artemisia</i>	3	3	3
<i>Anthopleura atodai</i>	4	4	3
<i>Anthopleura elegantissima</i>	6	8	5
<i>Anthopleura handi</i>	5	5	3
<i>Anthopleura krebsi</i>	3	2	2
<i>Anthopleura kurogane</i>	4	5	3
<i>Anthopleura midori</i>	4	3	2
<i>Anthopleura rosea</i>	6	7	4
<i>Anthosactis janmayenii</i>	2	5	3
<i>Aulactinia incubans</i>	3	4	5
<i>Aulactinia stella</i>	3	4	5
<i>Aulactinia vancouverensis</i>	5	4	5
<i>Aulactinia veratra</i>	6	7	4
<i>Aulactinia verrucosa</i>	3	2	2
<i>Bunodactis reynaudi</i>	3	2	2
<i>Bunodosoma grandis</i>	3	4	3
<i>Calliactis japonica</i>	1	1	1
<i>Calliactis parasitica</i>	3	5	3
<i>Calliactis polypus</i>	1	2	1
<i>Calliactis tricolor</i>	1	2	1
<i>Charisea saxicola</i>	0	1	1
<i>Cricophorus nutrix</i>	3	4	4
<i>Cyananthea hourdezi</i>	0	4	1
<i>Diadumene cincta</i>	2	4	5
<i>Diadumene leucolena</i>	0	0	0
<i>Diadumene leucolena</i>	0	0	0
<i>Diadumene sp.</i>	0	1	2
<i>Epiactis australiensis</i>	10	4	8
<i>Epiactis georgiana</i>	3	2	3
<i>Epiactis handi</i>	3	2	3
<i>Epiactis japonica</i>	3	2	3
<i>Epiactis lisbethae</i>	3	2	3
<i>Epiactis prolifera</i>	3	2	3
<i>Epiactis ritteri</i>	3	2	3
<i>Galatheanthemum profundale</i>	1	1	1
<i>Gyractis sesere</i>	4	4	2
<i>Haliplanella lineata</i>	4	10	10
<i>Haliplanella lucia</i>	4	7	7
<i>Hormathia armata</i>	3	5	4
<i>Hormathia lacunifera</i>	3	5	4
<i>Hormathia pectinata</i>	1	1	2

<i>Hormosoma scotti</i>	1	5	3
<i>Isactinia olivacea</i>	6	7	4
<i>Kadosactis antarctica</i>	0	4	1
<i>Metridium senile</i>	0	1	1
<i>Nemanthus nitidus</i>	2	3	2
<i>Ostiactis pearseae</i>	0	2	1
<i>Oulacis muscosa</i>	5	2	2
<i>Paracondylactis bertwigi</i>	4	5	4
<i>Paranthus niveus</i>	0	1	1
<i>Stephanthus antarcticus</i>	3	2	2
<i>Stomphia didemona</i>	4	9	7
<i>Triactis producta</i>	0	2	1
<i>Urticina coriacea</i>	3	4	3
<i>Urticina crassicornis</i>	3	2	4
<i>Urticina fecunda</i>	3	2	2
<i>Urticina grebelnyi</i>	7	2	3

An additional system for detection of *Diadumene cincta* is also recommended, in case *Diadumene lineata* have too distant a molecular evolutionary relationship with *Diadumene leucolena* (Table S49.2):

- Dia_cin_18S_F02: 5'-ACTGACCGAGCTGTCTTCT-3',
- Dia_cin_18S_R02: 5'-GTCGCAAGTCCGCCGCACA-3',
- Dia_cin_18S_P02: 5'-FAM-AAAGACCTCGTGTGCTCTTGACTGAG-BHQ-1-3',

Table S49.2: Species specific primer/probe assay for *Diadumene cincta* with, target gene, product size, melting temperature, primer/probe length, GC ratio (%), and number of mismatches between primer and probe region in closely related non-target species.

Species	Gene	Product size	Temp	Length	GC
Diadumene cincta	18S	70 bp			
Dia_cin_18S_F02		ACTGACCGAGCTGTCTTCT	61.2	20	55
Dia_cin_18S_R02		GTCGCAAGTCCGCCGCACA	59.8	20	55
Dia_cin_18S_P02		AAAGACCTCGTGTGCTCTTGACTGAG	68.2	26	54

Related species	Forward	Reverse	Probe
<i>Actinangle richardi</i>	1	5	4
<i>Actinia equina</i>	4	3	4
<i>Actinia fragacea</i>	4	2	4
<i>Actinia tenebrosa</i>	4	3	3
<i>Actinostola crassicornis</i>	2	3	2
<i>Adamsia palliata</i>	3	4	4
<i>Alicia mirabilis</i>	2	5	4
<i>Alicia sansibarensis</i>	2	5	4
<i>Allantactis parasitica</i>	1	5	4
<i>Anemonia sulcata</i>	5	2	1
<i>Anemonia viridis</i>	4	4	2
<i>Antholoba achates</i>	6	1	7
<i>Anthopleura artemisia</i>	5	3	4
<i>Anthopleura atodai</i>	6	2	2
<i>Anthopleura elegantissima</i>	4	5	6
<i>Anthopleura handi</i>	5	3	2
<i>Anthopleura krebsi</i>	5	4	3
<i>Anthopleura kurogane</i>	6	3	2
<i>Anthopleura midori</i>	6	3	3
<i>Anthopleura rosea</i>	5	3	3
<i>Anthosactis janmayenii</i>	4	5	4

<i>Aulactinia incubans</i>	5	3	2
<i>Aulactinia stella</i>	5	3	2
<i>Aulactinia vancouverensis</i>	7	3	2
<i>Aulactinia veratra</i>	5	3	3
<i>Aulactinia verrucosa</i>	3	4	3
<i>Bunodactis reynaudi</i>	5	4	3
<i>Bunodosoma grandis</i>	5	2	4
<i>Calliactis japonica</i>	1	5	4
<i>Calliactis parasitica</i>	3	5	4
<i>Calliactis polypus</i>	1	4	4
<i>Calliactis tricolor</i>	1	4	4
<i>Charisea saxicola</i>	2	5	4
<i>Cricophorus nutrix</i>	3	2	5
<i>Cyananthea hourdezi</i>	2	6	4
<i>Diadumene cincta</i>	0	0	0
<i>Diadumene leucolena</i>	2	4	5
<i>Diadumene leucolena</i>	2	4	5
<i>Diadumene sp.</i>	2	3	5
<i>Epiactis australiensis</i>	12	5	10
<i>Epiactis georgiana</i>	5	4	3
<i>Epiactis handi</i>	5	4	2
<i>Epiactis japonica</i>	5	4	2
<i>Epiactis lisbethae</i>	5	4	2
<i>Epiactis prolifera</i>	5	4	2
<i>Epiactis ritteri</i>	5	4	2
<i>Galatbeanthemum profundale</i>	1	5	4
<i>Gyractis sesere</i>	4	2	3
<i>Haliplanella lineata</i>	4	6	6
<i>Haliplanella lucia</i>	4	3	3
<i>Hormathia armata</i>	1	3	3
<i>Hormathia lacunifera</i>	1	3	3
<i>Hormathia pectinata</i>	1	5	4
<i>Hormosoma scotti</i>	3	3	4
<i>Isactinia olivacea</i>	5	3	3
<i>Kadosactis antarctica</i>	2	6	4
<i>Metridium senile</i>	2	5	4
<i>Metridium senile</i>	2	5	4
<i>Nemanthus nitidus</i>	2	7	5
<i>Ostiactis pearseae</i>	2	4	4
<i>Oulactis muscosa</i>	7	4	3
<i>Paracondylactis hertwigi</i>	6	3	3
<i>Paranthus niveus</i>	2	5	4
<i>Stephanthus antarcticus</i>	5	4	3
<i>Stomphia didemon</i>	4	5	4
<i>Triactis producta</i>	2	4	4
<i>Urticina coriacea</i>	5	4	2
<i>Urticina crassicornis</i>	5	4	3
<i>Urticina fecunda</i>	5	4	3
<i>Urticina grebelnyi</i>	9	4	2

Species 50: Bonnemaisonia hamifera

Full name: *Bonnemaisonia hamifera* Hariot 1891

In the genus *Bonnemaisonia* there are only four species. To ensure better assay design sequences from available species within the family *Bonnemaisoniales* were also included. Among the sequenced genes for the four species in the genus *Bonnemaisonia* the *rbcL* gene showed the best potential for assay design (table S50):

- Bon_ham_rbcL_F01: 5'-CAATTACTAGATTACCTGGGCA-3'
- Bon_ham_rbcL_R01: 5'-CTTTACAAAGTCCCGACCT-3'
- Bon_ham_rbcL_P01: 5'-FAM- GGCTTAGAGTCTATGGTTATGGCACGA-BHQ-1-3'

The total length of the product was 153 bp. The length may become a problem due to DNA degradation and therefore the possibility to extract fragments of such lengths for amplification.

Table S50: Species specific primer/probe assay for *Bonnemaisonia hamifera* with, target gene, product size, melting temperature, primer/probe length, GC ratio (%), and number of mismatches between primer and probe region in closely related non-target species

Species	Gene	Product size	Temp	Length	GC
<i>Bonnemaisonia hamifera</i>	<i>rbcL</i>	153 bp			
Bon_ham_rbcL_F01		CAATTACTAGATTACCTGGGCA	58.4	22	41
Bon_ham_rbcL_R01		CTTTACAAAGTCCCGACCT	56.4	20	45
Bon_ham_rbcL_P01		GGCTTAGAGTCTATGGTTATGGCACGA	68.5	28	46

Related species*	Forward	Reverse	Probe
<i>Bonnemaisonia asparagooides</i>	5	3	6
<i>Bonnemaisonia clarata</i>	5	3	6
<i>Bonnemaisonia geniculata</i>	5	5	5
<i>Asparagopsis armata</i>	2	7	5
<i>Asparagopsis taxiformis</i>	6	6	6
<i>Delisea flaccida</i>	5	5	4
<i>Delisea hypneoides</i>	5	5	4
<i>Delisea japonica</i>	5	4	missing
<i>Delisea pulchra</i>	5	3	4
<i>Ptilonia magellanica</i>	6	2	5
<i>Reticulocalyx mucosissimus</i>	5	5	7

* not necessary related but show strong sequence similarity to target species

References

- Aagaard, A.B. (2015): Comparison between genotype and phenotype and a potential determination of origin of the invasive North American mud crab *Rhithropanopeus harrisii*. BSc thesis, University of Copenhagen. Unpublished.
- Batista, F., A. Leitao, A. Huvet, S. Lapegue, S. Heurtebise & P. Boudry (2005): The taxonomic status and origin of the Portuguese oyster *Crassostrea angulata* (Lamark, 1819). International Oyster Symposium. <http://archimer.ifremer.fr/doc/00000/3433/>
- Blank, M. & R. Bastrop (2009): Phylogeny of the mud worm genus *Marenzelleria* (Polychaeta, Spionidae) inferred from mitochondrial DNA sequences. *Zoologica Scripta* 38(3):313-321.
- Bellan, G. (2001): Polychaeta, in: Costello, M.J. *et al.* (Ed.) (2001). European register of marine species: a check-list of the marine species in Europe and a bibliography of guides to their identification. Collection Patrimoines Naturels 50:214-231.
- Coan, E. V. & P. Valentich-Scott (2012): Bivalve seashells of tropical West America. Marine bivalve mollusks from Baja California to northern Peru. 2 vols, 1258 pp.
- Consortium TU (2007): The Universal Protein Resource (UniProt). *Nucleic Acids Research* 35:D193-D197.
- Cosel, R. von, (2009): The razor shells of the eastern Atlantic, part 2. Pharidae II: the genus *Ensis* Schumacher, 1817 (Bivalvia, Solenoidea). *Basteria*, 73:9-56.
- Coyne, K.J., S.M. Handy, E. Demir, E.B. Wherat, D.A. Hutchins, K.J. Portune, M.A. Doblin, S.C. Cary (2005): Improved quantitative real-time PCR assays for enumeration of harmful algal species in field samples using an exogenous DNA reference standard. *Limnology and Oceanography: Methods* 3:381-391.
- Dittami, S., I. Riisberg, B. Edvardsen (2013): Molecular probes for the detection and identification of ichthyotoxic marine microalgae of the genus *Pseudochattonella* (Dictyochophyceae, Ochrophyta). *Environmental Science and Pollution Research* 20:6824-6837.
- Eckford-Soper, L.K. and N. Daugbjerg (in press): A quantitative real-time PCR assay for identification and enumeration of the occasionally co-occurring ichthyotoxic *Pseudochattonella farcimen* and *P. verruculosa* and analysis of variation in gene copy numbers during the growth phase of single and mixed cultures. *Journal of Phycology*.
- Froese, R. & D. Pauly. (Eds.) (2015): FishBase. World Wide Web electronic publication. www.fishbase.org, (10/2015). Accessed Nov-2015.
- Goldberg, C.S., A. Sepulveda, A. Ray, J. Baumgardt & L.P. Waits (2013): Environmental DNA as a new method for early detection of New Zealand mudsnails (*Potamopyrgus antipodarum*). *Freshwater Science* 32(3):792–800.
- Goldstien, S.J., L. Dupont, F. Viard, P.J. Hallas, T. Nishikawa, D.R. Schiel, N.J. Gemmell. & JD. Bishop (2011): Global Phylogeography of the Widely Introduced North West Pacific Ascidian *Styela clava*. *PLoS ONE* 6(2): e16755.
- Hall, T.A. (1999): BioEdit: a user-friendly biological sequence alignment editor and analysis program for Windows 95/98/NT. *Nucl. Acids. Symp. Ser.* 41:95-98.

Jane, S.F., T.M. Wilcox, K.S. Mckelvey, M.K. Young, M.K. Schwartz, W.H. Lowe, B.H. Letcher & A.R. Whiteley (2014): Distance, flow and PCR inhibition: EDNA dynamics in two headwater streams. Molecular Ecology Resources 15:216–227.

Kantor, Y.I., M.V. Vinarski, A.A. Schileyko & A.V. Sysoev (2010): "Catalogue of the Continental Mollusks of Russia and Adjacent Territories" (PDF). Ruthenica, Russian Malacological Journal (online version). ISSN 2307-7336.

Komai, R., I. Yamasaki, S. Kobayashi, T. Yamamoto, S. Watanabe (2006): *Eriocheir ogasawaraensis* Komai, a new species of mitten crab (Crustacea: Decapoda: Brachyura: Varunidae) from the Ogasawara islands, Japan, with notes on the systematics of *Eriocheir* De Haan, 1835. Zootaxa 1168:1-20

Monniot, C. (2001): Ascidiacea & Sorberacea, in: Costello, M.J., C. Emblow & R.J. White (Eds.) European register of marine species: a check-list of the marine species in Europe and a bibliography of guides to their identification. Collection Patrimoines Naturels 50:352-355.

Muus, B.J., J.G. Nielsen, P. Dahlstrøm & B.O. Nyström (2006): Havfisk og fiskeri. Gyldendal.

Nathan, L.M., M. Simmons, B.J. Wegleitner, C.L. Jerde & A.R. Mahon (2014): Quantifying environmental DNA signals for aquatic invasive species across multiple detection platforms. Environmental Science & Technology 48(21):12800–12806.

Nathan, L.R., C.L. Jerde, M.L. Budny & A.R. Mahon (2015): The use of environmental DNA in invasive species surveillance of the Great Lakes commercial bait trade. Conservation Biology 29(2):430-439.

Pérez-Portela, R., J.D. Bishop, A.R. Davis & X. Turon (2009): Phylogeny of the families Pyuridae and Styelidae (Stolidobranchiata, Ascidiacea) inferred from mitochondrial and nuclear DNA sequences. Molecular Phylogenetics and Evolution 50(3):560-570.

Stefaniak, L., G. Lambert, A. Gittenberger, H. Zhang, S. Lin & R.B. Whitlatch (2009): Genetic conspecificity of the worldwide populations of *Didemnum vexillum* Kott, 2002. Aquatic Invasions 4(1):29-44.

Stepien, C.A. & E.M. Neilson (2013): What's in a name? Taxonomy and nomenclature of invasive gobies in the Great Lakes and beyond. Journal of Great Lakes Research 39:555-559.

Scorzetti, G., L.E. Brand, G.L. Hitchcock, K.S. Rein, C.D. Sinigalliano, J.W. Fell (2009): Multiple simultaneous detection of Harmful Algal Blooms (HABs) through a high throughput bead array technology, with potential use in phytoplankton community analysis. Harmful Algae 8:196-211.

Scriver, M., A. Marinich, C. Wilson, J. Freeland (2015): Development of species-specific environmental DNA (eDNA) markers for invasive aquatic plants. Aquatic Botany 122:27-31.

Turner, C.R., M. A. Barnes, C.C.Y. Xu, S.E. Jones, C.L. Jerde & D.M. Lodge (2014): Particle size distribution and optimal capture of aqueous microbial eDNA. Methods in Ecology and Evolution 5:676–684.

Ten Hove H., A. Harry & E.K. Kupriyanova (2009): Taxonomy of Serpulidae (Annelida, Polychaeta): The state of affairs. Zootaxa 2036, 126 pp.

Türkay, M. (2001): Decapoda, in: Costello, M.J., C. Emblow & R.J. White (Eds.) European register of marine species: a check-list of the marine species in Europe and a bibliography of guides to their identification. Collection Patrimoines Naturels 50:284-292.

van der Land, J. & J.H. den Hartog (2001): Hexacorallia, Actiniaria, in: Costello, M.J., C. Emblow & R.J. White (Eds.) European register of marine species: a check-list of the marine species in Europe and a bibliography of guides to their identification. Patrimoines Naturels 50:214-231.

Vierna J., K.T. Jensen, A.M. González-Tizón & A. Martínez-Lage (2012) Population genetic analysis of *Ensis directus* unveils high genetic variation in the introduced range and reveals a new species from the NW Atlantic. *Marine Biology* 159(10):2209–2227.

Wang, H., L. Qian, A. Wang and X. Guo (2013): Occurrence and Distribution of *Crassostrea sikamea* (Amemiya 1928) in China. *Journal of Shellfish Research* 32:439-446.

Wang, F., E. He, Y. Li, X. Cai & W. Ma(2015): Complete mitochondrial genome of the hybridized fish (*Oncorhynchus mykiss* x Atlantic salmon). *Mitochondrial DNA* 29:1-2.

WoRMS Editorial Board (2015). World Register of Marine Species. Available from <http://www.marinespecies.org> at VLIZ. Accessed november 2015

Yamasaki, I., G. Yoshizaki, M. Yokota, C.A. Strüssmann & S. Watanabe(2006): Mitochondrial DNA variation and population structure of the Japanese mitten crab *Eriocheir japonica*. *Fisheries science* 72:299-309.

Yan, J., S. Liu, Y. Sun, C. Zhang, K. Luo & Y. Liu (2005): RAPD and microsatellite analysis of diploid gynogens from allotetraploid hybrids of red crucian carp (*Carassius auratus*) x common carp (*Cyprinus carpio*). *Aquaculture* 243(1-4):49-60.

Yuan, J, Mi T, Zhen Y, Yu Z (2012): Development of a rapid detection and quantification method of Karenia mikimotoi by real-time quantitative PCR. *Harmful Algae* 17:83-91.

Appendix D: Ranking of species

Nr.	Group	Name	DCE	HELCOM	OSPAR	EU-forordning	NST 1	NST 2	KJ 1	KJ 2	PRM	Criteria 1 established NIS	Criteria 2 potential NIS	Criteria 3 invasive species	Criteria 4 ease of deter- mination	Criteria 5 eDNA	Score
1	Fish	<i>Neogobius melanostomus</i> (Pallas, 1814)	x	x	x			x	x			10		6	3	4	23
2	Inverte-brate	<i>Crassostrea gigas</i> (Thunberg, 1793)	x	x	x	x	x		x			10		6	3	3	22
3	Inverte-brate	<i>Crepidula fornicata</i> (Linnaeus, 1758)	x	x	x	x		x	x			10		6	3	3	22
4	Inverte-brate	<i>Teredo navalis</i> (Linnaeus, 1758)	x				x		x			10		6	3	3	22
5	Phyto	<i>Karenia mikimotoi</i> (G. Hansen & Moestrup, 2000)	x	x	x			x				10		6	2	3	21
6	Fish	<i>Oncorhynchus mykiss</i> (Walbaum, 1792)	x					x				10		4	3	4	21
7	Zoo-plankton	<i>Mnemiopsis leidyi</i> (A. Agassiz, 1865)	x		x		x		x			10		6	3	2	21
8	Inverte-brate	<i>Eriocheir sinensis</i> (H. Milne Edwards, 1853)	x		x		x		x			10		4	3	3	20
9	Inverte-brate	<i>Rhithropanopeus harrisii</i> (Gould, 1841)	x	x	x			x				10		4	3	3	20
10	Phyto	<i>Prorocentrum minimum</i> (Pavillard) (Schiller, 1933)	x					x				10		4	2	3	19
11	Sub. Veg.	<i>Gracilaria vermiculophylla</i> (Ohmi) (Papenfuss, 1967)	x	x	x		x		x			10		6	2	1	19
12	Sub. Veg.	<i>Sargassum muticum</i> (Yendo) (Fensholt, 1955)	x				x	x	x			10		6	2	1	19
13	Inverte-brate	<i>Cordylophora caspia</i> (Pallas, 1771)	x					x				10		4	2	3	19
14	Inverte-brate	<i>Styela clava</i> (Herdman, 1881)	x	x				x				10		4	2	3	19
15	Inverte-brate	<i>Dreissena polymorpha</i> (Pallas, 1771)	x	x			x	x				8		6	2	3	19
16	Fish	<i>Acipenser baerii</i>						x	8					4	3	4	19
17	Fish	<i>Acipenser gueldenstaedtii</i>						x	8					4	3	4	19

Nr.	Group	Name	DCE	HELCOM	OSPAR	EU-forordning	NST 1	NST 2	KJ 1	KJ 2	PRM	Criteria	Criteria	Criteria	Criteria	Criteria	Score
												1 established NIS	2 potential NIS	3 invasive species	4 ease of determination	5 eDNA	
18	Fish	<i>Acipenser ruthenus</i>									x	8		4	3	4	19
19	Fish	<i>Acipenser stellatus</i>									x	8		4	3	4	19
20	Fish	<i>Huso huso</i>									x	8		4	3	4	19
21	Fish	<i>Oncorhynchus gorbuscha</i>									x	8		4	3	4	19
22	Fish	<i>Salvelinus fontinalis</i> (Mitchill, 1814)								x		8		4	3	4	19
23	Sub. Veg.	<i>Colpomenia peregrina</i> (Sauvageau, 1927)	x						x	x		10		4	3	1	18
24	Invertebrate	<i>Ensis Americanus</i> (Gould)	x	x	x		x		x			8		4	3	3	18
25	Sub. Veg.	<i>Dasya baillouviana</i> (Gmelin) (Montagne, 1841)	x				x	x				10		4	2	1	17
26	Sub. Veg.	<i>Heterosiphonia japonica</i> (Yendo, 1920)	x				x	x				10		4	2	1	17
27	Sub. Veg.	<i>Spartina angelica</i>	x				x	x				10		4	2	1	17
28	Phyto	<i>Heterosigma akashiwo</i>	x									8		4	2	3	17
29	Phyto	<i>Pseudochattonella farcimen</i> (W. Eichrem, 2009)					x					8		4	2	3	17
30	Invertebrate	<i>Molgula manhattensis</i> (de Kay, 1843)					x				x	8		4	2	3	17
31	Zoo-plankton	<i>Cercopagis pengoi</i> (Ostroumov, 1891)	x	x			x	x			x	0	6	6	3	2	17
32	Invertebrate	<i>Homarus americanus</i> (H. Milne Edwards, 1837)				x	x	x	x	x		0	5	6	3	3	17
33	Invertebrate	<i>Paralithodes camtschaticus</i> (Tilesius, 1815)			x		x	x	x	x		0	5	6	3	3	17
34	Invertebrate	<i>Didemnum vexillum</i> (Kott, 2002)	x	x	x		x		x	x		0	8	4	2	3	17
35	Sub. Veg.	<i>Fucus evanescens</i> C. (Agardh, 1820)	x				x		x			10		4	1	1	16
36	Invertebrate	<i>Petricolaria pholadiformis</i> (Lamarck, 1818)	x				x		x			10		0	3	3	16
-	Parasites	<i>Anguillicola crassus</i> Kuwahara, (Niimi & Itagaki, 1974)	x				x		x			10		4	1	1	16

Nr.	Group	Name	DCE	HELCOM	OSPAR	EU-forordning	NST 1	NST 2	KJ 1	KJ 2	PRM	Criteria 1 established NIS	Criteria 2 potential NIS	Criteria 3 invasive species	Criteria 4 ease of deter- mination	Criteria 5 eDNA	Score
37	Fish	<i>Percoccottus glenii</i>			x							3	6	3	4	16	
38	Invertebrate	<i>Elminius modestus</i> (Darwin, 1854)	x	x				x			10		0	2	3	15	
39	Invertebrate	<i>Ficopomatus enigmatis</i> (Fauvel, 1923)	x	x	x			x			10		0	2	3	15	
40	Invertebrate	<i>Marenzelleria viridis</i> (Verrill, 1873)	x	x				x			10		0	2	3	15	
41	Invertebrate	<i>Ocenebra inornata</i> (Récluz, 1851)	x					x			10		0	2	3	15	
42	Invertebrate	<i>Potamopyrgus antipodarum</i> (J.E. Gray, 1843)	x					x			10		0	2	3	15	
43	Phyto	<i>Pseudochattonella verriculosum</i>	x	x	x						10		0	2	3	15	
44	Sub. Veg.	<i>Codium fragile</i> subsp. <i>fragile</i> (Suringar) (Hariot, 1889)					x	x			8		4	2	1	15	
45	Fish	<i>Carassius auratus auratus</i> (Linnaeus, 1758)				x	x				8		0	3	4	15	
46	Fish	<i>Cyprinus carpio</i> (Linnaeus, 1758)					x	x			8		0	3	4	15	
47	Invertebrate	<i>Mya arenaria</i> (Linnaeus, 1758)	x				x				10		0	2	2	14	
48	Zoo-plankton	<i>Penilia avirostris</i> (Dana, 1849)	x				x				10		0	2	2	14	
49	Invertebrate	<i>Diadumene lineata</i> (Verrill, 1869)					x				8		0	3	3	14	
50	Sub. Veg.	<i>Bonnemaisonia hamifera</i> (Hariot, 1891)	x				x	x			10		0	2	1	13	
51	Phyto	<i>Alexandrium catenella</i>	x	x							8		0	2	3	13	
52	Phyto	<i>Alexandrium minutum</i> (Halim, 1960)					x				8		0	2	3	13	
53	Phyto	<i>Alexandrium tamarensense</i> (Lebour) Balech, 1995					x				8		0	2	3	13	
54	Phyto	<i>Coscinodiscus wailesii</i> (Gran & Angst, 1931)	x	x			x				8		0	2	3	13	
55	Phyto	<i>Gymnodinium chlorophorum</i>	x								8		0	2	3	13	

Nr.	Group	Name	DCE	HELCOM	OSPAR	EU-forordning	NST		KJ		PRM	Criteria 1		Criteria 2		Criteria 3		Criteria 4		Score	
							1	2	1	2		established NIS	potential NIS	invasive species	ease of determination	eDNA					
56	Phyto	Odontella sinensis (Greville) (Grunow, 1884)						x			x	8		0	2	3	13				
57	Invertebrate	Balamus impovisus (Darwin)	x									8		0	2	3	13				
58	Invertebrate	Caprella mutica (Schurin, 1935)		x	x	x			x		x	8		0	2	3	13				
59	Invertebrate	Cauilleriella killariensis (Southern, 1914)							x		x	8		0	2	3	13				
60	Invertebrate	Neanthes succinea (Frey & Lauckart)	x									8		0	2	3	13				
61	Invertebrate	Pachycordyle navis (Millard, 1959)							x		x	8		0	2	3	13				
62	Invertebrate	Platorchestia platensis (Krøyer, 1845)							x		x	8		0	2	3	13				
63	Invertebrate	Telmatogeton japonicus (Tokunaga, 1933)							x		x	8		0	2	3	13				
64	Invertebrate	Limulus polyphemus (Linnaeus, 1758)							x		x	0	3	4	3	3	13				
65	Invertebrate	Gammarus tigrinus (Sexton, 1939)	x	x					x		x	0	4	4	2	3	13				
66	Zoo-plankton	Acartia tonsa (Dana, 1849)	x		x				x		x	10		0		2	12				
67	Sub. Veg.	Neosiphonia harveyi (J.W. Bailey) (M.-S. Kim, H.-G. Choi, Guiry & G.W. Saunders, 2001)	x						x		x	10		0	1	1	12				
68	Parasites	Gonionemus vertens (A. Agassiz, 1862)	x						x		x	10		0	1	1	12				
69	Parasites	Pseudodactylogyrus anguillae (Yin & Sproston, 1948)	x					x		x	x	10		0	1	1	12				
70	Invertebrate	Rapana venosa (Valenciennes, 1846)		x	x	x		x		x	x	0	7	0	2	3	12				
71	Parasites	Pseudodactylogyrus bini (Kikuchi, 1929)	x						x		x	10		0		1	11				
72	Sub. Veg.	Aglaothamnion halliae (F.S. Collins) (N.E. Aponte, D.L. Ballantine & J.N. Norris, 1997)	x						x		x	8		0	2	1	11				

Nr.	Group	Name	DCE	HELCOM	OSPAR	EU-forordning	NST		KJ		PRM	Criteria 1		Criteria 2		Criteria 3		Criteria 4		Score	
							1	2	1	2		established NIS	potential NIS	invasive species	ease of determination	eDNA					
73	Sub. Veg.	<i>Dictuota dichotoma</i> (Hudson) (J.V. Lamoutoux)	x									8		0	2	1	11				
74	Sub. Veg.	<i>Porphyra umbilicalis</i> (Kützing, 1843)						x				8		0	2	1	11				
75	Invertebrate	<i>Callinectes sapidus</i> (Rathbun, 1896)	x	x			x		x			0	5	0	3	3	11				
76	Invertebrate	<i>Corbicula fluminea</i> (O.F. Müller, 1774)	x	x					x			0	5	0	3	3	11				
77	Invertebrate	<i>Hydroides dianthus</i>	x	x								0	6	0	2	3	11				
78	Invertebrate	<i>Mytilopsis leucophaeta</i> (Conrad, 1831)	x	x					x			0	5	0	3	3	11				
79	Sub. Veg.	<i>Codium fragile</i> subsp. <i>Tomentosoides</i> (van Goor) (P.C. Silva)	x									8		0	1	1	10				
80	Zoo-plankton	<i>Hemimysis anomala</i> (G.O. Sars, 1907)	x	x					x			0	6	0	2	2	10				
81	Invertebrate	<i>Hemigrapsus sanguinus</i> (de Haan, 1835)	x	x					x			0	5	0	2	3	10				
82	Invertebrate	<i>Hemigrapsus takanoi</i> (Asakura & Watanabe, 2005)	x	x					x			0	5	0	2	3	10				
83	Invertebrate	<i>Marenzelleria neglecta</i> (Sikorski & Bick, 2004)	x	x					x			0	5	0	2	3	10				
84	Parasites	<i>Aphanomyces astaci</i> Schikora, 1906					x		x			8		0		1	9				
85	Invertebrate	<i>Dikerogammarus villosus</i> (Sowinsky, 1894)	x	x					x			0	4	0	2	3	9				
86	Invertebrate	<i>Venerupis philippinarum</i> (A. Adams & Reeve, 1850)							x			0	3	0	3	3	9				
87	Fish	<i>Gambusia holbrooki</i> (Girard, 1859)						x				0	2	0	3	4	9				
88	Phyto	<i>Alexandrium leei</i> (Balech, 1985)						x				0	3	0	2	3	8				
89	Phyto	<i>Gymnodinium catenatum</i> (L.W. Graham, 1943)							x			0	3	0	2	3	8				

Nr.	Group	Name	DCE	HELCOM	OSPAR	EU-forordning	NST		KJ		PRM	Criteria 1		Criteria 2		Criteria 3		Criteria 4		Criteria 5	Score
							1	2	1	2		established NIS	potential NIS	invasive species	ease of determination	eDNA					
90	Zoo-plankton	<i>Beroe ovata</i> (Bruguière, 1789)						x				0	3	0	3	2	8				
91	Inverte-brate	<i>Asterias amurensis</i>	x	x								0	2	0	3	3	8				
92	Inverte-brate	<i>Chelicorophium curvispinum</i> (G.O: Sars, 1895)						x				0	3	0	2	3	8				
93	Inverte-brate	<i>Gonionemus vertens</i> (A. Agassiz, 1862)						x				0	2	0	3	3	8				
94	Inverte-brate	<i>Mercenaria mercenaria</i> (Linnaeus, 1758)						x				0	3	0	2	3	8				
95	Inverte-brate	<i>Potamothrix heuscheri</i> (Bretschler, 1900)						x				0	3	0	2	3	8				
96	Phyto	<i>Dinophysis sacculus</i>	x	x								0	2	0	2	3	7				
97	Phyto	<i>Fibrocapsa japonica</i>	x	x								0	2	0	2	3	7				
98	Phyto	<i>Pfiesteria piscicida</i>	x	x								0	2	0	2	3	7				
99	Zoo-plankton	<i>Evadne anomyx</i> (G.O. Sars, 1897)						x				0	3	0	2	2	7				
100	Zoo-plankton	<i>Limnomysis benedeni</i> (Czerniavsky, 1882)						x				0	3	0	2	2	7				
101	Inverte-brate	<i>Anadara transversa</i>		x								0	1	0	3	3	7				
102	Inverte-brate	<i>Arcuatula senhousia</i>	x	x								0	2	0	2	3	7				
103	Inverte-brate	<i>Botrylloides violaceus</i> (Oka, 1927)						x				0	3	0	1	3	7				
104	Inverte-brate	<i>Bugula neritina</i> (Linnaeus, 1758)						x				0	3	0	1	3	7				
105	Inverte-brate	<i>Chama pacifica</i>		x								0	1	0	3	3	7				
106	Inverte-brate	<i>Chionoecetes opilio</i>		x								0	1	0	3	3	7				
107	Inverte-brate	<i>Corella eumyota</i> (Traustedt, 1882)						x				0	2	0	2	3	7				
108	Inverte-brate	<i>Dikerogammarus haemobaphes</i> (Eichwald, 1841)						x				0	2	0	2	3	7				

Nr.	Group	Name	DCE	HELCOM	OSPAR	EU-forordning	NST		KJ		PRM	Criteria 1		Criteria 2		Criteria 3		Criteria 4		Criteria 5	Score
							1	2	1	2		established NIS	potential NIS	invasive species	ease of determination	eDNA					
109	Invertebrate	<i>Dreissena bugensis</i> (Andrusov, 1897)	x	x					x			0	2	0		2	3	7			
110	Invertebrate	<i>Gmelinoides fasciatus</i> (Stebbing, 1899)							x			0	2	0		2	3	7			
111	Invertebrate	<i>Monocorophium sextonae</i> (Crawford, 1937)							x			0	2	0		2	3	7			
112	Invertebrate	<i>Mytilus galloprovincialis</i>	x	x								0	2	0		2	3	7			
113	Invertebrate	<i>Palaemon macrodactylus</i>	x	x								0	2	0		2	3	7			
114	Invertebrate	<i>Perophora japonica</i> (Oka, 1927)							x			0	2	0		2	3	7			
115	Invertebrate	<i>Pontogammarus robustoides</i> (G.O. Sars, 1894)							x			0	2	0		2	3	7			
116	Invertebrate	<i>Potamocorbula amurensis</i>	x	x								0	2	0		2	3	7			
117	Invertebrate	<i>Rangia cuneata</i>	x	x								0	2	0		2	3	7			
118	Fish	<i>Neogobius fluviatilis</i> (Pallas, 1814)						x				0	0	0		3	4	7			
119	Fish	<i>Neogobius gymnotrachelus</i> (Kessler, 1857)						x				0	0	0		3	4	7			
120	Fish	<i>Neogobius kessleri</i> , (Günther, 1861)						x				0	0	0		3	4	7			
121	Fish	<i>Proterorhinus marmoratus</i> , (Pallas, 1814)						x				0	0	0		3	4	7			
122	Phyto	<i>Alexandrium monilatum</i>		x								0	1	0		2	3	6			
123	Phyto	<i>Alexandrium ostenfeldii</i>	x									0	1	0		2	3	6			
124	Phyto	<i>Phaeocystis pouchetii</i>		x								0	1	0		2	3	6			
125	Sub. Veg.	<i>Undaria pinnatifida</i> (Harvey) (Suringar, 1873)	x	x				x				0	3	0		2	1	6			
126	Invertebrate	<i>Amphibalanus eburneus</i> , ex. <i>Balanus eburnus</i>		x								0	1	0		2	3	6			
127	Invertebrate	<i>Brachidontes pharaonis</i>		x								0	1	0		2	3	6			

Nr.	Group	Name	DCE	HELCOM	OSPAR	EU-forordning	NST		KJ		PRM	Criteria 1		Criteria 2		Criteria 3		Criteria 4		Criteria 5	
							1	2	1	2		established NIS	potential NIS	invasive species	ease of determination	eDNA					
128	Invertebrate	<i>Hydroides elegans</i>		x								0	1	0	2	3	6				
129	Invertebrate	<i>Microcosmus squamiger</i>		x								0	1	0	2	3	6				
130	Invertebrate	<i>Nematostella vectensis</i> Stephenson, 1935					x					0	0	0	3	3	6				
131	Invertebrate	<i>Nemopsis bachei</i> L. Agassiz, 1849						x				0	0	0	3	3	6				
132	Invertebrate	<i>Palaemon elegans</i>	x									0	1	0	2	3	6				
133	Invertebrate	<i>Tricellaria inopinata</i> d'Hondt & Occhipinti Ambrogi, 1985					x					0	2	0	1	3	6				
134	Invertebrate	<i>Urosalpinx cinerea</i> (Say, 1822)						x				0	1	0	2	3	6				
135	Zoo-plankton	<i>Maeotias marginata</i> (Modeer, 1791)						x				0	0	0	3	2	5				
136	Zoo-plankton	<i>Mertensia ovum</i> (Fabricius, 1780)					x					0	0	0	3	2	5				
137	Sub. Veg.	<i>Caulerpa taxifolia</i>	x									0	1	0	3	1	5				
138	Sub. Veg.	<i>Grateloupia turuturu</i>	x	x								0	2	0	2	1	5				
139	Invertebrate	<i>Branchiura sowerbyi</i> Beddard, 1892					x					0	0	0	2	3	5				
140	Invertebrate	<i>Diadumene cincta</i> Stephenson, 1925					x					0	0	0	2	3	5				
141	Invertebrate	<i>Incisocalliope aestuarius</i> (Watling & Maurer, 1973)						x				0	0	0	2	3	5				
142	Invertebrate	<i>Marenzelleria arctica</i> (Chamberlin, 1920)						x				0	0	0	2	3	5				
143	Invertebrate	<i>Obesogammarus crassus</i> (G.O. Sars, 1894)						x				0	0	0	2	3	5				
144	Invertebrate	<i>Paranais frici</i> Hrabe, 1941					x					0	0	0	2	3	5				
145	Invertebrate	<i>Potamoithrix bavaricus</i> (Oschmann, 1913)						x				0	0	0	2	3	5				

Nr.	Group	Name	DCE	HELCOM	OSPAR	EU-forordning	NST 1	NST 2	KJ 1	KJ 2	PRM	Criteria	Criteria	Criteria	Criteria	Criteria	Score
												1 established NIS	2 potential NIS	3 invasive species	4 ease of determination	5 eDNA	
146	Invertebrate	Potamothis bedoti (Piguet, 1913)						x				0	0	0	2	3	5
147	Invertebrate	Potamothis moldaviensis (Vejdovsky & Mrázek, 1903)							x			0	0	0	2	3	5
148	Invertebrate	Potamothis vejvodskyi (Hrabe, 1941)							x			0	0	0	2	3	5
149	Sub. Veg.	Gratelouphia doryphore					x					0	1	0	2	1	4
150	Sub. Veg.	Halophila stipulacea					x					0	1	0	2	1	4
151	Sub. Veg.	Lophocladia lallemandii					x					0	1	0	2	1	4
152	Sub. Veg.	Stypospodium schimperi					x					0	1	0	2	1	4
153	Invertebrate	Ameira divagans (Nicholls, 1939)						x				0	0	0	1	3	4
154	Invertebrate	Garveia franciscana (Torrey, 1902)						x				0	0	0	1	3	4
155	Invertebrate	Jaera (Jaera) sarsi (Vakkanov, 1936)						x				0	0	0	1	3	4
156	Invertebrate	Victorella pavida (Saville Kent, 1870)						x				0	0	0	1	3	4
157	Zoo-plankton	Paramysis (Mesomysis) intermedia (Czerniavsky, 1882)						x				0	0	0	1	2	3
158	Zoo-plankton	Paramysis (Serrapalpisis) lacustris (Czerniavsky, 1882)						x				0	0	0	1	2	3
159	Sub. Veg.	Caulerpa cylindracea					x					0	1	0	1	1	3
160	Parasites	Bonamia ostreae Pichot (Comps, Tigé, Grizel & Rabouin, 1980)						x				0	1	0	1	1	3
161	Parasites	Marteilia refringens (Grizel, Comps, Bonami, Cousserans, Duthoit & Le Pennec, 1974)						x				0	1	0	1	1	3
162	Parasites	Myicola ostreeae (Hoshina & Sugiura, 1953)						x				0	1	0	1	1	3
163	Parasites	Mytilicola orientalis (Mori, 1935)						x				0	1	0	1	1	3
164	Parasites	Pseudobacciger harengulae (Yamaguti, 1938)						x				0	1	0	1	1	3

NIVA Denmark is the name, water is our game

NIVA Denmark Water Research is a regional office of the Norwegian Institute for Water Research (NIVA) and has just recently been established to resolve environmental issues concerning the freshwater and marine systems that relate to Denmark.

NIVA Denmark has primary focus on research-based implementation of a number of EU's directives *inter alia* the Water Framework Directive and the Marine Strategy Framework Directive together with international conventions (HELCOM, OSPAR, BDC). We occasionally provide consultancy to authorities and small and medium-sized companies.

NIVA Denmark is a place for practice, observation, testing and synthesis. Key research and test areas include eutrophication, hazardous substances, biodiversity, and ecosystem health as well as the implications of multiple human activities in marine waters and in streams, rivers and lakes. We develop indicators, monitoring methods and tools to assess the state of an ecosystem in order to carry out analyses and contribute to evidence based and sustainable solutions to the challenges we and the environment face.

NIVA Denmark, as a regional office to NIVA has thus the backing of more than 200 dedicated researchers and experts.



Winghouse
Ørestads Boulevard 73
2300 Copenhagen S
Denmark
Telephone: +45 88 96 96 70
E-mail: post@niva-denmark.dk
CVR/VAT no.: 35431063
www.niva-denmark.dk