

Green Ocean Amazon 2014/15 High-Volume Filter Sampling: Atmospheric Particulate Matter of an Amazon Tropical City and its Relationship to Population Health Field Campaign Report

CMD Machado
EO Santos
KS Fernandes

JLA Neto
RAFS Souza

August 2016



DISCLAIMER

This report was prepared as an account of work sponsored by the U.S. Government. Neither the United States nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the U.S. Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the U.S. Government or any agency thereof.

**Green Ocean Amazon 2014/15 High-Volume
Filter Sampling: Atmospheric Particulate
Matter of an Amazon Tropical City and its
Relationship to Population Health
Field Campaign Report**

CMD Machado, Federal University of Amazonas
Principal Investigator

EO Santos, Federal University of Amazonas
KS Fernandes, Federal University of Amazonas
JLA Neto, Federal University of Amazonas
RAFS Souza, University of the State of Amazonas
Co-Investigators

August 2016

Work supported by the U.S. Department of Energy,
Office of Science, Office of Biological and Environmental Research

Acknowledgments

We acknowledge the support from the Central Office of the Large Scale Biosphere Atmosphere Experiment in Amazonia (LBA), the Instituto Nacional de Pesquisas da Amazonia (INPA), and the Universidade do Estado do Amazonia (UEA). The work was conducted under 001030/2012-4 of the Brazilian National Council for Scientific and Technological Development (CNPq).

Acronyms and Abbreviations

ARM	Atmospheric Radiation Measurement Climate Research Facility
CONAMA	Brazilian National Council of the Environment
CNPq	Brazilian National Council for Scientific and Technological Development
DOE	U.S. Department of Energy
EPA	Environmental Protection Agency
GoAmazon	Green Ocean Amazon 2014/15 field campaign
INPA	Instituto Nacional de Pesquisas da Amazonia
LBA	Central Office of the Large Scale Biosphere Atmosphere Experiment in Amazonia
PAH	polycyclic aromatic hydrocarbon
PM	particulate matter
TSP	total suspended particulate
UEA	Universidade do Estado do Amazonia
UFAM	Federal University of Amazonas
WHO	World Health Organization

Contents

Acknowledgments.....	iii
Acronyms and Abbreviations	iv
1.0 Summary.....	1
2.0 Results	2
3.0 Publications and References.....	3

Figures

1 High-volume samplers (PM _{2.5} and TSP) at the ARM site in Manacapuru, Amazonas.....	1
2 PM _{2.5} and TSP levels obtained at the ARM GoAmazon campaign site in Manacapuru from March to December 2015.....	2

1.0 Summary

Manaus, the capital of the Brazilian state of Amazonas, is developing very rapidly. Its pollution plume contains aerosols from fossil fuel combustion mainly due to vehicular emission, industrial activity, and a thermal power plant. Soil resuspension is probably a secondary source of atmospheric particles. The plume transports from Manaus to the U.S. Department of Energy (DOE) Atmospheric Radiation Measurement (ARM) Climate Research Facility ARM site at Manacapuru urban pollutants as well as pollutants from pottery factories along the route of the plume. Considering the effects of particulate matter on health, atmospheric particulate matter was evaluated at this site as part of the ARM Facility's Green Ocean Amazon 2014/15 (GoAmazon 2014/15) field campaign.

Aerosol or particulate matter (PM) is typically defined by size, with the smaller particles having more health impact. Total suspended particulate (TSP) are particles smaller than 100 μm ; particles smaller than 2.5 μm are called $\text{PM}_{2.5}$.

In this work, the $\text{PM}_{2.5}$ levels were obtained from March to December of 2015, totaling 34 samples and TSP levels from October to December of 2015, totaling 17 samples. Sampling was conducted with $\text{PM}_{2.5}$ and TSP high-volume samplers using quartz filters (Figure 1). Filters were stored during 24 hours in a room with temperature (21,1°C) and humidity (44,3 %) control, in order to do gravimetric analyses by weighing before and after sampling. This procedure followed the recommendations of the Brazilian Association for Technical Standards local norm (NBR 9547:1997). Mass concentrations of particulate matter were obtained from the ratio between the weighted sample and the volume of air collected. Defining a relationship between particulate matter ($\text{PM}_{2.5}$ and TSP) and respiratory diseases of the local population is an important goal of this project, since no information exists on that topic.



Figure 1: High-volume samplers ($\text{PM}_{2.5}$ and TSP) at the ARM site in Manacapuru, Amazonas.

Further chemical analyses (metals and polycyclic aromatic hydrocarbon [PAH]) will be done on these samples in order to identify major pollutants and to investigate the sources of these particles.

This project was supported by the National Council for Scientific and Technological Development (CNPq), whose funds were used to purchase equipment, and the Amazonas Research Foundation, which supported GOAmazon Campaign Science and students' fellowships. The campaign was carried out by

graduate and post-graduate students of the Chemistry Department of the Federal University of Amazonas (UFAM). We also had transport logistic support from UFAM throughout the sampling period at the ARM site.

This work contributed to the technical and scientific development of the Amazon region, and to the development of support for quality of life and environmental preservation.

2.0 Results

The results presented here are concerned with levels of particulate matter PM_{2.5} and TSP collected during the ARM GoAmazon field campaign. The range of PM_{2.5} concentration, 4,25 – 71,18 µg m⁻³, was higher in 70 % of samples than the World Health Organization (WHO) recommendation (25 µg m⁻³ 24-hour mean). Considering limits stated by the U.S. Environmental Protection Agency (EPA), PM_{2.5} levels were higher in 52% of samples. TSP levels varied from 22,69 to 92,00 µg m⁻³ during the period. Based on the Brazilian regulatory standard from the National Council of Environment (CONAMA), these concentrations were 82% lower than limits expected for TSP (Figure 2). In comparison to mean concentration obtained in the Manaus urban area, PM_{2.5} and TSP mean levels were 120% and 59% higher, respectively.




Figure 2: PM_{2.5} and TSP levels obtained at the ARM GoAmazon campaign site in Manacapuru from March to December 2015. WHO and EPA limits of PM_{2.5} concentration and the CONAMA limit of TSP level are shown for comparison.

Some of the samples showed PM_{2.5} levels similar to those found in São Paulo, a city very different than Manacapuru in respect to population and source of particulate matter. It became clear that seasonality influences particle concentration, once the rainy season exhibited low levels of PM_{2.5}, while during the dry season concentrations were about 3-5 times higher. The effects of El Niño contributed to an increase in vegetation fires (up to 2.000 from July through December) and to prolonging the dry season until December. It should be highlighted that in October the temperature inversion contributed to stagnation of the atmosphere, preventing the dispersion of pollutants.

Correlations between precipitation/vegetation fires and particulate matter have not been significant, indicating that others factors like continuous local sources and a low dispersion of pollutants during the dry season could have a great influence on particle levels.

Given the damage to human health caused by $PM_{2.5}$, especially in the elderly, children, and people with chronic respiratory diseases, we evaluated the relationship between the number of admissions due to respiratory ailments and the concentrations of $MP_{2.5}$. The results showed no significant correlations, suggesting that other factors may be causing these human health problems.

For now, the results are insufficient to establish a clear relationship between sources of particles and the atmospheric pollution at the Manacapuru ARM site, or to identify the major influences in its concentration. This is also true in relation to the effects of $PM_{2.5}$ on the health of the local population. In consequence, we will continue this project in the Manaus urban area to acquire data and learn more about these issues.

The opportunity provided by the GoAmazon campaign was very important to start our work in this initial stage of atmospheric research in the Amazon region. We expect to collaborate with the ARM team again.

3.0 Publications and References

ABNT. 1997. Material Particulado em Suspensão no Ar Ambiente - Determinação da Concentração Total pelo Método do Amostrador de Grande Volume. NBR 9547, Rio de Janeiro, p. 14.

CONAMA. Controle da poluição do ar - PRONAR.
<http://www.mma.gov.br/port/conama/res/res90/res0390.html>.

U.S. Environmental Protection Agency. 2014. *National Ambient Air Quality Standards (NAAQS)*.
<https://www.epa.gov/criteria-air-pollutants/naaqs-table>.

World Health Organization. 2014. *Ambient (Outdoor) Air Quality and Health*.
<http://www.who.int/mediacentre/factsheets/fs313/en/#>.

NETO, JLA. [AMB047]. Poster presentation at 39th Annual Meeting of the Brazilian Chemical Society, in June, 2016. <http://www.s bq.org.br/39ra/cdrom/indiceautor.htm>.



U.S. DEPARTMENT OF
ENERGY

Office of Science